


# **FIRST PHYSICS FROM COMPASS**


**Franco Bradamante**

*University of Trieste and INFN Trieste*

on behalf of the  
**COMPASS Collaboration**


Bloomington Indiana, May 28





- **INTRODUCTION**
- **SPECTROMETER AND DATA TAKING**
- **FIRST PHYSICS RESULTS (run 2002)**
  - $A_1^d$
  - **Transversity**
  - $A_{LL}$  from high  $p_T$
- **OTHER ONGOING ANALYSIS**
  - $A_{LL}$  from open charm
  - $\Lambda$  physics
  - **Exclusive  $\rho$  and  $\phi$ ,  $J/\Psi$**
  - **Flavour separation**
- **SUMMARY AND OUTLOOK**

**COmmon  
Muon and  
Proton  
Apparatus for  
Structure and  
Spectroscopy**




**NA58**

Finland, France, Germany, India, Israel, Italy, Japan,  
Poland, Portugal, Russia, Switzerland

Bielefeld, Bochum, Bonn, Burdwan, Calcutta, CERN,  
Dubna, Erlangen, Freiburg, Heidelberg, Helsinki, Lisbon,  
Mainz, Miyazaky, Moscow, Munich, Nagoya, Protvino,  
Saclay, Tel Aviv, Torino, Trieste, Warsaw

**28 Institutes, more than 200 physicists**

# THE COMPASS EXPERIMENT



- **experiment:** **thought of in** **April '94** **Trento workshop**  
**Nov. '94** **Trieste workshop**  
**Lol** **March '95**  
**encouraged** **June '95** **SPSLC in Cogne**  
**Proposal** **March '96**  
**recommended** **Sept. '96**  
**approved by RB** **Feb. '97** **as NA58**  
**Technical run** **2000**  
**Commissioning** **2001**


- **since 2002 taking data**

**with**

**a new spectrometer with outstanding performances**

- **merging of two programmes:** **HMC** **CHEOPS**  
**(muon beam)** **(hadron beam)**



# COMPASS programme with the muon beam




to determine the polarised **parton** density functions in a **polarised nucleon** from measurements of **hadron asymmetries** in semi-inclusive polarised DIS, **both longitudinal and transverse**

specifically,

- to measure the gluon polarisation  $\Delta G$  through open charm (Gluk and Reya, Altarelli and Stirling, 1988)



- to measure  $h_1$ , the new territory
- to measure the spin transfer in fragmentation from  $\Lambda$  production
- to remeasure with high statistics  $g_1$  and  $g_2$
- .....



## ▪ charmed hadrons

- production phenomena ( $p$ ,  $\pi$ ,  $K$ )
- leptonic decays
- semileptonic decays
- precision measurements of **c-baryon lifetimes**
- production and spectroscopy of **cc-baryons**



## ▪ gluonic states

- search for **glueballs** in Pomeron-Pomeron scattering
- search for **exotic states**

## ▪ hadron structure



- **polarizability** in Primakoff reactions

# THE COMPASS SPECTROMETER





# THE COMPASS SPECTROMETER

## 2002







# THE POLARISED TARGET




- ${}^6\text{LiD}$
- $\pm 50\%$  polarisation
- 40 % dilution factor
- 2.5 T
- 50 mK

$\mu$

# THE TARGET SYSTEM



# THE COMPASS/Oxford Danfysik MAGNET



## MoU CERN/DAPNIA/COMPASS

signed december 2003 :


- finalize instrumentation, full magnetic tests

2004:

- **delivery to Saclay**
- vacuum tests, cooling, max field, reversal/homogeneity
- **delivery to CERN**




2005: **installation, tests, magnetic measurements, polarization**





# RICH1

## Ring Imaging Cherenkov


- 90 m<sup>3</sup> (3 m C<sub>4</sub>F<sub>10</sub>)
- 116 VUV mirrors (focal length 3.3 m)
- 5.3 m<sup>2</sup> UV detectors
  - MWPC CsI cathods
  - 8x8 mm<sup>2</sup> pad
- 84k analog r/o channels
- K/π separation up to ~40 GeV



# RICH1




**116mirrors, 20 m<sup>2</sup>**




**PDs, 5.3 m<sup>2</sup>**

# MicroMegas


## Micro Mesh Gas Detectors



### Novel gaseous detector




**40x40 cm<sup>2</sup>**






# MicroMegas


time resolution below 10 ns



spatial resolution below 70  $\mu\text{m}$




efficiency larger than 97%



# 2002 RUN

---




- **160 GeV/c muons,  $2 \cdot 10^8 \mu^+ / 4 \text{ s}$  every 16.8 s,  $P_{\text{beam}} \cong 80 \%$**
- **$^6\text{LiD}$  target,  $P_{\text{target}} \cong 50 \%$**
- **polarization reversal by magnet field rotation every 8 h**
  
- **200 k readout channels, 35-40 kB/event**
- **data taking:**
  - **24 days setup (about 2/3 of equipment new)**
  - **57 days longitudinal target polarisation**
  - **19 days transverse target polarisation**
  
- **5 billion events recorded, 260 TByte total**
  
- **similar statistics in 2003**



# Central Data Recording




**Design value: 35MB/s**  
**3TB/day**




**2002**

**260 TByte in ~100 days**  
**5 billion events**




**2002**

# DATA PROCESSING at CERN




500 TB of raw data  
→ 500 000 jobs batch  
( 1 GByte ~ 8 h )

Processing in parallel  
as much as possible




# FIRST PHYSICS RESULTS

---




- $A_1^D$
- $A_{LL}$  from *high*  $p_T$
- Transversity



# DOUBLE SPIN ASYMMETRY $A_1^d$

virtual photon-deuteron asymmetry




$$A_1 = \frac{\sigma_{1/2} - \sigma_{3/2}}{\sigma_{1/2} + \sigma_{3/2}} \approx \frac{\sum_a e_a^2 \cdot (\mathbf{q}_a^+ - \mathbf{q}_a^-)}{\sum_a e_a^2 \cdot (\mathbf{q}_a^+ + \mathbf{q}_a^-)}$$

$$g_1 = \frac{1}{2} \cdot \sum_a e_a^2 \cdot (\mathbf{q}_a^+ - \mathbf{q}_a^-)$$

$$F_1 = \frac{1}{2} \cdot \sum_a e_a^2 \cdot (\mathbf{q}_a^+ + \mathbf{q}_a^-)$$

$A_1 \approx \frac{g_1}{F_1}$

# DOUBLE SPIN ASYMMETRY $A_1^d$



**COMPASS:**

**2002 data only**


6.5 Million DIS events


$Q^2 > 1 \text{ (GeV/c)}^2$

$0.1 < y < 0.9$

**expect x4 statistics  
by end of 2004**


Data displayed at experimental  $\langle Q^2 \rangle$  of every  $x_{Bj}$  bin





# $\Delta G/G$ at COMPASS

## Photon Gluon Fusion




$q = c$  cross section difference  
in charmed meson production  
→ *theory well understood*  
→ *experiment challenging*

$q = u, d, s$  cross section difference  
in 2+1 jet production  
in COMPASS: events with  
2 hadrons with high  $p_T$   
→ *experiment easy*  
→ *theory difficult*





# $\Delta G/G$ : pairs of high $p_T$ hadrons

## Photon Gluon Fusion



- **Current fragmentation**
  - $x_F > 0.1$
  - $z > 0.1$
- **2 high  $p_T$  hadrons**
  - $p_T > 0.7 \text{ GeV}/c$
  - $p_{T1}^2 + p_{T2}^2 > 2.5 (\text{GeV}/c)^2$
  - $m(h_1 h_2) > 1.5 \text{ GeV}/c^2$






# $\Delta G/G$ : pairs of high $p_T$ hadrons

$$A^{\gamma^*d} = \frac{1}{2P_T f DP_B} \left[ \frac{N_1^{\leftarrow} - N_2^{\leftarrow}}{N_1^{\rightarrow} + N_2^{\leftarrow}} + \frac{N_2^{\leftarrow} - N_1^{\leftarrow}}{N_2^{\rightarrow} + N_1^{\leftarrow}} \right]$$


$$\hat{a}_{LL}^{PGF} \approx -1 \text{ and } \hat{a}_{LL}^{Com} \approx 0.5$$

fractions of cross section determined by Monte Carlo


$$A^{\gamma^*d} = \frac{A_{LL}^{\mu N \rightarrow hh}}{D} \approx \left\langle \frac{\hat{a}_{LL}^{PGF}}{D} \right\rangle \left\langle \frac{\Delta G}{G} \right\rangle \frac{\sigma^{PGF}}{\sigma^{tot}} + \left\langle \frac{\hat{a}_{LL}^{Com}}{D} \right\rangle \left\langle \frac{\Delta q}{q} \right\rangle \frac{\sigma^{Com}}{\sigma^{tot}} + LODIS$$



Photon Gluon Fusion




QCD-Compton



Leading Order





# $\Delta G/G$ : pairs of high $p_T$ hadrons

Asymmetry in production of hadron pairs with high  $p_T$ :  
preliminary result from 2002 data

$$A^{\gamma^*d} = -0.065 \pm 0.036_{stat.} \pm 0.010_{syst.}$$

up to now systematic error contains only studies on  
false asymmetries due to target or spectrometer effects

assuming  $R_{PGF} \sim 1/4$      $\sigma(\Delta G/G) \sim 0.17$

... and  $\Delta G/G > 0$


## Improvements:

- 2003 and expected 2004 data will give a factor of 4 more data
- better reconstruction algorithm, ...  $\sigma(\Delta G/G) \rightarrow 0.05$



# TRANSVERSE SPIN PHYSICS


3 structure functions are necessary to describe the spin structure of the nucleon at LO:



all of equal importance!

$h_1(x)$  decouples from leading twist DIS because helicity of quark must flip



**NO MIXTURE WITH GLUON**



# COLLINS ASIMMETRY



$$\Phi_C = \Phi_h - \Phi_{s'}$$




$$N_h^\pm(\Phi_C) = N_h^0 \cdot \left\{ 1 \pm \mathbf{A}_1^h \cdot \sin\Phi_C \right\}$$

$$\mathbf{A}_{\text{Coll}} = \frac{1}{\mathbf{f} \cdot \mathbf{P}_T \cdot \mathbf{D}_{\text{nn}}} \cdot \mathbf{A}_1^h = \frac{\sum_a e_a^2 \cdot \Delta_T q_a \cdot \Delta D_a^h}{\sum_a e_a^2 \cdot q_a \cdot D_a^h}$$


# COLLINS ASYMMETRY



## MC vs DATA



# COLLINS ASYMMETRY





## RESULTS

$A_{\text{coll}}$  depends on  $p_{hT}$ ,  $z_h$ ,  $x_{Bj}$


with more statistics, the full analysis is foreseen

from 2002 data:

### $A_{\text{coll}}$ VS $x_{Bj}$





# COLLINS ASYMMETRY



## RESULTS


from 2002 data:

### $A_{\text{Coll}}$ vs $z_h$



all the tests we made are consistent with the fact that

**systematic effects, if present, are smaller than statistical errors**





# ONGOING ANALYSIS

---

- $\Lambda$  and  $\bar{\Lambda}$  hyperon production
- Vector meson production  $\rho$ ,  $\phi$  and  $J/\psi$
- Flavour decomposition of polarized PDF
- $\Delta G/G$  from open charm

# $\Lambda$ PRODUCTION




**Armenteros-Podolanski**

$$\alpha = \frac{P_L^+ - P_L^-}{P_L^+ + P_L^-}$$




# $\Lambda$ PRODUCTION

## DATA vs MONTECARLO





Data

Monte Carlo



# $\Lambda$ POLARIZATION ?



1/6 of 2002  
statistics

$$Q^2 > 1 \text{ GeV}^2$$


$$0.2 < y < 0.9$$

good potential  
for polarization  
measurement

# EXCLUSIVE $\rho$ and $\phi$ PRODUCTION




meson	mass cut	statistics (1/6 of 2002)
$\rho^0$	$0.5 < m_{\pi\pi} < 1 \text{ GeV}$	$1.3 \cdot 10^6$
$\phi$	$ m_{KK} - m_\phi  < 9 \text{ MeV}$	$42 \cdot 10^3$




- $|t'| < 0.5 \text{ GeV}^2$
- $7.5 < W < 16 \text{ GeV}$
- $Q^2 > 10^{-3} \text{ GeV}^2$

# EXCLUSIVE $\rho$ and $\phi$ PRODUCTION





## INVARIANT MASSES




- 16 % of total 2002 statistics
- no MC corrections yet

# EXCLUSIVE $\rho$ and $\phi$ PRODUCTION

## INTERFERENCE of $\rho^0$ and $\pi\pi$





- Söding parametrization
- No accept. corr.




# EXCLUSIVE $\rho$ and $\phi$ PRODUCTION

## ANGULAR DISTRIBUTIONS



$$p_T > 0.15 \text{ GeV}$$


$$Q^2 > 0.05 \text{ GeV}^2$$



# J/ψ PRODUCTION


$$J/\psi \rightarrow \mu^+ \mu^-$$

- first look
- mainly elastic





# FLAVOUR SEPARATION $\Delta q$



Looks very promising in particular for  $\Delta s$  !

Can the first moment of  $\Delta s$  be **positive**?

Low-x data **essential**!







# $\Delta G/G$ : OPEN CHARM

$$A_{\gamma N}^{c\bar{c}} = \frac{\Delta\sigma^{\gamma N \rightarrow c\bar{c}X}}{\sigma^{\gamma N \rightarrow c\bar{c}X}} = \frac{\int d\hat{s} \Delta\sigma^{\text{PGF}}(\hat{s}) \Delta G(x_G, \hat{s})}{\int d\hat{s} \sigma^{\text{PGF}}(\hat{s}) G(x_G, \hat{s})} \approx \langle a_{LL} \rangle \left\langle \frac{\Delta G}{G} \right\rangle$$

$$\hat{s} = M_{c\bar{c}}^2$$

$\Delta\sigma^{\text{PGF}}$  at NLO: Bojak, Stratmann NPB 540 (1999) 345; Contogouris *et al.*

## Photon-Gluon Fusion



$$A_{\text{raw}} = \frac{N_{c\bar{c}}^{\rightarrow\leftarrow} - N_{c\bar{c}}^{\leftarrow\rightarrow}}{N_{c\bar{c}}^{\rightarrow\leftarrow} + N_{c\bar{c}}^{\leftarrow\rightarrow}} = P_\mu P_T f D A_{\gamma N}^{c\bar{c}}$$

$$P_\mu \approx -0.76$$

$$f \approx 0.4$$

$$P_T \approx 0.5$$



$$D(y)$$

# D\* tagging: $D^* \rightarrow D^0 \pi$



$$D^* \rightarrow (K\pi)\pi$$

$$D^* \rightarrow (K\pi)\pi$$






# D\* tagging: $D^* \rightarrow D^0 \pi$



$$D^* \rightarrow (K\pi)\pi$$

$$D^0 \rightarrow K\pi$$





# PROJECTIONS FOR $\sigma(\Delta G/G)$

With


- *improved FOM and*
  - *assuming ~ 80 scheduled days (only L data) in 2004*
- we expect **for all the deuteron data (2002, 2003, 2004)**

**$\sigma(\Delta G/G) = 0.24$  from open charm**

**$\sigma(\Delta G/G) = 0.05$  from high  $p_T$  all  $Q^2$**

**$\sigma(\Delta G/G) = 0.16$  from high  $p_T$   $Q^2 > 1$**

resolved  $\gamma$   
 $\sigma_{\text{syst}} ?$



# SUMMARY AND OUTLOOK

---

- CERN is again contributing to the **NUCLEON SPIN PUZZLE**
- a technically challenging new experiment is **IN OPERATION SINCE 2002**
  - “LHC” technologies     *detectors*  
   *read-out*  
   *data handling*
- a privileged situation at CERN
- **FIRST PHYSICS RESULTS** have been produced  
**MANY MORE IN THE PIPE-LINE**
- **COMPASS** is foreseen to run up to the end of the present mid-term plan of CERN (2010)

**BIG DISCOVERY POTENTIAL**

thank you

and  
see you all at  
spin2004

F. Bradamante, May 28



organizing committee

- |                       |                            |
|-----------------------|----------------------------|
| g. altarelli          | bern, switzerland          |
| m. anselmino          | torino, italy              |
| d. barbari            | humburg, germany           |
| r. bertini            | turin, italy               |
| f. bradamante         | trieste, italy (host)      |
| s. dalla torre        | trieste, italy             |
| p.f. dalpiaz          | ferenza, italy             |
| e. de sanctis         | innsbruck, austria         |
| m. figner             | prague, czech rep.         |
| t. johansson          | uppsala, sweden            |
| e. leader             | oxford, england            |
| d. peullier           | strasbourg, france         |
| a. martin             | trieste, italy (cosmology) |
| w. meyer              | böblingen, germany         |
| s. noruchev           | pryazino, russia           |
| r. payer              | trieste, italy             |
| a. penzo              | trieste, italy             |
| a. cipriakian         | duisburg, germany          |
| j. soifer             | marseille, france          |
| a. van der steenhoven | nijmegen, netherlands      |
| t. walcher            | mainz, germany             |

international committee

- |                 |                            |
|-----------------|----------------------------|
| t. rose         | brn, usa (inv)             |
| a.d. kriech     | michigan, usa (inv, chair) |
| f. bradamante   | trieste, italy             |
| o. chamberlain* | berkeley, usa              |
| d.g. crabb      | virginia, usa              |
| e.d. courant*   | brn, usa                   |
| a.v. afremov    | inn, russia                |
| g. fidicaro*    | bern, switzerland          |
| w. hebeler*     | washington, usa            |
| k. hatanaka     | riepi, japan               |
| k. imai         | kyoto, japan               |
| r.g. milner     | mit, usa                   |
| y. mori         | kek, japan                 |
| w.f.h. van oers | manitoba, canada           |
| c.y. prescott   | slac, usa                  |
| f. rathmann     | bose, germany              |
| yu.m. shatunov  | novosibirsk, russia        |
| v. soergel      | heidelberg, germany        |
| e. steffens     | erlangen, germany          |
| n.e. tyurin     | ibep, russia               |

\*honorary members

conference secretariat

e. novacco  
sezione infn di trieste  
via a. valerio, 7  
34127 trieste, italy  
ph. +39 040 5583367  
fax +39 040 5583350

trieste  
october 10\_16 2004

organized by  
the trieste section  
of infn association  
of italia members

hosted at  
the abbas adami  
international center  
for theoretical physics

sponsored by  
international union  
of pure and applied physics  
international spin committee  
physics consortium trieste  
consorzio trieste

