

# $\Delta G/G$ at COMPASS



Yann Bedfer  
Saclay - DAPNIA/SPhN  
On behalf of the COMPASS collaboration

## Physics Motivation

$$\frac{1}{2} = \frac{1}{2}\Delta\Sigma + \Delta G + L_q + L_g$$

- $\Delta\Sigma$  @ SMC, SLAC, HERMES, JLab :
  - Ellis-Jaffe sum rule violated.
  - $\Delta\Sigma$  contributes little to nucleon spin.
  - Where is the nucleon spin ?
- $\Delta G$  :
  - Next candidate contributor.
  - Large  $\Delta G$ ,  $\sim 2-3$  at SMC  $Q^2$  would mask quark spin via axial anomaly  
*Efremov, Teryaev, JINR Report E2-88-287 (1988).*  
*Altarelli, Ross, Phys.Lett. B212, 391 (1988).*


## COMPASS Spectrometer

- Muon Beam

- 160 GeV.
- $2 \cdot 10^8 \mu$  per Spill of  $\sim 15$  s.
- 76-80% polarisation.

- Target

- $LiD : f \simeq 40\%$ ,  $P_T \simeq 50\%$ .
- Cf. Kisseelev's talk.



- Spectrometer      *NIMA 577 (2007) 455*

- Two stages,  
each with HCalorimeter and  $\mu$ -filter.
- RICH : in stage 1.
- ECalorimeter : 2 (since mid-2004), 1 (in 2006).

## Asymmetry Measurement

- Two oppositely polarized target cells : *upstream*, *downstream*
- Polarization reversal by field rotation every 8 hours :

$$\frac{A^{\parallel}}{D} = \frac{1}{|P_T P_\mu| f D} \frac{1}{2} \left( \frac{N_{\textcolor{blue}{u}}^{\uparrow\downarrow} - N_{\textcolor{green}{d}}^{\uparrow\uparrow}}{N_{\textcolor{blue}{u}}^{\uparrow\downarrow} + N_{\textcolor{green}{d}}^{\uparrow\uparrow}} + \frac{N_{\textcolor{green}{d}}^{\uparrow\downarrow} - N_{\textcolor{blue}{u}}^{\uparrow\uparrow}}{N_{\textcolor{green}{d}}^{\uparrow\downarrow} + N_{\textcolor{blue}{u}}^{\uparrow\uparrow}} \right) \quad \textcolor{magenta}{D} = \text{Depolarization factor}$$

$$\textcolor{magenta}{P_T} \times \textcolor{magenta}{P_\mu} \times \textcolor{magenta}{f} \times \textcolor{magenta}{D} \simeq 50\% \times 80\% \times 40\% \times 60\% \simeq 10\%$$

- Weighted asymmetry

$$\frac{A^{\parallel}}{D} = \frac{1}{P_T} \frac{1}{2} \left( \frac{\sum_{\textcolor{blue}{u}}^{\uparrow\downarrow} w - \sum_{\textcolor{green}{d}}^{\uparrow\uparrow} w}{\sum_{\textcolor{blue}{u}}^{\uparrow\downarrow} w^2 + \sum_{\textcolor{green}{d}}^{\uparrow\uparrow} w^2} + \frac{\sum_{\textcolor{green}{d}}^{\uparrow\downarrow} w - \sum_{\textcolor{blue}{u}}^{\uparrow\uparrow} w}{\sum_{\textcolor{green}{d}}^{\uparrow\downarrow} w^2 + \sum_{\textcolor{blue}{u}}^{\uparrow\uparrow} w^2} \right) \quad w_i = \langle \textcolor{magenta}{P_\mu} f D \rangle_i$$

$$\Rightarrow \text{Gain in precision} = \sqrt{\langle w^2 \rangle / \langle w \rangle^2}$$

- Microwave reversal (once *per* ~month) cancels acceptance *vs.* target field correlation.
- 2006 : 3-cell target  $\Rightarrow$  Even better false asymmetry suppression. Rotation once *per* day.

## Data Taking



- Luminosity in the longitudinal mode :

	2002	2003	2004	2006
Integrated Luminosity ( $fb^{-1}$ )	0.43	0.58	0.92	0.85

- 2006 upgrade :
  - Larger acceptance : 70 mrd → 180 mrd.
  - Better RICH PID
  - Electromagnetic calorimetry.
- Only partially analyzed : 2002 ↔ 2004 (*most of the time*)
- COMPASS resumed data taking in 2007,  
w/ a polarized proton ( $NH_3$ ) target.


## $\Delta G$ via scaling violation in $g_1$

- COMPASS QCD fit :
    - Cf. talk of Helena Santos.
- ⇒ World data described well by two solutions, with  $\Delta G > 0$  and  $\Delta G < 0$ , resp. :



- ⇒ Small sensitivity to  $\Delta G$  ⇒ Need direct measurement
- ⇒ Sensitivity to assumed functional shape.

## $\Delta G/G$ direct measurement : Photon Gluon Fusion



**$q = c$**  : Open Charm production

- Triggered by PGF at LO  
(neglecting Intrinsic Charm)
- ⇒ Theory Golden Channel
- Experimentally difficult
- pQCD scale set by  $\hat{s} > 4m_c^2$



⇒ Explore all  $Q^2$

**$q = u,d,s,c$**  : High  $p_T$  Hadrons

- Competing LO-DIS, QCD-Compton and resolved photon processes.
- ⇒ Theoretical uncertainties.
- Higher statistics
- pQCD scale can be set by  $p_T$

## Open charm : $D^o$ meson reconstruction



- $D^o \rightarrow K\pi$ 
  - Thick target  
⇒ No Charm decay vertex reconstruction



- RICH PID
- + Kinematical cuts
  - Momentum fraction  $z_{D^o}$
  - $D^o$  decay angle

$S_{\text{eff}} = \text{effective Signal} = S^2 / (S + B)$   
where S and B Signal and Background counts.

- Favorable case :  $D^o$  from  $D^* \rightarrow D^o\pi \rightarrow K\pi\pi$



- 1/3 of  $D^o$ 's
- $D^*$  tagging by cut on 3-body invariant mass
- Bump =  $D^o \rightarrow K\pi\pi^o$  missing  $\pi^o$


## Open charm : Extraction of $\Delta G/G$

$$\frac{N^{\uparrow\downarrow} - N^{\uparrow\uparrow}}{N^{\uparrow\downarrow} + N^{\uparrow\uparrow}} = P_T P_\mu f \textcolor{blue}{a_{LL}} \frac{\textcolor{magenta}{S}}{\textcolor{magenta}{S} + \textcolor{magenta}{B}} \frac{\Delta G}{G} + \textcolor{green}{A}_B$$

- $A_B$  determined to be negligible on side bands.
- In order to minimize statistical error use event weighting  
 $w = P_\mu f \textcolor{blue}{a_{LL}} \textcolor{magenta}{S}/(\textcolor{magenta}{S} + \textcolor{magenta}{B})$
- Needed inputs :
  - Signal purity  $\textcolor{magenta}{S}/(\textcolor{magenta}{S} + \textcolor{magenta}{B})$  : derived from fit.
  - $\textcolor{blue}{a_{LL}}$  from a parameterization derived from MC simulation.

## Open charm : Analyzing power $a_{LL}$

$$\langle a_{LL} \rangle \frac{\Delta G}{G} \simeq \frac{\int \Delta \sigma^{PGF}(\hat{s}) \Delta G(x_g, \hat{s}) d\hat{s}}{\int \sigma^{PGF}(\hat{s}) G(x_g, \hat{s}) d\hat{s}}$$




*Correlation factor  $\simeq 82\%$*

- Hard scattering kinematics
  - Needs MC information
  - MC = AROMA. Checked *vs.* data.
- Parametrization with :  $y, Q^2, z_{D^o}, p_T^{\gamma D^o}$
- Using neural network

$\Rightarrow$  LO extraction of  $\Delta G/G$ .

## Open charm : Result



- Preliminary result from COMPASS 2002-2004 data

Systematics include :

- False asymmetry : Upper bound from high statistics including background.
- Background asymmetry.
- Fitting procedure.
- MC : charm mass and PDFs.


$$\Delta G/G = -0.57 \pm 0.41(\text{stat.}) \pm 0.17(\text{syst.})$$

$$\langle x_g \rangle = 0.15$$


$$\mu^2 \simeq 13 \text{ GeV}^2$$

## High $p_T$ hadrons


- Competing processes :







PGF



QCDC

LO ( $p_T$  via hadronization)

- + Resolved photons :

 $qq \rightarrow qq$  $qg \rightarrow qg$  $qg \rightarrow qg$  $gg \rightarrow gg$ 

(Introduce a dependence upon the poorly known polarized structure of the photon.)

- Factorization granted by hard scale.

- 2 different methods :

I) At LO :  $A_{||} \simeq ( R_{PGF} \langle a_{LL}^{PGF} \rangle + \sum R_i \langle a_{LL}^i \Delta q/q \rangle ) \Delta G/G + A_{Background}$

Determine  $R$ ,  $\langle a_{LL} \rangle$  and  $A_{Background}$  from MC simulation.

II) Asymmetry *vs.*  $p_T$  compared to theoretical prediction based on  $\Delta G$  parametrization.  
Can be NLO.


## High $p_T$ measurement via MC generator

- $Q^2 < 1 \text{ GeV}^2$ 
  - PYTHIA used :  
*pQCD + resolved photon + model dependent low scale processes.*
  - Hard scale defined by  $p_T$
  - Highest statistics.
- $Q^2 > 1 \text{ GeV}^2$ 
  - LEPTO used :  
*pQCD alone  $\Rightarrow$  better controlled.*
  - Hard scale defined by  $Q^2$
  - Lower statistics.
  - Resolved photons assumed negligible.
- Cf. presentation by K. Klimaszewski

## NLO calculation : high $p_T$ photoproduction

- Calculations by group of *BNL/Regensburg*.
  - Single high  $p_T$  hadron  
*Jäger, Stratmann, Vogelsang, Eur.Phys.J. C44 (2005) 533-543.*
  - Pair of high  $p_T$  hadrons  
*Hendlmeier, Stratmann, Schäfer, arXiv :0706.3766 [hep-ph]*
- Photoproduction :  $Q^2 < 0.5 \text{ GeV}^2$
- $\Delta G$  independent of MC model.
- Dependent upon functional shape  $\Delta g(x)$
- Need to validate calculation on unpolarized cross section

## General Trend in NLO Calculations at low $\sqrt{s}$




- Everything is fine at  $\sqrt{s} = 200$  GeV.
- Discrepancy grows at lower  $\sqrt{s}$ .
- All-order resummations of large logarithms greatly bridge the gap  
*Phys. Rev. D71 (2005) 114004.*
- Resummations not yet done in COMPASS case,  $\sqrt{s} = 17.3$  GeV.  
For the time being, assume they do not impact on the asymmetries.


Eur.Phys.J. C36 (2004) 371-374

## High $p_T$ photoproduction : Projections

- Measurement not released yet.
- Projections (*compared to GRSV scenarios*) :




Single hadron : Data analysis  
of  $\sim 1/3$  of recorded data.



Single hadron : Jäger et al.,  
*Eur.Phys.J. C44 (2005) 533-543*


## $\Delta G/G$ : Summary of results




- COMPASS disfavors the large values needed for  $\Delta G$  to account for the  $g_1$  results *via* the axial anomaly.
- But compatible with  $|\Delta G/G| \simeq 0.2 : 0.3$

## 2006 Upgrade : RICH

- Central part : CsI+MWPC photodetectors → MAPMTs
- Outer part : Faster electronics.





$\mu s$  range : Event dominated by halo muons.




$ns$  time resolution.

## 2006 Upgrade : RICH + Acceptance

- Combined effect on tracking of :
  - RICH improvements
  - + Enlarged acceptance : 70 mrd → 180 mrd



## 2006 Upgrade : ECalorimetry




- Electromagnetic calorimeter in first stage : 1500 lead glass blocks.
  - Installed in 2006. Included into trigger system in 2007.
  - 2006 data being analyzed.

## 2006 Upgrade : Impact on $\Delta G/G$ channels


- Gain in  $\mathcal{F}oM$  per incident muon.

(Caveat : -  $\mathcal{F}oM(\propto \delta\Delta G/G^{-2})$  evaluated on effective #events, ignoring event weighting.)

- Open charm  $D^*$  :  $\times 2.25$



(Extra gain from RICH PID below  $K$  threshold.)



$D^*$ 's in 1/2 of 2006 data.

- High  $p_T Q^2 > 1$  :  $\times 1.8$



## Conclusion

- Open charm  
 $\Delta G/G = -0.57 \pm 0.41(\text{stat.}) \pm 0.17(\text{syst.}) \quad x_g \simeq 0.15 \quad \mu^2 \simeq 13 \text{ GeV}^2$
- High  $p_T Q^2 > 1$       (*2002-2003 data*)  
 $\Delta G/G = 0.06 \pm 0.31(\text{stat.}) \pm 0.06(\text{syst.}) \quad x_g \simeq 0.13$
- High  $p_T Q^2 < 1$       (*PLB 633 (2006) 25-32*)  
 $\Delta G/G = 0.016 \pm 0.058(\text{stat.}) \pm 0.055(\text{syst.}) \quad x_g \simeq 0.085 \quad \mu^2 \simeq 3 \text{ GeV}^2$
- Favors low value of  $\Delta G$
- NLO extraction from high  $p_T$  photoproduction to be released.
- 2006 :
  - 1/2 more statistics.
  - Larger impact on  $\mathcal{FOM}$  due to upgrade.
- 2007 : Polarized proton target.  
(*Not optimum for  $\Delta G$  since  $fP_T$  reduced.*)
- 2008 : Hadron beam.

# SPARES

## COMPASS QCD fit : Results for $p$ , $n$ and $d$

- World data at  $Q^2 = 3 \text{ GeV}^2$ .  $p$ ,  $n$  and  $d$  with  $\Delta G > 0$ , and  $d$  with  $\Delta G < 0$   
 $\Delta G < 0$  preferred at low  $x$ .



## Open charm : Extraction of $\Delta G/G$

$$N_i = a\Phi(\sigma_S + \sigma_B)(1 + P_T P_\mu f(a_{LL} \frac{\Delta G}{G} \frac{\sigma_S}{\sigma_S + \sigma_B} + A_B \frac{\sigma_B}{\sigma_S + \sigma_B}))$$

for each of  $i = (u, d) \times (\uparrow\uparrow, \uparrow\downarrow)$

- Look at double ratio  $\frac{u^{\uparrow\downarrow} d^{\uparrow\uparrow}}{u^{\uparrow\uparrow} d^{\uparrow\downarrow}}$ 
  - $\Phi$  same for  $u$  and  $d$   $\Rightarrow$  cancels out.
  - Assume stable acceptance  $u/d$  ratio = $_i$   $a$  cancels out.
- Determine  $A_B = 0$  from side bands.
- Solve for  $\Delta G/G$
- Needed inputs :  $a_{LL}$  and purity.

## 2006 Upgrade : RICH + Acceptance

