



**The Abdus Salam
International Centre for Theoretical Physics**



1942-43

Sixth International Conference on Perspectives in Hadronic Physics

12 - 16 May 2008

Highlights from the COMPASS experiment at CERN.

F. Bradamante
*University of trieste / INFN
Trieste
Italy*

HIGHLIGHTS FROM THE COMPASS EXPERIMENT @ CERN

F. Bradamante

University of Trieste and INFN Trieste



on behalf of the *COMPASS* Collaboration



Sixth International Conference on Perspectives in Hadronic Physics
Trieste, May 16, 2008

**COmmon
Muon and
Proton
Apparatus for
Structure and
Spectroscopy**



NA58

**Czech Republic, Finland, France, Germany, India, Israel, Italy,
Japan, Poland, Portugal, Russia**

*Bielefeld, Bochum, Bonn, Burdwan, Calcutta, CERN,
Dubna, Erlangen, Freiburg, Heidelberg, Helsinki, Lisbon,
Mainz, Miyazaky, Moscow, Munich, Nagoya, Prague, Protvino,
Saclay, Tel Aviv, Torino, Trieste, Warsaw*

28 Institutes, ~230 physicists

COMPASS

- **experiment:** **thought of in** **April '94** **Trento workshop**
Nov. '94 **Trieste workshop**
@ICTP
Lol **March '95**
encouraged **June '95** **SPSLC in Cogne**
Proposal **March '96**
recommended **Sept. '96**
approved by RB **Feb. '97** **as NA58**
Technical run **2000**
Commissioning **2001**

- **since 2002 taking data with**
a new spectrometer with outstanding performances

- **merging of two programmes:** **HMC** **CHEOPS**
(muon beam) **(hadron beam)**




Physics program of COMPASS

- **Experiments with muon beam**
 - $\Delta G/G$
 - g_1
 - Transverse spin effects
 - Flavor decomposition of spin distribution functions
 - Vector meson production
 - Spin transfer in Λ -hyperon production
- **Experiments with hadron beams**
 - Pion and Kaon polarizabilities
 - Diffractive production of exotic states
 - Search for glueballs
 - Light meson spectroscopy
 - Production of double charmed baryons




longitudinally polarised muon beam
longitudinally or transversely polarised
target
calorimetry
particle identification

luminosity: $\sim 5 \cdot 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$
beam intensity: $2 \cdot 10^8 \mu^+/\text{spill}$ (4.8s/16.2s)
beam momentum: 160 GeV/c



The Spectrometer for the Muon Programme



TWO STAGE SPECTROMETER:
 Polarized beam and target
 SAT, LAT, PID
 $0.003 < x < 0.5$
 $10^{-3} < Q^2 < 10 \text{ GeV}^2$



WHERE ARE WE ?

- in 2002, 2003, 2004, 2006 and 2007 COMPASS has taken data in the **muon program configuration**
 - 160 GeV, polarized μ beam
 - 2002-2006 ${}^6\text{LiD}$ polarized target (\sim polarized deuterons)
 - 2007 NH_3 polarized target (\sim polarized protons)
- **2000 TB $\sim 5 \cdot 10^{10}$ events**
- **pilot run in 2004 for hadron program**
- **2008: hadron beam at 190 GeV for diffractive and central production**

physics results

1. $\Delta G/G$
2. $\Delta\Sigma$
3. Transversity
4. Cahn asymmetry
5. Pentaquark
6. Exclusive ρ^0
7. Λ physics

1. pion polarizability
2. PWA in diffractive scattering



THE COMPASS MUON PROGRAM


TWO CLASSES OF PHENOMENA:

- LONGITUDINAL SPIN CASE
- TRANSVERSE SPIN CASE



LONGITUDINAL SPIN CASE: the beginning

EMC 1988



$$\Gamma_1^p = 0.123 \pm 0.013 \pm 0.019$$

$$\Delta\Sigma = 0.12 \pm 0.17$$

→ SPIN CRISIS

LONGITUDINAL SPIN CASE

from polarised lepton – polarised nucleon DIS

$$d\sigma = d\bar{\sigma} \pm d\Delta\sigma$$

$$\frac{d\Delta\sigma}{dx dy} = \frac{e^4}{4\pi^2 Q^2} \cdot \left\{ \cos\alpha \cdot \left[\left(1 - \frac{y}{2} - \frac{y^2}{4} \cdot \gamma^2 \right) \cdot \mathbf{g}_1 - \frac{9}{2} \cdot \gamma^2 \cdot \mathbf{g}_2 \right] - \sin\alpha \cdot \cos\varphi \cdot \sqrt{1 - \frac{y}{2} - \frac{y^2}{4} \cdot \gamma^2} \cdot \gamma \cdot \left(\frac{y}{2} \cdot \mathbf{g}_1 + \mathbf{g}_2 \right) \right\}$$

with $\mathbf{g}_1(\mathbf{x}) \approx \sum_q e_q^2 \cdot [\Delta\mathbf{q}(\mathbf{x}) + \Delta\bar{\mathbf{q}}(\mathbf{x})]$ and $\Delta\mathbf{q} = \vec{\mathbf{q}} - \vec{\bar{\mathbf{q}}}$

first moments: $\Gamma_1 = \int \mathbf{g}_1(\mathbf{x}) dx$ $\Delta\mathbf{q} = \int \Delta\mathbf{q}(\mathbf{x}) dx$

from Γ_1^p measurement of EMC in 1988 and using complementary information from neutron and hyperon β -decay one obtained

$$\Delta\Sigma = \Delta\mathbf{u} + \Delta\mathbf{d} + \Delta\mathbf{s} = 0.12 \pm 0.17$$

at variance with naïve expectation

since $\frac{1}{2} = \frac{1}{2} \Delta\Sigma + \Delta\mathbf{G} + L_{q,g}$

necessity for measuring Γ_1^n

SMC, SLAC, HERMES

$\Delta\mathbf{q}$ and $\Delta\bar{\mathbf{q}}$ in SIDIS

SMC, HERMES, COMPASS

$\Delta\mathbf{G}$ in SIDIS

HERMES, COMPASS


LONGITUDINAL SPIN CASE

physics results

$\Delta G/G$



MEASUREMENTS OF THE GLUON POLARIZATION




FOUR LINES OF ATTACK:

1. Double spin asymmetry of the OPEN CHARM cross-section in high energy μ D scattering
2. Double spin asymmetry of the HIGH- p_t HADRON PAIRS in high energy μ D DIS ($Q^2 > 1 \text{ GeV}^2$)
3. Double spin asymmetry of the high- p_t hadron pairs in high energy μ D scattering ($Q^2 < 1 \text{ GeV}^2$)
4. Measurement of g_1 of the deuteron and QCD fit of all the world data

$\Delta G/G$ at COMPASS

Photon Gluon Fusion



$q = c$ cross section difference
in charmed meson production
→ theory well understood
→ experiment challenging

$q = u, d, s$ cross section difference
in 2+1 jet production
in COMPASS: events with
2 hadrons with high- p_t
→ experiment easy
→ theory more difficult


$\Delta G/G$ from Open Charm



D mass spectra

$$D^0 \rightarrow K + \pi$$

$$D^* \rightarrow D^0 + \pi_S \rightarrow K + \pi + \pi_S$$



APS, 13 April 2008

G.K. Mallot



$\Delta G/G$ from open charm

2002 – 2006 data $D^0 + D^*$

$$\Delta G/G = -0.49 \pm 0.27 \text{ (stat)} \pm 0.11 \text{ (syst)}$$

@ $\langle x_g \rangle \sim 0.11$, $\langle \mu^2 \rangle \sim 13 \text{ (GeV/c)}^2$

preliminary



$\Delta G/G$




from High- p_t hadron pairs



$\Delta G/G$ from High- p_t hadrons, $Q^2 > 1$ (GeV/c) 2

PGF and background

$$\frac{A_{LL}}{D} \approx \frac{a_{LL}^{PGF}}{D} \frac{\Delta G}{G} \frac{\sigma^{PGF}}{\sigma^{tot}} + A_1 \frac{a_{LL}^{LO}}{D} \frac{\sigma^{LO}}{\sigma^{tot}} + A_1 \frac{a_{LL}^{QCD-C}}{D} \frac{\sigma^{QCD-C}}{\sigma^{tot}}$$



$\Delta G/G$ from High- p_t hadrons, $Q^2 > 1 \text{ (GeV/c)}^2$

2002 – 2004 data: High p_T , $Q^2 > 1 \text{ GeV/c}^2$


$$\Delta G/G = 0.08 \pm 0.10 \text{ (stat)} \pm 0.05 \text{ (syst)}$$

@ $\langle x_g \rangle = 0.082$ (range: 0.055 – 0.123), $\mu^2 \sim 3 \text{ (GeV/c)}^2$

preliminary



$\Delta G/G$ from High- p_t hadrons



$\Delta G/G$ from High- p_t hadrons, $Q^2 < 1 \text{ (GeV/c)}^2$

2002 – 2004 data: High p_T , $Q^2 < 1 \text{ GeV/c}^2$


$$\Delta G/G = 0.016 \pm 0.058 \text{ (stat)} \pm 0.055 \text{ (syst)}$$

$$@ \langle x_g \rangle = 0.085, \mu^2 = 3 \text{ GeV}^2$$

preliminary



Glueon Polarization



COMPASS preliminary results

high- p_T pairs, $Q^2 > 1 \text{ GeV}^2$:

2002–2004

$$\Delta G/G = 0.08 \pm 0.10 \text{ (stat)} \pm 0.05 \text{ (syst)}$$

@ $\langle x_g \rangle = 0.082$ (range: 0.055 – 0.123), $\mu^2 \sim 3 \text{ (GeV/c)}^2$

high- p_t pairs, $Q^2 < 1 \text{ GeV}^2$:

2002–2004

$$\Delta G/G = 0.016 \pm 0.058 \text{ (stat)} \pm 0.055 \text{ (syst)}$$

@ $\langle x_g \rangle = 0.085$, $\mu^2 = 3 \text{ GeV}^2$


open charm:

2002–2006

$$\Delta G/G = -0.49 \pm 0.27 \text{ (stat)} \pm 0.11 \text{ (syst)}$$

@ $\langle x_g \rangle \sim 0.11$, $\langle \mu^2 \rangle \sim 13 \text{ (GeV/c)}^2$

Summary of results




g_1^d



g_1 of the deuteron (2002-2003)

Phys Lett B 612 (2005) 154



- most precise measurement for $0.004 < x < 0.03$
- new NLO QCD fit, precision of a_0 improves factor 2 ($Q^2 = 4 \text{ GeV}^2$)

$$a_0 = \Delta\Sigma(\overline{MS}) = 0.237^{+0.024}_{-0.029}$$



CONCLUSION from ΔG MEASUREMENTS:

ΔG SMALL

more precise measurements will come soon

COMPASS 2006

RHIC RUN6

....

$$\frac{1}{2} = \frac{1}{2} \Delta \Sigma + \Delta G + L_{q,g}$$

interest in
orbital angular momentum

GPD's

Ji's SUM RULE $J^q(t) = \frac{1}{2} \int dx x (H^q + E^q)$

more on **LONGITUDINAL SPIN CASE**

MEASUREMENT OF VALENCE QUARK POLARISATION



valence quark polarisation



hadron asymmetries

Semi-inclusive asymmetries

$$A^+ = \frac{\sigma_{\uparrow\downarrow}^{h+} - \sigma_{\uparrow\uparrow}^{h+}}{\sigma_{\uparrow\downarrow}^{h+} + \sigma_{\uparrow\uparrow}^{h+}} \quad A^- = \frac{\sigma_{\uparrow\downarrow}^{h-} - \sigma_{\uparrow\uparrow}^{h-}}{\sigma_{\uparrow\downarrow}^{h-} + \sigma_{\uparrow\uparrow}^{h-}}$$


$$A_1^h(x) = \frac{\sum_q e_q^2 (\Delta q(x) D_q^h + \Delta \bar{q}(x) D_{\bar{q}}^h)}{\sum_q e_q^2 (q(x) D_q^h + \bar{q}(x) D_{\bar{q}}^h)}$$

Difference asymmetry


$$A^+ = \frac{(\sigma_{\uparrow\downarrow}^{h+} - \sigma_{\uparrow\downarrow}^{h-}) - (\sigma_{\uparrow\uparrow}^{h+} - \sigma_{\uparrow\uparrow}^{h-})}{(\sigma_{\uparrow\downarrow}^{h+} - \sigma_{\uparrow\downarrow}^{h-}) + (\sigma_{\uparrow\uparrow}^{h+} - \sigma_{\uparrow\uparrow}^{h-})}$$

$$A_d^{\pi^+ - \pi^-}(x) = A_d^{K^+ - K^-}(x) = \frac{\Delta u_v(x) + \Delta d_v(x)}{u_v(x) + d_v(x)}$$



- Fragmentation functions $D_q^h = \int D_q^h(z) dz$ are **poorly known**
- Difference asymmetry originally **was proposed in:**
L.Frankfurt *et al.*, Phys. Lett. B230 (1989) 141
- First **was used in SMC:** B. Adeva *et al.*, Phys. Lett. B369 (1996) 93.
- Meaningful physics results for the deuteron target in LO QCD even **without hadron identification**



valence quark polarisation



comparison with other experiments




The Transverse Spin Case



Transverse Spin case

Large effects observed in hadronic interactions



VIEW LETTERS

10 MAY 1976






FIG. 3. Three components and magnitude of the $\Lambda^0 \rightarrow p + \pi^-$ asymmetry as a function of Λ^0 transverse momentum.



LETTERS B

1 August 1991



$$A_N = \frac{1}{P_{\text{beam}}} \frac{N_{\text{left}} - N_{\text{right}}}{\text{sum}}$$




Fig. 4. A_N versus x_F for π^+ , π^- and π^0 data.


Transverse Spin case

Large effects observed in hadronic interactions

Theoretical developments:

at leading order a third PDF is necessary for a complete description of the structure of the nucleon

R.L. Jaffe and X. Ji, Phys. Rev. Lett. **67** (1991) 552



- $\Delta_T q(x)$ being chiral-odd, it can be measured only in conjunction with another chiral-odd partner:

DY $\Delta_T q \otimes \Delta_T \bar{q}$

SIDIS $\Delta_T q \otimes FF$

Collins function

measurable in

$e+e \rightarrow$ hadrons

- relevance of transverse momentum dependent (TMD) PDF and FF

Sivers function


Many Workshops in recent years on *Transverse Momentum, spin, and position distributions of partons in hadrons*

Transversity – single hadron - 1

Collins and Sivers angles

$$\Phi_C = \phi_h - \phi_{S'}$$

$$\Phi_S = \phi_h - \phi_S$$



$\phi_{S'}$, azimuthal angle of spin vector of fragmenting quark ($\phi_{S''} = \pi - \phi_{S'}$)

ϕ_h azimuthal angle of hadron momentum


Transversity – single hadron - 2

first measurements of transverse spin asymmetries in DIS of high energy muons on a transversely polarized **deuteron target**

published single hadron asymmetries from 2002-2004 runs

- **Collins:** related to transverse quark distributions
- **Sivers:** related to intrinsic k_T

Phys Rev Lett 94 (2005) 202002
Nucl Phys B765 (2007) 31



Collins asymmetry for pions and kaons



preliminary

2002-2004 data

proton

(virtual photon asymm)

(lepton beam 2002-05 → DIS07)



final CERN-PH-EP/2008-002


hep-ex/0802.2160 (PRL)

2003-2004 data

deuteron

(virtual photon asymm)

COMPASS
sign convention



Independent Measurement of the Collins FF

measurable in e^+e^- annihilation


- first attempts to measured it from the correlation between the azimuthal angles of π 's from e^+e^- annihilation using LEP data

last years: great news from BELLE

the Collins FF is being measured in e^+e^- annihilation, and it is different from zero!

measurement of the correlation between the azimuthal angles of π 's in the near jet and in the far jet from e^+e^- annihilation

- 547 fb^{-1} charm corrected data sample,
- UL and UC double ratios



Collins asymmetries: SUMMARY

The facts:

- HERMES has measured on a proton target non-zero Collins asymmetries for π^+ and π^-
- COMPASS has measured on a deuteron target Collins asymmetries compatible with zero
- BELLE has produced the first results on Collins FF

Conclusion:

- Collins mechanism is a real phenomenon
- universality of Collins FF
- transversity can be measured in SIDIS


Present picture

- Collins: $\Delta_T u \sim -\Delta_T d$
 $\Delta_T^0 D(\text{fav.}) \sim -\Delta_T^0 D(\text{unfav})$

To extract TMD DF and FF GLOBAL ANALYSIS are necessary

Transversity

HERMES,
COMPASS,
BELLE



- This is the extraction of **transversity** from new experimental data.
- Compared to previous extraction
PRD75:054032,2007
- $\Delta_T u(x) > 0$ and $\Delta_T d(x) < 0$ The errors are diminished significantly.
- $\Delta_T u(x)$ became larger than that of the previous fit.


Sivers asymmetry for pions and kaons




preliminary
2002-2005 data
proton
(DIS07)



CERN-PH-EP/2008-002
hep-ex/0802.2160 (PRL)
2003-2004 data
deuteron



First moment of the Sivers functions



$$\diamond \Delta^N f_q^{(1)}(x) \equiv \int d^2 k_\perp \frac{k_\perp}{4m_p} \Delta^N f_{q/p\uparrow}(x, k_\perp) = -f_{1T}^{\perp(1)q}(x)$$

Sivers asymmetry

the measured asymmetry on **deuteron** compatible with zero
has been interpreted as

Evidence for the Absence of Gluon Orbital Angular Momentum in the Nucleon

S.J. Brodsky and S. Gardner, PLB643 (2006) 22

The approximate cancellation of the SSA measured on a deuterium target suggests that the gluon mechanism, and thus the orbital angular momentums carried by gluons in the nucleon, is small.

SUMMARY AND OUTLOOK

- a technically challenging new experiment is **IN OPERATION SINCE 2002**

“LHC” technologies *detectors*
 read-out
 data handling

- a privileged situation at CERN
- **MANY PHYSICS RESULTS** have been produced
MANY MORE IN THE PIPE-LINE
- **COMPASS** is foreseen to run up to the end of the present mid-term plan of CERN (2010)

BIG DISCOVERY POTENTIAL

- with some upgrade **COMPASS** might be an interesting option even in the second decade of this century



- SPARE SLIDES


Transversity: two hadrons - 1

$$\phi_{RS} = \phi_R - \phi_{S'} = \phi_R + \phi_S - \pi$$

$$\cos \phi_R = \frac{(\mathbf{q} \times \mathbf{l}) \cdot (\mathbf{q} \times \mathbf{R}_T)}{|\mathbf{q} \times \mathbf{l}| |\mathbf{q} \times \mathbf{R}_T|}$$

$$\sin \phi_R = \frac{(\mathbf{l} \times \mathbf{R}_T) \cdot \mathbf{q}}{|\mathbf{q} \times \mathbf{l}| |\mathbf{q} \times \mathbf{R}_T|}$$


$$\mathbf{R} = \frac{\mathbf{P}_1 - \mathbf{P}_2}{2} \quad \mathbf{P} = \mathbf{P}_1 + \mathbf{P}_2 \quad \mathbf{R}_T \perp \mathbf{P}$$



$$\frac{A_{UT}^{\sin \phi_{RS}}}{D_{NN} \cdot f \cdot P} = A_{RS} = \frac{\sum_i e_i^2 \Delta_T q_i(\mathbf{x}) H_i^{\zeta n}(z, M_h^2)}{\sum_i e_i^2 q_i(\mathbf{x}) D_i^h(\vec{z}, M_h^2)}$$

Transversity: two hadrons - 2

all +/- combinations





- precise measurement of few %
- systematics seems well under control
- also compatible with zero
- interesting to see **proton** in 2006



Exclusive ρ^0 production




Exclusive ρ^0 production (2002 data) - 1



Exclusive ρ^0 production - 2

ANGULAR DISTRIBUTIONS

$0.01 < Q^2 < 0.05$ $0.05 < Q^2 < 0.3$ $0.3 < Q^2 < 0.6$ $0.6 < Q^2 < 2.0$ $2.0 < Q^2 < 10 \text{ GeV}^2$



preliminary, statistical errors only, limited by MC

corrected for acceptance, smearing and efficiency (MC:DIPSI gen)
background not subtracted



Exclusive ρ^0 production - 3

measurement

of r_{00}^{04}

distribution :

$$W(\cos\theta) = \frac{3}{4} \left[(1 - r_{00}^{04}) + (3r_{00}^{04} - 1)\cos^2\theta \right]$$




Spin density matrix element:

$$r_{00}^{04} \sim \frac{|T_{01}|^2 + (\varepsilon + \delta)|T_{00}|^2}{\sigma_T + (\varepsilon + \delta)\sigma_L}$$

$T_{\lambda\rho\lambda\gamma}$ are helicity amplitudes
 meson photon

$$\frac{\text{SCHC}}{\sigma_T} \rightarrow \frac{\sigma_L}{\sigma_T}$$

$0.01 < Q^2 < 0.05$ $0.05 < Q^2 < 0.3$ $0.3 < Q^2 < 0.6$ $0.6 < Q^2 < 2.0$ $2.0 < Q^2 < 10 \text{ GeV}^2$



Exclusive ρ^0 production - 4

measurement

of r_{1-1}^{04}

and $\text{Im } r_{1-1}^3$

distribution :




$$W(\varphi) = \frac{1}{2\pi} [1 - 2r_{1-1}^{04} \cos 2\varphi + 2\text{Im } r_{1-1}^3 \underbrace{P_\mu}_{\text{beam polarisation}} \sqrt{1 - \varepsilon^2} \sin 2\varphi]$$

Spin density matrix element:

$$r_{1-1}^{04} \approx \frac{\text{Re}(T_{11} T_{-11}^*) - (\varepsilon + \delta) \cdot |T_{10}|^2}{\sigma_T + (\varepsilon + \delta) \cdot \sigma_L} \stackrel{\text{if SCHC holds}}{=} 0 \quad \text{weak violation}$$

$$\text{Im } r_{1-1}^3 = \dots = 0$$

0.01 < Q² < 0.05 0.05 < Q² < 0.3 0.3 < Q² < 0.6 0.6 < Q² < 2.0 2.0 < Q² < 10 GeV²



Pentaquark search



$\Phi(1860)$ Pentaquark search - 1


motivated by NA49 report of pentaquark candidate

- **COMPASS has large sample of the double-strange Ξ^- baryon (18000 Ξ^- , 11000 Ξ^+) from 2002-2003 data**
- **search for $\Xi^- \pi^-$ resonance**

$$\Phi(1860)^{- -} \rightarrow \Xi^- \pi^- \rightarrow \Lambda \pi^- \pi^- \rightarrow \rho \pi^- \pi^- \pi^-$$



$\Phi(1860)$ Pentaquark search - 2



Eur Phys J C41 (2005) 469

compare to yields

- **opposite-sign pairs:** $\Xi(1530)^0 \rightarrow \Xi^- \pi^+$, 1700 and 920 evts
- **like-sign pairs:** evts <79 and <89 at 99% CL
(expected ~400)





Fit: HERMES data

► HERMES data \diamond fit

$ep \rightarrow e\pi X$

$p_{lab} = 27.57 \text{ GeV}/c$

$ep \rightarrow eK X$



\diamond Diefenthaler, hep-ex/0612010 (2006)


Fit: COMPASS data

► COMPASS data \diamond fit

$eD \rightarrow e\pi X$

$p_{lab} = 160 \text{ GeV}/c$



$eD \rightarrow eKX$



\diamond A. Martin (COMPASS), Czech. J. Phys. 56, F33 (2006)



Predictions: COMPASS Hydrogen target

► Predictions for the COMPASS hydrogen target



Transversity: Λ polarization

2002-2003 data





Longitudinal polarization of Λ and anti Λ in DIS






Λ POLARIZATION (2002 data) - 1

Event selection



	$N(\Lambda)$	$N(\bar{\Lambda})$
E665	750	650
NOMAD	8087	649
HERMES 1996-2000	10568	1687
COMPASS 2002	7919	5062
COMPASS 2003	34789	20651

Λ POLARIZATION



$$\frac{dN}{d \cos \theta} = \frac{N_{tot}}{2} (1 + \alpha P \cos \theta)$$

2002 data



Λ POLARIZATION

2002 data

