



Azimuthal asymmetries in SIDIS off unpolarized targets at COMPASS

Andrea Bressan
(University and INFN - Trieste)

On behalf the COMPASS collaboration





Outline

- Introduction
- The COMPASS experiments
- Analysis and extraction of the asymmetries
- Results
- Conclusions







Experimental status

- Azimuthal modulations in $l p \rightarrow l' h X$ measured by

- EMC
- E665



Fits from M. Anselmino, V. Barone, E. Boglione, U. D'Alesio, F. Murgia, A. Prokudin, A. Kotzinian, and C. Turk

- Large modulations up to 40% for $\cos\phi$, while $\cos 2\phi \sim 5\%$
(with ϕ or ϕ_h the the hadron azimuthal angle in GNS)
- More recently ZEUS in the high- p_T (pQCD region)

Since last year, new data from COMPASS and HERMES





Cahn effect – just a reminder

The unpolarized SIDIS cross section is:

$$d\sigma^{lp \rightarrow l'hX} = \sum_q f_q(x, Q^2) \otimes d\sigma^{lp \rightarrow l'q} \otimes D_q^h(z, Q^2)$$

with f the PDF and D the FF

In collinear PM than the elementary xSection is

$$d\sigma^{lp \rightarrow l'q} \propto \hat{s}^2 + \hat{u}^2 \propto x(1 + (1-y)^2)$$

i.e. no dependence on ϕ_h . Taking into account the parton transverse momentum in the kinematics leads to:

$$\hat{s} = sx \left[1 - \frac{2k_T}{Q} \sqrt{1-y} \cdot \cos \phi \right] + O\left(\frac{k_T^2}{Q}\right) \quad \hat{u} = sx(1-y) \left[1 - \frac{2k_T}{Q\sqrt{1-y}} \cdot \cos \phi \right] + O\left(\frac{k_T^2}{Q}\right)$$

Resulting in the $\cos \phi_h$ and $\cos 2\phi_h$ modulations observed in the azimuthal distributions





Unpolarised target SIDIS cross-section

$$\frac{d\sigma}{dx dy d\psi dz d\phi_h dP_{h\perp}^2} = \frac{\alpha^2}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x}\right) \left\{ F_{UU,T} + \varepsilon F_{UU,L} + \sqrt{2\varepsilon(1+\varepsilon)} \cos\phi_h F_{UU}^{\cos\phi_h} \right. \\ \left. + \varepsilon \cos(2\phi_h) F_{UU}^{\cos 2\phi_h} + \lambda_e \sqrt{2\varepsilon(1-\varepsilon)} \sin\phi_h F_{LU}^{\sin\phi_h} \right\}$$

$$F_{LU}^{\sin\phi_h} = \frac{2M}{Q} \mathcal{C} \left[-\frac{\hat{h} \cdot k_T}{M_h} \left(x e H_1^\perp + \frac{M_h}{M} f_1 \frac{\tilde{G}^\perp}{z} \right) + \frac{\hat{h} \cdot p_T}{M} \left(x g^\perp D_1 + \frac{M_h}{M} h_1^\perp \frac{\tilde{E}}{z} \right) \right]$$

Cahn effect + Boer-Mulders DF

Boer-Mulders DF

$$F_{UU}^{\cos\phi_h} = \frac{2M}{Q} \mathcal{C} \left[-\frac{\hat{h} \cdot k_T}{M_h} \left(x h H_1^\perp + \frac{M_h}{M} f_1 \frac{\tilde{D}^\perp}{z} \right) - \frac{\hat{h} \cdot p_T}{M} \left(x f^\perp D_1 + \frac{M_h}{M} h_1^\perp \frac{\tilde{H}}{z} \right) \right]$$

the nucleon spin in an unpolarised

$$F_{UU}^{\cos 2\phi_h} = \mathcal{C} \left[-\frac{2(\hat{h} \cdot k_T)(\hat{h} \cdot p_T) - k_T \cdot p_T}{MM_h} h_1^\perp H_1^\perp \right]$$

clean **Boer-Mulders x Collins FF + Cahn effect**
and R. Sassot, Nucl. Phys. B



Common Muon and Proton

Apparatus for Structure and Spectroscopy

NA58

Czech Republic, Finland, France, Germany, India, Israel, Italy,
Japan, Poland, Portugal, Russia

*Bielefeld, Bochum, Bonn, Burdwan, Calcutta, CERN,
Dubna, Erlangen, Freiburg, Heidelberg, Helsinki, Lisbon,
Mainz, Miyazaky, Moscow, Munich, Prague, Protvino,
Saclay, Tel Aviv, Torino, Trieste, Warsaw, Yamagata*

28 Institutes, ~230 physicists




PALACIO DE CONGRESOS DE MADRID

DIS2009

Andrea Bressan

longitudinally polarised muon beam
longitudinally or transversely polarised target
calorimetry
particle identification

luminosity: $\sim 5 \cdot 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$
beam intensity: $2 \cdot 10^8 \mu^+/\text{spill}$ (4.8s/16.2s)
beam momentum: 160 GeV/c




COMPASS

LHC

SPS




The Spectrometer for the Muon Programme



TWO STAGE SPECTROMETER:

$$0.003 < x < 0.5$$

$$10^{-3} < Q^2 < 10 \text{ (GeV/c)}^2$$



SciFi Silicon

PALACIO DE CONGRESOS DE MADRID

And



Data used for this analysis

- part of the 2004 (${}^6\text{LiD}$ target) data collected with longitudinal (L) and transverse (T) polarization
- with both target orientation configurations to cancel possible polarization effects

Event selection:


DIS events...

- $Q^2 > 1 \text{ (GeV/c)}^2$
- $0.1 < y < 0.9$
- $W > 5 \text{ (GeV/c}^2\text{)}$

Hadrons





- $0.2 < z < 0.85$
- $0.1 < p_T < 1.5 \text{ (GeV/c)}$

Statistics of this analysis:





Mean kinematical values







unpolarised target SIDIS cross-section


to extract the asymmetries the azimuthal distributions have to be corrected by the apparatus acceptance

→ Final distributions fitted with the following function

$$N_{corr}(\phi) = N_0 \{ 1 + A_{\cos\phi} \cdot \cos\phi + A_{\cos 2\phi} \cdot \cos 2\phi + A_{\sin\phi} \cdot \sin\phi \}$$



final azimuthal distribution





Systematic Error

- The systematic error is evaluated from:
 - compatibility of results with L and T target polarization (different experimental conditions, different MCs)
 - comparison of results obtained using two different MCs with different settings for each data set (LEPTO default, standard COMPASS high pt; ~extreme cases)
 - compatibility of results from subsamples corresponding to:
 - different periods
 - different geometrical regions for the scattered muon






Results: $\sin\phi$ modulation

$$A_{\sin\phi} / \varepsilon_s$$

$$\varepsilon_s = \frac{2y\sqrt{1-y}}{1+(1-y)^2}$$



error bars:
statistical
errors

bands:
systematical
errors





Results: $\cos\phi$ modulation




$$A_{\cos\phi} / \epsilon_c$$

$$\epsilon_c = \frac{2(2-y)\sqrt{1-y}}{1+(1-y)^2}$$



What was expected



M. Anselmino, M. Boglione, A. Prokudin, C. Türk
Eur. Phys. J. A 31, 373-381 (2007)
does not include Boer – Mulders contribution






results: $\cos 2\phi$ modulation


$$A_{\cos 2\phi} / \varepsilon_2$$

$$\varepsilon_2 = \frac{2(2-y)}{1+(1-y)^2}$$





Predictions



— total Boer Mulders
- - - Cahn pQCD

V.Barone, A.Prokudin, B.Q.Ma
arXiv:0804.3024 [hep-ph]



PALACIO DE CONGRESOS DE MADRID


DIS2009

Andrea Bressan

17



Summary of the results





Summary

First results on unpolarized asymmetries:

- Results obtained separately for + and - hadrons
- $\sin\phi$ modulation compatible with 0
- $\cos\phi$ modulation up to 20% (for large z or p_T) and the overall trend is reproduced by the predictions
- $\cos 2\phi$ modulation smaller (10% at most). Overall good agreement with the predictions
- There is a difference between +h and -h asymmetries on $\cos\phi/\cos 2\phi$

All in all: new input for deeper understanding of the nucleon structure





Thank You

