
**AZIMUTHAL ASYMMETRIES of CHARGED HADRONS in SIDIS off
LONGITUDINALLY POLARIZED DEUTERON at COMPASS**

A.V. Efremov JINR, Dubna.

On behalf of the COMPASS Collaboration.


OUTLINE

- 1. Introduction: Theoretical framework & Motivation**
- 2. Method of analysis**
- 3. Data selection**
- 4. Results**
- 5. Conclusions and prospects**

Supported by the RFFI-CERN grant 08-02-91013. Published arXiv:1007.1562

Hadron azimuthal distributions are sources of information on PDFs and PFFs, characterizing longitudinal and transverse nucleon spin structure, e.g.:

$$d\sigma_h / d\phi \sim h_1(x) \otimes H_1^\perp(z) \cdot \sin(\phi + \phi_s) + \dots \quad \ell + N \rightarrow \ell' + X + h$$



A number of PDF's and PFF's enter in total SIDIS cross section

The cross section and asymmetry of SIDIS:

$$d\sigma = d\sigma_{00} + P_\mu d\sigma_{L0} + P_L (d\sigma_{0L} + P_\mu d\sigma_{LL}) + |P_T| (d\sigma_{0T} + P_\mu d\sigma_{LT}),$$

$$a(\phi) = \frac{d\sigma^{\leftrightarrow} - d\sigma^{\leftarrow\leftarrow}}{|P_L| (d\sigma^{\leftrightarrow} + d\sigma^{\leftarrow\leftarrow})} = \frac{d\sigma_{0L} + P_\mu d\sigma_{LL} - \tan\theta_\gamma (d\sigma_{0T} + P_\mu d\sigma_{LT})}{d\sigma_{00} + P_\mu d\sigma_{L0} (\approx f_1(x) \otimes D_1(z))}$$

where contributions to σ^{ij} (i=beam, j= target polarizations) from each quark and antiquark (up to the order of (M/Q)) have forms:

worm-gear-1

$$d\sigma_{0L} \propto \epsilon x h_{1L}^\perp(x) \otimes H_1^\perp(z) \sin(2\phi) + \sqrt{2\epsilon(1-\epsilon)} \frac{M}{Q} x^2 \left[h_L(x) \otimes H_1^\perp(z) + f_L^\perp(x) \otimes D_1(z) \right] \sin(\phi),$$

Twist 3

helicity

$$d\sigma_{LL} \propto \sqrt{1-\epsilon^2} x g_{1L}(x) \otimes D_1(z) - \sqrt{2\epsilon(1-\epsilon)} \frac{M}{Q} x^2 \left[g_L^\perp(x) \otimes D_1(z) + e_L(x) \otimes H_1^\perp(z) \right] \cos(\phi),$$

Twist 3

transversity

$$d\sigma_{0T} \propto \epsilon [x h_1(x) \otimes H_1^\perp(z) \sin(\phi + \phi_S) + x h_{1T}^\perp(x) \otimes H_1^\perp(z) \sin(3\phi - \phi_S)] - x f_{1T}^\perp(x) \otimes D_1(z) \sin(\phi - \phi_S)$$

Sivers

pretzelosity

Mulders&Tangerman
Boer&Mulders
Bucchetta et al.

\otimes =convolution in k_T

$$d\sigma_{LT} \propto \sqrt{1-\epsilon^2} x g_{1T}(x) \otimes D_1(z) \cos(\phi - \phi_S).$$

worm-gear-2



$\phi_S = 0$ for L-target

$$\text{tg } \theta_\gamma \cong \frac{M}{Q} x$$


Summary:

- $a(\phi)$ expected to be small, $\leq 1\%$.
- Methods of analysis should be adequate.
- Several asymmetry modulations should be seen in $a(\phi)$.
- Aims: search for $a(\phi)$, its possible $\sin(2\phi)$, $\sin(\phi)$ (Sivers + Transversity), $\sin(3\phi)$ (Pretzelosity) and $\cos(\phi)$ modulations and \mathbf{x} , \mathbf{z} and \mathbf{p}_h^T dependence of corresponding amplitudes.


The polarized ${}^6\text{LiD}$ -Target



4 target polarizations:



After few weeks



Polarization: ~ 50%

$$R_f(\varphi) = \frac{N_{+f}^U(\varphi)}{N_{-f}^D(\varphi)} \times \frac{N_{+f}^D(\varphi)}{N_{-f}^U(\varphi)} = \frac{\cancel{C_f^U(\varphi)} \cancel{L_{+f}^U} \sigma_+(\varphi)}{\cancel{C_f^D(\varphi)} \cancel{L_{-f}^D} \sigma_-(\varphi)} \times \frac{\cancel{C_f^D(\varphi)} \cancel{L_{+f}^D} \sigma_+(\varphi)}{\cancel{C_f^U(\varphi)} \cancel{L_{-f}^U} \sigma_-(\varphi)} = \frac{\sigma_+(\varphi)^2}{\sigma_-(\varphi)^2},$$

where

$N_{pf}^t(\varphi)$ is a number of events,

$t = U$ or D for Upper or Down cell,

$p = +$ or $-$ polarization (along or opposite to the beam),

$f = +$ or $-$ solenoid field direction (along or opposite to beam),

$C_f^t(\varphi)$ target acceptance factor (source of false asymmetries),

$L_{pf}^t = \Phi_{pf}^t n^t$ product of beam flux and target density,

σ_p spin dependent cross sections.

$L_{\pm f}^t$ and $C_f^t(\varphi)$ cancel if beam crosses both cells and if one combines periods with the same f .

$$R_f(\varphi) = \frac{(1 + P_{+f}^U a_f(\varphi))(1 + P_{+f}^D a_f(\varphi))}{(1 - P_{-f}^D a_f(\varphi))(1 - P_{-f}^U a_f(\varphi))},$$

$$a_f(\varphi) \cong \frac{R_f(\varphi) - 1}{P_{+f}^U + P_{+f}^D + P_{-f}^U + P_{-f}^D}$$

$$a_+(\varphi) \approx a_-(\varphi)$$

$$a(\varphi) = a_+(\varphi) \oplus a_-(\varphi) \rightarrow \text{final results}$$

Summary:

- the DR method has been tested using a part of data,
- possible ϕ -dependent false asymmetries, connected with the acceptance, are canceled,
- the DR method can be used for studies of small modulations of ϕ -asymmetries, of order 0.2% or smaller, the analysis of the full set of COMPASS L-data is in progress, first the data of 2002-2004 from deuterium, presented in this talk.

AIM: TO HAVE A CLEAN SAMPLE OF HADRONS

(1) Selection of “GOOD EVENTS” out of pre-selected sample of events with $Q^2 > 1 \text{ GeV}^2$ and $y > 0.1$
(=167.5 M from 2002, 2003, 2004 data taking)

EXCLUDED EVENTS:

- originated from bad spills,
- with a number of rec.prim.vertex > 1 ,
- $\chi^2/\text{NDF} > 2$,
- Z vertex outside the fiducial volume U or D- cell,
- $140 \text{ GeV} > E(\text{muon}) > 180 \text{ GeV}$,
- invariant mass $W < 5 \text{ GeV}$,
- $y > 0.9$.

= 58% of initial sample

(2) Selection of “GOOD TRACKS” from “GOOD EVENTS”.

Total number of tracks from “GOOD EVENTS” = 290 M

Excluded tracks:

- identified as muons,
- with z-variable >1 ,
- with $p_T^h < 0.1$ GeV -----→ “GOOD TRACKS” = 157 M

(3) Selection of “GOOD HADRONS” from “GOOD TRACKS”.



Each track should:

- hit one of the hadron calorimeters HCAL1 or HCAL2,
- have an associated energy cluster $E_{hcal1} > 5$ GeV or $E_{hcal2} > 7$ GeV,
- energy cluster coordinates compatible with the track coordinates,
- energy cluster compatible with the momentum of the track →
“GOOD HADRONS” = 53 M (25 M h^- + 28 M h^+)

(4) Each “GOOD HADRON” enters in considerations of asymmetries in restricted region

$$x = 0.004 - 0.7, \quad z = 0.2 - 0.9, \quad p_T^h = 0.1 - 1 \text{ GeV}/c$$

The weighted sum of azimuthal asymmetries $a(\phi) = a_+(\phi) \otimes a_-(\phi)$ for h^- (left) and h^+ (right) averaged over all kinematical variables :



Fit parameters	h^-	h^+	h^-	h^+
$\times 10^4$				
a^{const}	23 ± 17	40 ± 15	23 ± 16	35 ± 15
$a^{\sin \phi}$	15 ± 23	-30 ± 21	-	-
$a^{\sin 2\phi}$	30 ± 23	-24 ± 21	-	-
$a^{\sin 3\phi}$	40 ± 24	-10 ± 21	-	-
$a^{\cos \phi}$	-4 ± 24	38 ± 22	-	-
$\chi^2/\text{n.d.f.}$	6.1/5	1.0/5	10.4/9	7.0/9

Fit function :

$$a(\phi) = a^{\text{const}} + a^{\sin \phi} \sin(\phi) + a^{\sin 2\phi} \sin(2\phi) + a^{\sin 3\phi} \sin(3\phi) + a^{\cos \phi} \cos(\phi) \quad \text{OR} \quad a(\phi) = a^{\text{const}}$$


—Within stat. precision of about 1-1.5 σ , ϕ -dependent amplitudes are compatible with zero; fits by constants: OK,

—Parameters a^{const} are different from zero and about equal for h^+ and h^-

REMINDE: $a^{\text{const}} \propto d\sigma_{LL} \propto g_{1L}(x)/f_1(x)$, where g_{1L} is helicity PDF convoluted with non-polarized PFF. For isoscalar D-target it expected to be weakly dependent on the hadron charge.

—Qualitatively (to be confirmed by higher statistics) $a^{\sin \phi}$, $a^{\sin 2\phi}$ and $a^{\sin 3\phi}$ have opposite signs for h^- and h^+ predicted by quark models of nucleon due to WW-relation between h_{1L} and h_1 and relation $h_{1T}^+ h_1 = -1/2[h_{1L}^+]^2$.


Dependence of the parameter a^{const} for h^+ and h^- on kinematical variables:



- $a^{\text{const}}(x)/|P_\mu|D_0(x) \equiv A_d^h(x)$ (D_0 is a virtual photon depolarization factor) is in agreement with COMPASS published data (PLB660(2008)458),
- $a^{\text{const}}(z, p_T^h)$ for h^- and for h^+ : small and flat.

— Statistical errors are shown, systematic ones are estimated to be smaller: global systematic multiplicative errors are smaller than 6%.

Dependence of the parameter $a^{\sin\phi}$ for h^+ and h^- on kinematic variables:



— $a^{\sin\phi}(x)$ are less pronounced than the HERMES ones [Phys.Lett. B562 (203)182],


— $a^{\sin\phi}(z, p_T^h)$ is flat and do not confirm the HERMES trends.

REMINDE: $a^{\sin\phi} \propto -d\sigma_{0L} + \tan\theta_\gamma d\sigma_{0T} \propto -\frac{M}{Q} x \left((h_L(x) - h_T(x)) \otimes H_1^\perp(z) + (f_L^\perp(x) + f_{TT}^\perp) \otimes D_1(z) \right)$

where $h_L(x)$ and $f_L^\perp(x)$ are pure twist-3 PDF.

NOTE: HERMES data are for identified π^+ and π^- and at smaller $\langle Q^2 \rangle$.

Dependence of the parameter $a^{\sin 2\phi}$ for h^+ and h^- on kinematic variables:




— $a^{\sin 2\phi}(x)$ are small and in general agree with HERMES and theoretical predictions by H.Avakian et al., Phys.Rev. D77 (2008) 014023,

— $a^{\sin 2\phi}(z, p_T^h)$ - no other data.

REMINDE: $a^{\sin 2\phi} \propto d\sigma_{0L} \propto xh_{1L}^\perp(x) \otimes H_1^\perp(z)$, where h_{1L}^\perp is a PDF not seen yet. It is linked with the transversity PDF h_1 by a relation of the Wandzura-Wilczek type.

$$h_{1L}^{\perp(1)}(x) \approx -x^2 \int_x^1 \frac{dy}{y^2} h_1(y)$$


Dependence of the parameter $a^{\sin 3\phi}$ for h^+ and h^- on kinematic variables:



— $a^{\sin 3\phi}(x)$ are small, compatible with zero. But some peculiarities: points for h^- are mostly positive while these for h^+ are mostly negative as for the COMPASS results for the amplitude of the $\sin(3\phi - \phi_s)$ modulation extracted from the data with transversally polarized D-target. (A.Kotzinian, DIS2007 [arXiv:07093253])

REMINDE: $a^{\sin 3\phi} \propto d\sigma_{0T} \propto x h_{1T}^\perp \otimes H_1^\perp(z)$, where h_{1T}^\perp is pretzelosity PDF additionally suppressed by $\tan(\theta_\gamma) \sim \frac{M}{Q}$.

Dependence of the parameter $a^{\cos\phi}$ for h^+ and h^- on kinematic variables:



- $a^{\cos\phi}(x)$ increasing with x in absolute value,

- $a^{\cos\phi}(z)$ and $a^{\cos\phi}(p_T^h)$ small, flat and consistent with zero,

- studied for the first time.

Remind:

$$a^{\cos\phi} \propto -d\sigma_{LL} + \tan\theta_\gamma d\sigma_{LT} \propto \frac{M}{Q} x \left((g_L^\perp(x) - g_{1T}(x)) \otimes D_1(z) + e_L(x) \otimes H_1^\perp(z) \right)$$

g_L^\perp and e_L are pure twist-3 PDF (analog to Cahn effect in unpolarized SIDIS)




1. Integrated over x , z and p_h^T , all ϕ -modulation are consistent with zero within errors, while ϕ -independent parts differ from zero and are almost equal for h^- and h^+ .
2. In study of amplitudes over range $0.004 < x < 0.7$, $0.2 < z < 0.9$ and $0.1 < p_h^T < 1$ GeV/c it was found:
 - the ϕ -independent parts $\alpha^{const}(x)/D_0$ are in agreement with COMPASS data on $A_d^h(x)$, calculated by another method and using different cuts;
 - the amplitudes $\alpha^{\sin\phi}(x, z, p_h^T)$ are small and in general compatible with the HERMES data, if one takes into account the difference in x , W and Q^2 between the two experiments;
 - other amplitudes are consistent with zero within statistical errors of about 0.5% (systematical errors are estimated to be much smaller).
3. Tests have shown that lower cut, $z > 0.05$, gives identical results with smaller errors.
4. Data of 2006 from deuterium target will be added. Data of 2007 from hydrogen target will be interesting to compare with effects observed by COMPASS and HERMES on transversally polarized targets.

Thank you!


BACK UP SLIDES. STABILITY of RESULTS



Z - stability of results






"Pulls" distribution for a_i



Expected: Mean = 0.
 Sigma = 1.

ϕ - stability of rates



$$F = F_+ \oplus F_-$$

$$F_+ = \frac{N_{++}^u}{N_{++}^D} \cdot \frac{N_{-+}^D}{N_{-+}^U} = \frac{L_{++}^2}{L_{-+}^2}$$

$$F_1 = \frac{N_{+-}^U}{N_{+-}^D} \cdot \frac{N_{--}^D}{N_{--}^U} = \frac{L_{+-}^2}{L_{--}^2}$$

BACK UP SLIDES. PDF & FF IN SIDIS.



$f_1(x) \equiv q(x)$ is the PDF of non-polarized quarks in a non-polarized target,

$g_{1L}(x) \equiv g_1(x) \equiv \Delta q(x)$ is the PDF of the longitudinally polarized quarks in the longitudinally polarized target (helicity PDF),

$g_{1T}(x)$ is the same as $g_1(x)$ but in the transversely polarized target,

$h_1(x)$ is the PDF of the transversely polarized quark with polarization parallel to that one of a transversely polarized target (so-called transversity PDF),

$h_{1L}^\perp(x)(h_{1T}^\perp)$ is the PDF of the transversely polarized quark in direction of transverse momentum in longitudinally (transversely) polarized target (so-called pretzelosity PDF),

$h_1^\perp(x)$ is the PDF of the transversely polarized quark (perpendicular to transverse momentum) in the non-polarized target (so-called Boer-Mulders PDF),

$f_{1T}^\perp(x)(f_{1L}^\perp)$ is the PDF responsible for a left-right asymmetry in the distribution of the non-polarized quarks in the transversely (longitudinally) polarized target (so-called Sivers PDF),

$D_1(z)$ is the PFF of the non-polarized quark in the non-polarized or spinless produced hadron,

$H_1^\perp(z)$ is the PFF responsible for a left-right asymmetry in the fragmentation of a transversely polarized quark into a non-polarized or spinless produced hadron (so-called Collins PFF),

$e, e_L, g^\perp, g_L^\perp, h, h_L, f^\perp$ and f_L^\perp are pure twist-3 terms entering the cross section with a factor M/Q and having no clear physical interpretation.

Sources of false asymmetries are identified using a simplified expression for

$$A(\phi) \approx \frac{dN_+(\phi)}{N_+ d\phi} - \frac{dN_-(\phi)}{N_- d\phi} \approx p_0 + p_1 \sin(\phi) + p_2 \sin(2\phi) + p_3 \sin(3\phi) + p_4 \cos(\phi) + \dots$$

$$\approx p_0 + p_1 \sin(\phi).$$

Remind, that $\phi = \arccos \left(\frac{(\vec{\ell} \times \vec{\ell}') \cdot (\vec{q} \times \vec{p}_h)}{|\vec{\ell} \times \vec{\ell}'| |\vec{q} \times \vec{p}_h|} \right) \cdot \text{sign} \left[\vec{p}_h \cdot (\vec{\ell} \times \vec{\ell}') \right].$


$\sin(\phi)$ appears from the vector product $(\vec{\ell} \times \vec{p}_h)$, which is pseudo-vector.
But it could appear only being multiplied by another pseudo-vector:

spin of the target	\vec{S} , with a fraction p_s	} physics asym.
spin of the muon	$\vec{\mu}$, " p_μ	
target magnetic field	\vec{H} , " p_H	} false asym., due to incomplete knowledge of \vec{H} and/or misalignments
product	$(\vec{H} \times \vec{\mu})$, " $p_{H\mu}$	
product	$(\vec{H} \times \vec{S})$, " p_{HS}	


So, $p_1 \sim p_s + p_\mu + p_{H\mu} + p_{HS} + p_H$, where false asym depend on:

- directions of the field,
 - track extrapolations,
 - sign of particles.
- } different for U and D cells.

BACK UP SLIDES.KINEMATICS



The distribution of events, passed all data selection cuts, vs. Q^2 (left) and vs. y (right).



The distribution of identified good charged hadrons vs. z (left) and vs. p_T^h (right).₂₀

x bins	z bins	p_T^h bins (GeV)
	0.05(0.120)0.200	
0.004(0.010)0.012	0.200(0.216)0.234	0.100(0.177)0.239
0.012(0.020)0.022	0.234(0.253)0.275	0.239(0.289)0.337
0.022(0.031)0.035	0.275(0.299)0.327	0.337(0.385)0.433
0.035(0.053)0.076	0.327(0.361)0.400	0.433(0.485)0.542
0.076(0.098)0.132	0.400(0.455)0.523	0.542(0.610)0.689
0.132(0.190)0.700	0.523(0.661)0.900	0.689(0.814)1.000

The size of each bin is optimized to have ≥ 1 M of events

The first z bin (0.05 – 0.2) has been used for tests only