

Longitudinal polarization of the Λ and $\bar{\Lambda}$ hyperons in DIS at COMPASS (2003-2004). Dependence on the target polarization.

Salmina Ekaterina

JINR, Dubna
On behalf of COMPASS collaboration

04.10.2010

- Longitudinal polarization of Λ and $\bar{\Lambda}$ hyperons in DIS (averaged on target polarization). ¹
- Dependence of Λ and $\bar{\Lambda}$ longitudinal polarization on the target polarization.

¹ COMPASS Collab. EPJC 64 (2009) 171-179

Physical Motivation


Longitudinal polarization of Λ and $\bar{\Lambda}$ in DIS is sensitive to:

- $s(x), \bar{s}(x)$
- polarization of strange quarks Δs


$$\Delta s = \int dx [s_{\uparrow}(x) - s_{\downarrow}(x) + \bar{s}_{\uparrow}(x) - \bar{s}_{\downarrow}(x)]$$

- Λ spin structure

Example of quark spin transfer to Λ in DIS





COMPASS Spectrometer setup



- Year 2003:
 $P_b = -0.76 \pm 0.04$
- Year 2004:
 $P_b = -0.80 \pm 0.04$
- 160 GeV μ^+ beam
- $2.8 \cdot 10^8 \mu/\text{spill}$ ($4.8 \text{ s}/16.8 \text{ s}$)
- $Q^2 > 1 (\text{GeV}/c)^2$: $31.2 \cdot 10^7$ events

Polarized target



- target material: ${}^6\text{LiD}$
- polarisation: $> 50\%$
- dilution factor: ~ 0.4
- Dynamic Nuclear Polarization
- solenoid field: 2.5 T
acceptance: 70 mrad
- ${}^3\text{He}/{}^4\text{He}$: $T_{min} \approx 50 \text{ mK}$
- two 60 cm long target cells with opposite polarization
- regular polarization reversal by field rotation



Production of Λ , $\bar{\Lambda}$ and K

- $\mu^+ + d \rightarrow \mu^+ + \Lambda + X$
 $\Lambda \rightarrow p + \pi^-$
- $\mu^+ + d \rightarrow \mu^+ + \bar{\Lambda} + X$
 $\bar{\Lambda} \rightarrow \bar{p} + \pi^+$
- $\mu^+ + d \rightarrow \mu^+ + K_S + X$
 $K_S \rightarrow \pi^+ + \pi^-$

No PID identification

Event selection

- Primary vertex inside the target
- Secondary vertex: 5 cm downstream of the last target cell
- $p_T > 23 \text{ MeV}/c$
- $\theta < 0.01 \text{ rad}$
- $Q^2 > 1 \text{ (GeV}/c)^2$
- $0.2 < y < 0.9$
- $p_{\pm} > 1 \text{ GeV}/c$
- $0.05 < x_F < 0.5$
- $-1 < \cos \theta < 0.6$




Statistics: comparison with other experiments

Experiment	Λ	$\bar{\Lambda}$
E665	750	650
NOMAD	8087	649
HERMES	7300	1687
RHIC	13000	10000
COMPASS	70000	42000


COMPASS has the largest number of Λ and $\bar{\Lambda}$.

Invariant mass example: year 2004, Λ and $\bar{\Lambda}$

Λ , 2004 DATA



$\bar{\Lambda}$, 2004 DATA




Sideband subtraction method was used to obtain $\cos \theta$ angular distribution.

Bands regions: $(-5; -3)$, $(-1.5; 1.5)$, $(3; 5) \sigma$ from mass peak.

Longitudinal Λ ($\bar{\Lambda}$) polarisation

$$\frac{1}{N_{tot}} \frac{dN}{dcos\theta} = \frac{1}{2}(1 + \alpha P_L \cos \theta)$$



P_L - longitudinal polarisation of hyperon.

$\alpha = +(-)0.642 \pm 0.013$ – Λ ($\bar{\Lambda}$) decay parameter.



By definition longitudinal spin transfer is:

$$P_L = D_{LL} P_b D(y),$$

Depolarisation factor


$$D(y) = \frac{1-(1-y)^2}{1+(1-y)^2}$$

Example of angular distribution fits



- Angular dependencies for $\Lambda, \bar{\Lambda}$
- 2004 year events

Results: Comparison of Λ and $\bar{\Lambda}$: x




- $D_{LL}^{\Lambda} \neq D_{LL}^{\bar{\Lambda}}$
- $D_{LL}^{\Lambda} \simeq 0$

$$D_{LL}^{\Lambda} = -0.012 \pm 0.047 \pm 0.024$$

$$D_{LL}^{\bar{\Lambda}} = 0.249 \pm 0.056 \pm 0.049$$

Results: Comparison of Λ and $\bar{\Lambda}$: x_F




- $D_{LL}^{\bar{\Lambda}}$ rises with x_F

$$D_{LL}^{\Lambda} = -0.012 \pm 0.047 \pm 0.024$$


$$D_{LL}^{\bar{\Lambda}} = 0.249 \pm 0.056 \pm 0.049$$

Results: Comparison with other experiments: Λ






- COMPASS results agree with other experiments.

Comparison with theory ($\bar{\Lambda}$): CTEQ5 and GRV98





- Data for $\bar{\Lambda}$ are sensitive to the $s(x)$ distribution

Dependence on the target polarisation



+ : target pol. is same to μ pol.

- : target pol. is opposite to μ pol.



Averaged over full kinematics :

$$\Delta P^\Lambda = P_+^\Lambda - P_-^\Lambda = -0.01 \pm 0.04$$
$$\Delta P^{\bar{\Lambda}} = P_+^{\bar{\Lambda}} - P_-^{\bar{\Lambda}} = +0.01 \pm 0.05$$

No significant dependence is found.

Summary:

- The presented data are the most precise measurements to date of the longitudinal spin transfer to Λ and $\bar{\Lambda}$ in DIS.

$$D_{LL}^{\Lambda} = 0 : -0.012 \pm 0.047 \pm 0.024$$

$$D_{LL}^{\bar{\Lambda}} \neq 0 : 0.249 \pm 0.056 \pm 0.049$$

$$D_{LL}^{\Lambda} \neq D_{LL}^{\bar{\Lambda}}$$

- First measurement of the $\Lambda(\bar{\Lambda})$ polarization for different target polarization. No significant dependence is found.
- Comparison with theory:
Spin transfer to $\bar{\Lambda}$ is sensitive to $\bar{s}(x)$


Backup slides

Estimation of systematic errors

	Λ	$\bar{\Lambda}$
Spin transfer to kaons, $\delta(MC_1)$	0.016	0.016
Variation of selection cuts, $\delta(MC_2)$	0.016	0.044
Uncertainty of the ss-method, $\delta(ss)$	0.010	0.016
σ_{syst}	0.024	0.049


Invariant mass of Λ on $\cos\theta$

COMPASS 2003, Preliminary




Kaon background is important at large $\cos\theta$

Example of diquark spin transfer to Λ in DIS



Comparison with theory (Λ): CTEQ5 and GRV98



- CTEQ5 – solid line
- GRV98 – dashed line
- $D_{LL}(s) = 0$ BG and SU(6) models – 2 lower lines

Polarization of Λ from quark fragmentation

Λ polarization from struck quark fragmentation in parton model:

$$P_\Lambda = \frac{\sum_q e_q^2 [P_b D(y) q(x) + P_T \Delta q(x)] \Delta D_q^\Lambda(z)}{\sum_q e_q^2 [q(x) + P_b P_T D(y) \Delta q(x)] D_q^\Lambda(z)}$$

- $P_b D(y) q(x)$ – spin transfer from polarized muon
- $P_T \Delta q(x)$ – spin transfer from polarized quark

A. Kotzinian, A. Bravar, D. von Harrach, *Eur.Phys.J.* **C2**, 329-337 (1998), hep-ph/9701384

Spin transfer of Λ and $\bar{\Lambda}$: $P_T = 0$

$$D_{LL}^{\Lambda}(x, z) \approx \frac{1}{9} \frac{s(x)\Delta D_s^{\Lambda}(z)}{\sum_q e_q^2 q(x) D_q^{\Lambda}(z)},$$
$$D_{LL}^{\bar{\Lambda}}(x, z) \approx \frac{1}{9} \frac{\bar{s}(x)\Delta D_{\bar{s}}^{\bar{\Lambda}}(z)}{\sum_q e_q^2 q(x) D_q^{\bar{\Lambda}}(z)}$$

$$D_{LL}(\bar{\Lambda}) > D_{LL}(\Lambda)$$

- $s(x) \neq \bar{s}(x)$
- $D_q^{\Lambda}(z) > D_q^{\bar{\Lambda}}$

Longitudinal polarization and spin transfer equations

$$\frac{dN}{d\Omega} = \frac{N_{tot}}{4\pi} (1 + \alpha \vec{P} \cdot \vec{k})$$

$\alpha = +(-)0.642 \pm 0.013$ – Λ ($\bar{\Lambda}$) decay parameter.

$$\frac{1}{N_{tot}} \frac{dN}{dcos\theta} = \frac{1}{2} (1 + \alpha P_L \cos \theta)$$

By definition longitudinal spin transfer is:


$$P_L = D_{LL} P_b D(y),$$

where P_b – beam polarization and $D(y)$ – depolarization factor.

$$D(y) = \frac{1 - (1 - y)^2}{1 + (1 - y)^2}$$


Measurement of $\cos\theta$ distribution gives access to P_L and so to D_{LL}

Results: Comparison with other experiments: Λ



- COMPASS results agree with other experiments.

Results: Comparison with other experiments: $\bar{\Lambda}$




- COMPASS data are in agreement with NOMAD data
- it is the only data on x_F - dependence of $\bar{\Lambda}$

Production of Λ , $\bar{\Lambda}$ and K

- $\mu^+ + d \rightarrow \mu^+ + \Lambda + X$
 $\Lambda \rightarrow p + \pi^-$
- $\mu^+ + d \rightarrow \mu^+ + \bar{\Lambda} + X$
 $\bar{\Lambda} \rightarrow \bar{p} + \pi^+$
- $\mu^+ + d \rightarrow \mu^+ + K_S + X$
 $K_S \rightarrow \pi^+ + \pi^-$

No PID identification

Kinematic distributions for the selected Λ sample



Mean values:

$$\langle x \rangle = 0.05$$





$$\langle x_F \rangle = 0.23$$

$$\langle y \rangle = 0.46$$

$$\langle Q^2 \rangle = 3.31 \text{ (GeV}/c)^2$$

Good agreement between data and MC
Wide range of x_{Bj}

Kinematic distributions for the selected $\bar{\Lambda}$ sample



Mean values:

$$\langle x \rangle = 0.05$$

$$\langle x_F \rangle = 0.22$$



$$\langle y \rangle = 0.48$$

$$\langle Q^2 \rangle = 3.27 \text{ (GeV/c)}^2$$

Λ and $\bar{\Lambda}$ have similar kinematic regions

Fit for $\Delta P/P$

Dependence on pol. PDFs



$$\Delta P/P = \frac{P_- - P_+}{(P_- + P_+)/2}, \quad \text{on y axis } \frac{\langle D(y) P_b \rangle}{2fP_T} \frac{\Delta P^{\bar{\Lambda}}(x)}{P^{\bar{\Lambda}}(x)}$$

$\Delta P/P$ changes a sign in x_{Bj} region

Longitudinal Λ ($\bar{\Lambda}$) polarisation

Unpolarised target:

$$P_\Lambda = \frac{\sum_q e_q^2 P_b D(y) q(x) \Delta D_q^\Lambda(z)}{\sum_q e_q^2 q(x) D_q^\Lambda(z)}$$

Polarised target:


$$P_\Lambda = \frac{\sum_q e_q^2 [P_b D(y) q(x) + P_T \Delta q(x)] \Delta D_q^\Lambda(z)}{\sum_q e_q^2 [q(x) + P_b P_T D(y) \Delta q(x)] D_q^\Lambda(z)}$$

By definition longitudinal spin transfer is:

$$P_L = D_{LL} P_b D(y),$$


$$\text{Depolarisation factor } D(y) = \frac{1 - (1-y)^2}{1 + (1-y)^2}$$

Results: Comparison with other experiments: Λ



- COMPASS results agree with other experiments.

Results: Comparison with other experiments: $\bar{\Lambda}$



- COMPASS data are in agreement with NOMAD data
- it is the only data on x_F - dependence of $\bar{\Lambda}$