


Vector-Meson & DVCS Measurements in *COMPASS*-II

Heiner Wollny
CEA-Saclay Irfu/SPhN
on behalf of COMPASS





DIS 2011
XIX International Workshop on Deep Inelastic Scattering and Related Subjects

Motivation

Deep Inelastic Scattering:



$$\mu p \rightarrow \mu' X$$



PDFs: $q(x), \Delta q(x)$

Deeply Virtual Compton Scattering:

$$\mu p \rightarrow \mu' p' \gamma$$



GPDs: $H(x, \xi, t), \tilde{H}(x, \xi, t)$





COMPASS-II will start in 2012


230 physicists, 10 countries, 25 institutes

COMPASS II


SPS









- using existing COMPASS detector
- Upgrades of existing em. calorimeter ECAL1&2








- using existing COMPASS detector
- Upgrades of existing em. calorimeter ECAL1&2
- ~ 2.5 m Liquid hydrogen target
- ~ 4 m Recoil proton detector (**CAMERA**)
- New large angle em. calorimeter in front of SM1 (**ECAL0**)



- μ^+ and μ^- beams
- momentum: 100 – 190 GeV/c
- polarization: 80 %
opposite for μ^+ and μ^-
- coverage of intermediate x_{Bj}
 \rightsquigarrow unexplored region between
 ZEUS+H1 and HERMES+JLab




$\mu p \rightarrow \mu' p' \gamma$: Interference with Bethe-Heitler



both processes interfere on level of amplitudes

$$d\sigma_{(\mu p \rightarrow \mu' p' \gamma)} \propto |T^{BH}|^2 + \text{Interference Term} + |T^{DVCS}|^2$$


$\mu p \rightarrow \mu' p' \gamma$: Interference with Bethe-Heitler



both processes interfere on level of amplitudes

$$d\sigma(\mu p \rightarrow \mu' p' \gamma) \propto |T^{BH}|^2 + \text{Interference Term} + |T^{DVCS}|^2$$


$0.005 < x_{Bj} < 0.01$



BH dominates


~> reference yield

$0.01 < x_{Bj} < 0.03$



Interference

$x_{Bj} > 0.03$




DVCS dominates

MC simulation
for COMPASS
without ECALO


Beam Tests @ COMPASS during hadron programme:

- 2008 (8 hours)
- 2009 (10 times statistics of 2008)



Target Setup for the Hadron Programme (2008-2009):


- Target: 40 cm LH₂
- Recoil Detector (1 m long)
- ECAL1 & ECAL2




~> Observation of exclusive single photon production

- $\epsilon_{\mu p \rightarrow \mu' p' \gamma} = 0.32 \pm 0.13$
- SPS & COMPASS availability
- DAQ dead time
- Trigger efficiency

global efficiency: $\epsilon_{global} = 0.13 \pm 0.05$






- confirmation of global efficiency






- confirmation of global efficiency

↪ realistic projection of errors



$$\begin{aligned}
 d\sigma_{(\mu p \rightarrow \mu' p' \gamma)} &= d\sigma^{BH} + d\sigma_{unpol}^{DVCS} + P_{\mu} d\sigma_{pol}^{DVCS} \\
 &+ e_{\mu} a^{BH} \text{Re}(T^{DVCS}) + e_{\mu} P_{\mu} a^{BH} \text{Im}(T^{DVCS})
 \end{aligned}$$






$$d\sigma_{(\mu p \rightarrow \mu' p' \gamma)} = d\sigma^{BH} + d\sigma_{unpol}^{DVCS} + P_{\mu} d\sigma_{pol}^{DVCS} + e_{\mu} a^{BH} \text{Re}(T^{DVCS}) + e_{\mu} P_{\mu} a^{BH} \text{Im}(T^{DVCS})$$

- Beam charge and Spin **sum**:

$$\mathcal{S}_{CS,U} = d\sigma^{+\leftarrow} + d\sigma^{-\rightarrow} = 2 (d\sigma^{BH} + d\sigma_{unpol}^{DVCS} + e_{\mu} P_{\mu} a^{BH} \text{Im}(T^{DVCS}))$$

- ϕ dependence gives access to GPD H

$$\propto \sin \phi$$



$$d\sigma_{(\mu p \rightarrow \mu' p' \gamma)} = d\sigma^{BH} + d\sigma_{unpol}^{DVCS} + P_\mu d\sigma_{pol}^{DVCS} + e_\mu a^{BH} \text{Re}(T^{DVCS}) + e_\mu P_\mu a^{BH} \text{Im}(T^{DVCS})$$


- Beam charge and Spin **sum**:

$$S_{CS,U} = d\sigma^{+\leftarrow} + d\sigma^{-\rightarrow} = 2 (d\sigma^{BH} + d\sigma_{unpol}^{DVCS} + e_\mu P_\mu a^{BH} \text{Im}(T^{DVCS}))$$


- ϕ dependence gives access to GPD H
- Integration over ϕ and subtracting BH:

$$d\sigma^{DVCS}/dt \sim \exp(-B|t|)$$

$$\propto \sin \phi$$




(a)

(b) $x \sim 0.003$ $x \sim 0.03$ $x \sim 0.3$

- Integration over ϕ and subtracting BH:

$$d\sigma^{DVCS}/dt \sim \exp(-B|t|)$$

$$B(x_B) \sim \frac{1}{2} \langle r_{\perp}^2(x_B) \rangle$$



Input for projections:

- $L = 1222 \text{ pb}^{-1}$
- 2 years of data
- $\epsilon_{global} = 10\%$
- 160 GeV/c muon beam
- 2.5 m LH_2 target

- Integration over ϕ and subtracting BH:

$$d\sigma^{DVCS}/dt \sim \exp(-B|t|)$$

$$B(x_B) \sim \frac{1}{2} \langle r_{\perp}^2(x_B) \rangle$$




- Ansatz at small x_B : ($x \sim x_B$)

$$B(x_B) = b_0 + 2\alpha' \ln(x_0/x_B)$$




Accuracy $> 2.5\sigma$

for: $\alpha' > 0.26$ (with ECAL 1+2)

for: $\alpha' > 0.125$ (with ECAL 0+1+2)



$$\begin{aligned}
 d\sigma_{(\mu p \rightarrow \mu' p' \gamma)} &= d\sigma^{BH} + d\sigma_{unpol}^{DVCS} + P_{\mu} d\sigma_{pol}^{DVCS} \\
 &+ e_{\mu} a^{BH} \text{Re}(T^{DVCS}) + e_{\mu} P_{\mu} a^{BH} \text{Im}(T^{DVCS})
 \end{aligned}$$






$$d\sigma_{(\mu p \rightarrow \mu' p' \gamma)} = d\sigma^{BH} + d\sigma_{unpol}^{DVCS} + P_\mu d\sigma_{pol}^{DVCS} + e_\mu a^{BH} \text{Re}(T^{DVCS}) + e_\mu P_\mu a^{BH} \text{Im}(T^{DVCS})$$

- Beam charge and Spin difference:

$$D_{CS,U} = d\sigma^{+\leftarrow} - d\sigma^{-\rightarrow} = 2 (P_\mu d\sigma_{pol}^{DVCS} + e_\mu a^{BH} \text{Re}(T^{DVCS}))$$

⇒ BH contribution cancels

↪ control detector acceptance and beam flux with high precision



$$d\sigma_{(\mu p \rightarrow \mu' p' \gamma)} = d\sigma^{BH} + d\sigma_{unpol}^{DVCS} + P_\mu d\sigma_{pol}^{DVCS} + e_\mu a^{BH} \text{Re}(T^{DVCS}) + e_\mu P_\mu a^{BH} \text{Im}(T^{DVCS})$$

- Beam charge and Spin difference:


$$D_{CS,U} = d\sigma^{+\leftarrow} - d\sigma^{-\rightarrow} = 2 \left(P_\mu d\sigma_{pol}^{DVCS} + e_\mu a^{BH} \text{Re}(T^{DVCS}) \right)$$

⇒ BH contribution cancels

↪ control detector acceptance and beam flux with high precision

$$\propto c_0^{Int} + c_1^{Int} \cos \phi$$


$$c_{0,1}^{Int} \propto \text{Re}(F_1 \mathcal{H})$$



Input for projections:

- $L = 1222 \text{ pb}^{-1}$
- 2 years of data
- $\epsilon_{global} = 10\%$
- 160 GeV/c muon beam
- 2.5 m LH₂ target

$$\text{Re}(\mathcal{H}(\xi, t)) = \sum_f e_f^2 \left[\mathcal{P} \int dx H^f(x, \xi, t) \left(\frac{1}{x-\xi} \mp \frac{1}{x+\xi} \right) \right]$$



Cross section measurements:


- Pseudo-scalar: $\pi, \eta, \dots \Rightarrow \tilde{H}$
- Vector meson: $\rho, \omega, \phi, \dots \Rightarrow H$

Allow for flavour separation:

$$H_{\rho^0} = \frac{1}{\sqrt{2}} \left(\frac{2}{3} H^u + \frac{1}{3} H^d + \frac{3}{8} H^g \right)$$

$$H_{\omega} = \frac{1}{\sqrt{2}} \left(\frac{2}{3} H^u - \frac{1}{3} H^d + \frac{1}{8} H^g \right)$$

$$H_{\phi} = -\frac{1}{3} H^s - \frac{1}{8} H^g$$



Cross section measurements:

- Pseudo-scalar: $\pi, \eta, \dots \Rightarrow \tilde{H} \text{ \& \; } \tilde{E}$
- Vector meson: $\rho, \omega, \phi, \dots \Rightarrow H \text{ \& \; } E$

Vector meson production from transversely polarized target:


- Asymmetry $\propto E/H$

Allow for flavour separation:

$$H_{\rho^0} = \frac{1}{\sqrt{2}} \left(\frac{2}{3} H^u + \frac{1}{3} H^d + \frac{3}{8} H^g \right)$$

$$H_{\omega} = \frac{1}{\sqrt{2}} \left(\frac{2}{3} H^u - \frac{1}{3} H^d + \frac{1}{8} H^g \right)$$

$$H_{\phi} = -\frac{1}{3} H^s - \frac{1}{8} H^g$$



Cross section measurements:

- Pseudo-scalar: $\pi, \eta, \dots \Rightarrow \tilde{H} \text{ \& \; } \tilde{E}$
- Vector meson: $\rho, \omega, \phi, \dots \Rightarrow H \text{ \& \; } E$

Vector meson production from transversely polarized target:

- Asymmetry $\propto E/H$

Allow for flavour separation:

$$H_{\rho^0} = \frac{1}{\sqrt{2}} \left(\frac{2}{3} H^u + \frac{1}{3} H^d + \frac{3}{8} H^g \right)$$

$$H_{\omega} = \frac{1}{\sqrt{2}} \left(\frac{2}{3} H^u - \frac{1}{3} H^d + \frac{1}{8} H^g \right)$$

$$H_{\phi} = -\frac{1}{3} H^s - \frac{1}{8} H^g$$

Also studied without RPD

used data with transversely polarized targets:

${}^6\text{LiD}$: 2002-2004

NH_3 : 2007 & 2010


HEMP: Transversely Polarized Target

Transverse target spin asymmetry: $A_{UT}^{\sin \phi - \phi_S}$




$$A_{UT}^{\sin \phi - \phi_S}(\rho^0) \propto \sqrt{|-t'|} \frac{\text{Im}(\mathcal{E}^* \mathcal{H})}{|\mathcal{H}|^2}$$

- \mathcal{E} and \mathcal{H} are weighted sums of GPDs $E^{q,g}$ and $H^{q,g}$

\rightsquigarrow provide access to GPD E



Goloskokov and Kroll
Eur. Phys. J. C 59 4 (2009)





- COMPASS-II will investigate quark GPDs with DVCS
 - Covered x_B/Q^2 regime not accessible to any other experiment in the near future
 - Change of beam charge and polarization - **UNIQUE**
 - Study nucleon transversal dimension as function of x_B (Tomography)
 - Constrain GPD H through ϕ dependence of $\mathcal{D}_{CS,U}$ and $\mathcal{S}_{CS,U}$
- Complementary information from hard exclusive meson production
- In a second phase measurement with transversely polarized target and RPD



Thank You!



Back Up



- 1 Motivation
- 2 COMPASS Experiment
- 3 DVCS: Testrun 2008 and 2009
- 4 Deeply Virtual Compton Scattering
- 5 Hard Exclusive Meson Production
- 6 Summary
- 7 Back Up
- 8 Table of Content