

COMPASS: present and future



Barbara Badelek
University of Warsaw

Low x 2012

Paphos, June 27 – 30, 2012

What has COMPASS done so far




- Is taking data since 2002 (2005 break due to SPS shutdown);
- muon and hadron programmes;
- longitudinally polarised muon beam of $160 \text{ GeV}/c$
off longitudinally and transversely polarised targets: ${}^6\text{LiD}$ (d), NH_3 (p);
- $190 \text{ GeV}/c$ hadron beams: π , K, p
off unpolarised targets: liquid H_2 , Pb, Ni, Cu, W;
- originally planned until 2009;
- addendum 2010-2011: transverse and longitudinal NH_3 ;
- **muon results:** quarks contribute 30% to the nucleon's spin,
gluons contribution small in the measured x range,
all 3 leading twist PDF (f_1, g_1, h_1) investigated
- **hadron results:** search for exotics, hybrids, glueballs;
measurements of pion polarisabilities.



COMPASS Spectrometer (muon run)

Nucl. Instr. Meth. A577 (2007) 455

Two stages, ~ 350 planes
Calorimetry
Particle identification
Large, solid state polarised target




COMPASS polarised targets




- * Two (three) target cells, oppositely polarised
- * Polarisation reversed every 8 h (less frequent after 2005) by field rotation
- * Material: solid $^6\text{LiD}(\text{NH}_3)$
- * Polarisation: $\sim 50\%$ ($\sim 90\%$), by the Dynamical Nuclear Polarisation
- * Dilution: $f \sim 0.4$ (~ 0.15)
- * Polar acceptance: ~ 70 mrad (~ 180 mrad after 2005)

COMPASS RICH



Before upgrade: white distribution
After upgrade: yellow distribution



RICH2007

Federica Sozzi



19

COMPASS II Proposal

- CERN–SPSC–2010–014 (SPSC–P–340) of May 17, 2010
www.compass.cern.ch/compass/proposal/compass-II_proposal/compass-II_proposal.pdf
- Approved in December 2010 initially 3 years data taking (**Phase 1**)
- **Flavour separation and fragmentation in spin-averaged SIDIS**
(strange sector !)
- Focus on transverse structure of the nucleon
 - **GPD**, transverse size and parton orbital angular momentum
 - **T-odd TMD** (Sivers, Boer-Mulders distributions)
 - **Drell-Yan** process and TMD sign change SIDIS \iff DY
- **π/K polarisabilities and tests of ChPT**
in the Compton scattering via Primakoff reaction.
- Addendum foreseen (spin-dependent GPD), **Phase 2**.


Spin-averaged SIDIS

- Spin-averaged SIDIS, $\mu p \rightarrow \mu h X$, will be recorded simultaneously with DVCS/DVMP, on a long LH₂ target (NH₃ target from COMPASS I is difficult!)
- Charge and nature of “h” will be determined (π^\pm , π^0 , K^\pm , K^0 , Λ , $\bar{\Lambda}$)
- Combined with COMPASS I data on ${}^6\text{LiD} \implies$ q_f separation in global QCD fits, constraints on FF (hadron multiplicities) and LO determination of $s(x)$ at $0.001 < x < 0.2$.
- Two years of data taking: dependence on x , Q^2 , p_T^2 , z and asymmetries $A^{\cos \phi}$, $A^{\cos 2\phi}$, $A^{\sin \phi}$ of hadrons on unpol. proton target. They are sensitive to T-odd TMD Boer-Mulders function and Cahn effect. Data exist for ${}^6\text{LiD}$.




3D picturing of the proton *via* GPD


D. Mueller, X. Ji, A. Radyushkin, A. Belitsky, ...
M. Burkardt, ... Interpretation in impact parameter space



Proton form factors,
transverse charge &
current densities




Correlated quark momentum
and helicity distributions in
transverse space - **GPDs**



Structure functions,
quark **longitudinal**
momentum & helicity
distributions

Access GPD through the DVCS/DVMP mechanism




$Q^2 \rightarrow \infty$,
fixed $x_B, t \implies |t|/Q^2$ small

- 4 GDPs ($H, E, \tilde{H}, \tilde{E}$) for each flavour and for gluons
- Factorisation proven for σ_L only
- All depend on 4 variables: x, ξ, t, Q^2 ; DIS @ $\xi = t = 0$;
Later Q^2 dependence omitted. **Careful! Here $x \neq x_B$!**
- H, \tilde{H} conserve nucleon helicity
 E, \tilde{E} flip nucleon helicity
- $\underline{H}, \underline{E}$ refer to unpolarised distributions
 \tilde{H}, \tilde{E} refer to polarised distributions
- $H^q(x, 0, 0) = q(x), \tilde{H}^q(x, 0, 0) = \Delta q(x)$



- $\underline{H}, \underline{E}$ accessed in vector meson production *via* A_{UT} asymmetries
- \tilde{H}, \tilde{E} accessed in pseudoscalar meson production *via* A_{UT} asymmetries
- All 4 accessed in DVCS (γ production) in $A_C, A_{LU}, A_{UT}, A_{UL}$
- Integrals of $H, E, \tilde{H}, \tilde{E}$ over x give Dirac-, Pauli-, axial vector- and pseudoscalar vector form factors respectively.
- **Important:** $J_z^q = \frac{1}{2} \int dx x [H^q(x, \xi, t=0) + E^q(x, \xi, t=0)] = \frac{1}{2} \Delta \Sigma + L_z^q$ (X. Ji)

Why GPD at COMPASS ?



- CERN high energy muon beam
 - 100 - 190 GeV
 - 80% polarisation, opposite for μ^+ , μ^-
 - $\mu^+ \leftarrow$ and $\mu^- \rightarrow$ beams
- Kinematic range
 - between HERA and HERMES/JLab12
 - intermediate x (sea and valence)
- Separation
 - pure B-H @ low x_B
 - predominant DVCS @ high x_B
- Plans
 - DVCS
 - DVMP
- Goals
 - from unpolarised target: H (Phase 1)
 - from \perp polarised target: E (Phase 2)

DVCS/DVMP: $\mu p \rightarrow \mu p \gamma$ (M); what do we measure?



$$d\sigma^{\mu p \rightarrow \mu p \gamma} = d\sigma^{BH} + (d\sigma_{unpol}^{DVCS} + P_\mu d\sigma_{pol}^{DVCS}) + e_\mu (\text{Re}I + P_\mu \text{Im}I) \quad 4 - \text{dim.} : x_B, Q^2, t, \phi$$

Observables (Phase 1):

$$\bullet S_{CS,U} \equiv \mu^{+\leftarrow} + \mu^{-\rightarrow} = 2 \left(d\sigma^{BH} + d\sigma_{unpol}^{DVCS} + e_\mu P_\mu \text{Im}I \right)$$


$$\bullet D_{CS,U} \equiv \mu^{+\leftarrow} - \mu^{-\rightarrow} = 2 \left(P_\mu d\sigma_{pol}^{DVCS} + e_\mu \text{Re}I \right)$$

$$\bullet A_{CS,U} \equiv \frac{\mu^{+\leftarrow} - \mu^{-\rightarrow}}{\mu^{+\leftarrow} + \mu^{-\rightarrow}} = \frac{D_{CS,U}}{S_{CS,U}}$$

• Each term ϕ -modulated

If ϕ -dependence integrated over \implies twist-2 DVCS contribution;

if ϕ -dependence analysed: $\implies \text{Im}(F_1H)$ and $\text{Re}(F_1H)$




Analogously for transversely polarised target (Phase 2): $S_{CS,T}, D_{CS,T}, A_{CS,T} \implies E$



DVCS/DVMP: experimental requirements

New hardware items needed:

- 2.5 m long liquid H₂ target (LH₂), ready 2012
- 4 m long, tof recoil proton detector (CAMERA), ready 2012, with 2 barrels and a readout of 1GHz digitalisation (Gandalf)
- large angle electromagnetic calorimeter (ECAL0) just downstream target; prototype ready 2012




Simulations @ 160 GeV and present setup (no ECAL0); normalisation of BH @ small x_B



DVCS tests in 2008/2009

- Data taken at 160 GeV with μ^+ and μ^- (flux (μ^+)/flux (μ^-) = 3 at this energy)
- 40 cm long H_2 target (one vertex with μ, μ')
- Short recoil proton detector (one proton, $p_p < 1$ GeV/c)
- No ECAL0 (only one γ)

Results for μ^+ beam:




- MC normalised to yield @ low x_B
- A clear signal of DVCS at $x_B > 0.03$



In 2012 ~ 10 x larger statistics!

DVCS: projected data sets and results

- Simulations for:
 - 2 years of data taking
 - 10% global efficiency
 - $L = 1222 \text{ pb}^{-1}$




- $S_{CS,U}, D_{CS,U}, A_{CS,U}$ measured in $6 x_B \times 4 Q^2$ bins as function of ϕ
- Azimuthal dependence $A_{CS,U}$ and $D_{CS,U}$ compared to models:



Nucleon transverse structure (“tomography”)

In the GPD limiting case $\xi = 0$, $t \equiv -\Delta^2 = -\Delta_{\perp}^2$ and

$$q^f(x, \mathbf{b}_{\perp}) = \int \frac{d^2\Delta_{\perp}}{(2\pi)^2} e^{-i\Delta_{\perp} \cdot \mathbf{b}_{\perp}} H^f(x, 0, -\Delta_{\perp}^2)$$




Get the nucleon transverse size as a function of longitudinal momentum fraction

Nucleon transverse structure – projected results


Case $x = \xi$

$$\frac{d\sigma_{\text{unpol}}^{\text{DVCS}}}{dt} \propto e^{-B(x_B)|t|} \quad \text{where at low } x_B : B(x_B) \approx \frac{1}{2} \langle r_{\perp}^2(x_B) \rangle$$




Here a simple ansatz: $B(x_B) = B_0 + 2\alpha' \log \frac{x_0}{x_B}$ was assumed.


Transverse Momentum Dependent (TMD) distributions



- parton intrinsic k_T taken into account
- allow accessing quark $L!$
- at COMPASS studied in 2 ways:
 - semi-inclusive DIS (polarised muons on unpolarised/transversely polarised target)
 - Drell-Yan process (pion beam on unpolarised/transversely polarised target)



SIDIS



DY


From Cédric Lorcé (Orsay), IWHSS2012

DVCS vs. SIDIS

Incoherent scattering




DVCS



$$x = \frac{k^+}{P^+} \quad -2\xi = \frac{\Delta^+}{P^+} \quad t = \Delta^2$$

SIDIS



Factorization

Compton form factor
Cross section

=

hard

μ_F

\otimes

soft

- process dependent
- perturbative

- « universal »
- non-perturbative


TMD distributions...cont'd

- In LT and considering k_T , 8 PDF describe the nucleon
- QCD-TMD approach valid $k_T \ll \sqrt{Q^2}$
- After integrating over k_T only 3 survive: f_1, g_1, h_1
- TMD accessed in SIDIS and DY by measuring azimuthal asymmetries
- SIDIS: e.g. $A_{\text{Sivers}} \propto \text{PDF} \otimes \text{FF}$
- DY: e.g. $A_{\text{Sivers}} \propto \text{PDF}^{\text{beam}} \otimes \text{PDF}^{\text{target}}$
- OBS! Boer-Mulders and Sivers PDF are T-odd, i.e. process dependent

$$h_1^\perp(\text{SIDIS}) = -h_1^\perp(\text{DY})$$

$$f_{1T}^\perp(\text{SIDIS}) = -f_{1T}^\perp(\text{DY})$$

- OBS! transversity PDF is chiral-odd
- Boer-Mulders, Sivers and transversity ($h_1^\perp, f_{1T}^\perp, h_1$) will be measured in COMPASS II




Boer-Mulders (h_1^\perp) and Sivers (f_{1T}^\perp) DF in SIDIS

- Boer–Mulders asymmetry,

$$A_{\cos 2\phi} \equiv A_{LU}^{\cos 2\phi},$$



was found $\neq 0$ on deuteron

- can be measured on the proton together with DVCS





- Sivers asymmetry, $A_{Siv}^p \equiv A_{LT}^{\sin \Phi_S}$, measured on d and p targets

- Found positive for h^+ on proton but less strong than in HERMES




Drell-Yan process: $\pi^- p^\uparrow \rightarrow \mu^+ \mu^- X$ @ COMPASS

- Clean partonic process
- No fragmentation functions involved!
- Convolution of two Parton Distribution Functions
 $\sigma^{\text{DY}} \propto f_{\bar{u}|\pi^-} \otimes f'_{u|p}$, $\sigma^{\text{DY}} = \sigma^{\text{DY}}(x_\pi, x_p)$
- Gives an access to azimuthal modulations of 4 PDF: transversity, pretzelosity, Boer–Mulders and Sivers.
- Ideal: $\bar{p}p$; good compromise: $\pi^- p$
- Here dominated by annihilation of valence \bar{u} from π^- and valence u from p
- COMPASS has large acceptance in the valence region of p and π (large SSA expected).
 Example of covered kinematics (in blue):
 π^- beam, 190 GeV/c, NH_3 target, \perp polarised dimuon mass range: 4 – 9 GeV/c² (low bckg.)
- QCD TMD approach justified by:
 $M_{\mu\mu} \gg p_T^{\mu\mu} \approx 1 \text{ GeV}$




Drell-Yan @ COMPASS: experimental requirements

- intense π beam (up to 10^9 /spill)
- hadron absorber downstream target
- di-muon trigger








Drell-Yan @ COMPASS: Results from 2009 beam test



- 3 days of data taking
- $8 \cdot 10^7 \pi^- / 9.6 \text{ s spill}$
- 2 cells of CH_2 of 40-20-40 cm
- temporary absorber
- simple trigger

Drell–Yan @ COMPASS: Projections for azimuthal asymmetries

- projections for $4 \text{ GeV}/c^2 < M_{\mu\mu} < 9 \text{ GeV}/c^2$
- 2 years of data taking
- $6 \cdot 10^8$ pions /9.6 s spill
- 1.1 m long, polarised NH_3 target





Goals: change of sign between DIS and SIDIS in h_1^\perp ,
 J/Ψ production mechanism,...

Future Drell–Yan experiments

Facility	Type	s (GeV ²)	Time-line
RHIC (STAR, PHENIX)	collider, $p^\uparrow p^\uparrow$	$200^2, 500^2$	> 2014
RHIC (internal target)	fixed target, $p^\uparrow p^\uparrow$	500	> 2015
RHIC (AnDY)	collider, $p^\uparrow p^\uparrow$	500^2	<i>cancelled</i>
E906 (Fermilab)	fixed target, pp	226	> 2010
J-PARC	fixed target, pp^\uparrow	$60 \div 100$	> 2015
GSI (PAX)	collider, $\bar{p}^\uparrow p^\uparrow$	200	> 2017
GSI (Panda)	fixed target, $\bar{p}p$	30	> 2016
NICA	collider, $p^\uparrow p^\uparrow, d^\uparrow d^\uparrow$	676	> 2014
COMPASS II	fixed target, $\pi^- p^\uparrow$	$300 \div 400$	> 2012


Studies of Chiral Perturbative Theory

Primakoff ($\pi^- Z \rightarrow \pi^- Z \gamma$) and Compton ($\pi \gamma \rightarrow \pi \gamma$) processes



- Breaking of chiral symmetry \implies Goldstone bosons (pions, kaons)
- ChPT predicts *e.g.* pion electromagnetic polarisabilities
 \implies deviations of $\sigma(\pi\gamma \rightarrow \pi\gamma)$ from QED from point-like, spin 0 object


$$\frac{d\sigma_{\pi\gamma}}{d\Omega_{\text{cm}}} = \left[\frac{d\sigma_{\pi\gamma}}{d\Omega_{\text{cm}}} \right]_{\text{point-like}} + C \frac{s - m_\pi^2}{s^2} P(\alpha_\pi, \beta_\pi)$$



$$P(\alpha_\pi, \beta_\pi) = (1 - \cos \theta_{\text{cm}})^2 (\alpha_\pi - \beta_\pi) + (1 + \cos \theta_{\text{cm}})^2 (\alpha_\pi + \beta_\pi) f_1(s) + (1 - \cos \theta_{\text{cm}})^3 (\alpha_\pi - \beta_\pi) f_2(s)$$

Studies of Chiral Perturbative Theory,...cont'd

- 2-loop ChPT prediction: $\alpha_\pi - \beta_\pi = (5.7 \pm 1.0) \cdot 10^{-4} \text{ fm}^3$
- Measurements: $\alpha_\pi - \beta_\pi = (4 - 14) \cdot 10^{-4} \text{ fm}^3$
- COMPASS II: measurements of both **pion and kaon polarisabilities!**



In 120 days (90 with π , 30 with μ beams)	$\alpha_\pi - \beta_\pi$ (10^{-4} fm^3)	$\alpha_\pi + \beta_\pi$ (10^{-4} fm^3)	$\alpha_2 - \beta_2$ (10^{-4} fm^5)
2-loop ChPT prediction	5.7 ± 1.0	0.16 ± 0.10	16
COMPASS sensitivity	± 0.66	± 0.025	± 1.94

Tentative time table for COMPASS II

Proposal: CERN-SPSC-2010-014 (SPSC-P-340) of May 17, 2010
Approved in December 2010, initially 3 years data taking, 2014-16 (Phase 1)
www.compass.cern.ch/compass/proposal/compass-II_proposal/compass-II_proposal.pdf

- **2012 setup and tests:** Primakoff with π , K beams, 8 weeks
 \implies test of χ PT;
 DVCS with μ^+ , μ^- beams on unpolarised protons, 4 weeks;
- **2013 SPS shutdown;** polarised target installation;
- **2014 Drell-Yan** with π beam \implies TMD;
- **2015 DVCS** with μ^+ , μ^- beams on unpolarised protons,
– **2016** \implies constrain GPD H and t -slope parameter B .
 Parallely SIDIS \implies PDF, TMD, FF (especially for s-quark).
- **≥ 2017 Addendum ???** DVCS with μ^+ , μ^- beams on \perp polarised protons
 \implies constrain GPD E (Phase 2).