

OZI rule violation in vector meson production at COMPASS

HK23.5 DPG Frühjahrstagung Mainz

Johannes Bernhard¹

Institut für Kernphysik Mainz

for the COMPASS collaboration

March 21th 2011



bmb+f - Förderschwerpunkt

COMPASS

Großgeräte der physikalischen
Grundlagenforschung

¹Contact: johannes.bernhard@cern.ch


Motivation

Okubo-Zweig-Iizuka rule¹: processes with disconnected quark lines suppressed
 $\phi(1020)$ to $\omega(782)$ production ratios (A and B non-strange hadrons), not corrected for phase-space²:

$$\sigma(AB \rightarrow \phi X) / \sigma(AB \rightarrow \omega X) \simeq \tan^2(\theta - \theta_0) \simeq 4.2 \cdot 10^{-3}$$

Numerous violations observed, possible explanations:



- reactions on nucleons: $s\bar{s}$ production due to strangeness content of proton
- intermediate (gluon-rich) states



¹S. Okubo, Phys. Lett. 5(1963)165, G. Zweig, CERN report TH-401(1964), J. Iizuka, Prog.Theor.Suppl.38(1966)21

²H.J. Lipkin, Phys. Lett. B 60 (1976) 371

Violations of the OZI rule / COMPASS



A. Sibirtsev and W. Cassing, Eur.Phys.J.A7(2000)407

No data available for higher energies


Study at COMPASS:

Compare $\phi(1020) \rightarrow K^+K^-$ to $\omega(782) \rightarrow \pi^+\pi^-\pi^0$ production

The COMPASS spectrometer at CERN

190 GeV/c π^\pm /K $^\pm$ /p beam

2 stage high resolution spectrometer
with large acceptance





beam PID with CEDAR detectors
particle ID with RICH and Calorimetry


Event selection

Interest in $p p \rightarrow p (\pi^+ \pi^- \pi^0) / (K^+ K^-) p$ final states


- select event topology (charged tracks, reaction inside target volume, recoil proton etc.)
- ID K^+ with RICH, π^0 with ECALs
- conservation of charge, exclusivity




Reaction Kinematics



Invariant mass distributions ($K^+ K^-$)




Invariant mass distributions ($\pi^+ \pi^- \pi^0$)




Test OZI violation: Analysis


$0.6 < x_F < 0.7$






$0.7 < x_F < 0.8$



$0.8 < x_F < 0.9$




ω



- ① fit invariant mass distributions with Breit-Wigner folded with Gaussian plus polynomial background in x_F bins \Rightarrow yields
- ② correct for acceptance and branching \Rightarrow corrected yields
- ③ calculate $R = \frac{\text{Number of } \phi}{\text{Number of } \omega}$

Test OZI violation: Result



N.B.: Included only systematics from fit and ECAL reconstruction, additional effects are still under investigation

Outlook and Conclusions


Preliminary results from 2008 proton campaign (one week):
OZI violation of a factor 3 at 190 GeV beam energy

Ongoing:

- 2009 data sample ($> 5x$ statistics of 2008)
- OZI tests w.r.t. t'
- Improved Monte-Carlo (multi-dim. acceptance)
- ω/ϕ spin alignment \Rightarrow production mechanisms


Spares

Exclusivity



Selection of exclusive events: energy balance $191 \text{ GeV} \pm 6 \text{ GeV}$


Production mechanism



Ongoing: binning the ratio $R = \frac{\text{Number of } \phi \text{ in } t}{\text{Number of } \omega \text{ in } t}$

Background

Composition 2008 data sample: exclusive vs. non-exclusive



Important for background studies