

Likelihood Methods for Beam Particle Identification at the COMPASS Experiment

Tobias Weisrock
for the COMPASS Collaboration

Graduate School Symmetry Breaking
Johannes Gutenberg-Universität Mainz

SPIN-Praha-2012
July 7, 2012



bmb+f - Förderschwerpunkt

COMPASS

Großgeräte der physikalischen
Grundlagenforschung



JOHANNES GUTENBERG
UNIVERSITÄT MAINZ

Symmetry
→
Breaking



Outline

The COMPASS Experiment

Hadron Physics @ COMPASS


Beam Particle Identification @ COMPASS

Beam Particle Identification using Likelihoods

Efficiency and Purity of the Likelihood Method


The COMPASS Experiment

- ▶ **CO**mmun **M**uon and **P**roton **A**pparatus for **S**tructure and **S**pectroscopy
- ▶ Located at SPS at CERN




The COMPASS Experiment

- ▶ **CO**mmun **M**uon and **P**roton **A**pparatus for **S**tructure and **S**pectroscopy
- ▶ Located at SPS at CERN




The COMPASS Experiment

- ▶ **CO**mmun **M**uon and **P**roton **A**pparatus for **S**tructure and **S**pectroscopy
- ▶ Located at SPS at CERN




The COMPASS Experiment

- ▶ **CO**mmun **M**uon and **P**roton **A**pparatus for **S**tructure and **S**pectroscopy
- ▶ Located at SPS at CERN



The COMPASS Experiment

- ▶ **CO**mmun **M**uon and **P**roton **A**pparatus for **S**tructure and **S**pectroscopy
- ▶ Located at SPS at CERN



Hadron Physics @ COMPASS

- ▶ Very broad program



Hadron Physics @ COMPASS

- ▶ Very broad program
- ▶ Spectroscopy
 - ▶ Different final states
 - ▶ 3π , 5π , $K\pi\pi$, . . .
 - ▶ Search for exotic states



Hadron Physics @ COMPASS

- ▶ Very broad program
- ▶ Spectroscopy
 - ▶ Different final states
 - ▶ $3\pi, 5\pi, K\pi\pi, \dots$
 - ▶ Search for exotic states
- ▶ Primakoff program
 - ▶ $\pi p \rightarrow \pi\gamma p$
 - ▶ Measure pion/kaon polarizabilities



Hadron Physics @ COMPASS

- ▶ Very broad program
- ▶ Spectroscopy
 - ▶ Different final states
 - ▶ $3\pi, 5\pi, \mathbf{K}\pi\pi, \dots$
 - ▶ Search for exotic states
- ▶ Primakoff program
 - ▶ $\pi p \rightarrow \pi\gamma p$
 - ▶ Measure pion/kaon polarizabilities



Example: $\mathbf{K}\pi\pi$ analysis

Analysis of diffractive dissociation of \mathbf{K}^- into $\mathbf{K}^-\pi^+\pi^-$ on a liquid hydrogen target at the COMPASS spectrometer, PhD Thesis P. Jasinski (JGU Mainz)




Event Selection

- ▶ Incoming kaon
- ▶ Primary vertex in target with 3 outgoing charged tracks
- ▶ Four momentum transfer $0.07 < t'[\text{GeV}^2/c^2] < 0.7$
- ▶ One negative particle identified with RICH




Partial Wave Analysis

- ▶ disentangle $K\pi\pi$ spectrum
- ▶ several resonances found
- ▶ further investigation needed (mass dependent fits)



Partial Wave Analysis


- ▶ disentangle $K\pi\pi$ spectrum
- ▶ several resonances found
- ▶ further investigation needed (mass dependent fits)



J^P	mass (GeV/c ²)	possible state
0^-	1.30	$K(1460)$
1^+	1.25	$K_1(1270)$
1^+	1.35	$K_1(1400)$
1^+	1.80	$K_1(1650)$
1^-	1.75	$K^*(1680)$
2^+	1.44	$K_2^*(1430)$
2^-	1.70	$K_2(1770)$
2^-	1.85	$K_2(1820)$
2^-	1.9 – 2.2	several

Partial Wave Analysis

- ▶ disentangle $K\pi\pi$ spectrum
- ▶ several resonances found
- ▶ further investigation needed (mass dependent fits)



J^P	mass (GeV/c ²)	possible state
0^-	1.30	K(1460)
1^+	1.25	K₁(1270)
1^+	1.35	K₁(1400)
1^+	1.80	K₁(1650)
1^-	1.75	K*(1680)
2^+	1.44	K₂*(1430)
2^-	1.70	K₂(1770)
2^-	1.85	K₂(1820)
2^-	1.9 – 2.2	several

Only $\approx 50\%$ of kaons used due to bad efficiency of beam particle identification.


Beam Particle Identification @ COMPASS

- ▶ Two CEDAR detectors
- ▶ **30 m** upstream of the target




How does a CEDAR work?

- ▶ CEDAR = Čerenkov Differential counters with Acromatic Ring focus
- ▶ Fast charged particles emit Čerenkov light with angle $\cos(\theta) = \frac{1}{n\beta}$



How does a CEDAR work?

- ▶ Čerenkov light detected with 8 PMTs
- ▶ Particle identification using multiplicities, e.g. 6 of 8 PMTs




Influence of Beam Divergence

- ▶ beam tuned to traverse full spectrometer
- ▶ beam is not parallel in the CEDAR region


Influence of Beam Divergence

- ▶ beam tuned to traverse full spectrometer
- ▶ beam is not parallel in the CEDAR region



Influence of Beam Divergence


- ▶ beam tuned to traverse full spectrometer
- ▶ beam is not parallel in the CEDAR region



- ▶ Kaon ring leaves acceptance, pion ring enters
- ▶ Multiplicity method does not work for divergent beams

Influence of Beam Divergence

- ▶ beam tuned to traverse full spectrometer
- ▶ beam is not parallel in the CEDAR region



- ▶ Kaon ring leaves acceptance, pion ring enters
- ▶ Multiplicity method does not work for divergent beams
- ↪ Find a method which takes divergence into account

General Idea

- ▶ Look at response of single PMTs for kaon and pion separately
- ▶ Take care of beam divergence
- ▶ Identify beam particles using likelihoods



General Idea

- ▶ Look at response of single PMTs for kaon and pion separately
- ▶ Take care of beam divergence
- ▶ Identify beam particles using likelihoods




5 steps to take

1. Measure beam divergence
2. Create a pure kaonsample and a pure pionsample
3. Determine probabilities to have hits in PMTs for pion and kaon
4. Calculate likelihoods from probabilities
5. Use likelihoods to identify particles



Step 1: Measure beam divergence

- ▶ Measure beam position in front of (x_1, y_1) and behind (x_2, y_2) CEDARs
- ▶ Calculate relative displacement $\Delta_x = \frac{x_2 - x_1}{1283,4 \text{ cm}}$
- ▶ Divergence $\theta_x = \arctan(\Delta_x) \approx \Delta_x$



Step 2: Create a pure Kaonsample and a pure Pionsample

Create a Kaonsample and a Pionsample

▶ Kaonsample

→ Use free Kaon decay $\mathbf{K}^- \rightarrow \pi^- \pi^- \pi^+$

- ▶ 3 outgoing particles with correct charged
- ▶ Primary vertex outside of the target
- ▶ Cut on transverse momentum and Kaon mass

▶ Pionsample

→ Use diffractive production $\pi^- \mathbf{p} \rightarrow \pi^- \pi^- \pi^+ \mathbf{p}$

- ▶ 3 outgoing particles with correct charge
- ▶ Primary vertex inside the target
- ▶ Small angle to beam direction, similar momenta → same mass

In addition: Take a beamsample without any filtering for testing the method

Step 3: Determine probabilities to have hits in PMTs for π and K

Example: Particle with divergence θ_x, θ_y produces signal in PMT i

Question: Is it a kaon?

→ Use Bayes' Theorem:

$$P_{\theta_x, \theta_y}^i(\text{Kaon}|\text{Signal}) = \frac{\dots}{\dots}$$

Here:



Step 3: Determine probabilities to have hits in PMTs for π and K

Example: Particle with divergence θ_x, θ_y produces signal in PMT i

Question: Is it a kaon?

→ Use Bayes' Theorem:

$$P_{\theta_x, \theta_y}^i(\text{Kaon}|\text{Signal}) = \frac{P_{\theta_x, \theta_y}^i(\text{Signal}|\text{Kaon}) \cdot \dots}{\dots}$$

Here:

$P_{\theta_x, \theta_y}^i(\text{Signal}|\text{Kaon})$: Probability that Kaon at θ_x, θ_y (→ Kaonsample) produces signal in PMT i

Step 3: Determine probabilities to have hits in PMTs for π and K

Example: Particle with divergence θ_x, θ_y produces signal in PMT i

Question: Is it a kaon?

→ Use Bayes' Theorem:

$$P_{\theta_x, \theta_y}^i(\text{Kaon}|\text{Signal}) = \frac{P_{\theta_x, \theta_y}^i(\text{Signal}|\text{Kaon}) \cdot P_{\theta_x, \theta_y}(\text{Kaon})}{\dots}$$

Here:

$P_{\theta_x, \theta_y}^i(\text{Signal}|\text{Kaon})$: Probability that Kaon at θ_x, θ_y (→ Kaonsample) produces signal in PMT i

$P_{\theta_x, \theta_y}(\text{Kaon})$: Probability that Kaon has divergence θ_x and θ_y (→ Kaonsample)

Step 3: Determine probabilities to have hits in PMTs for π and K

Example: Particle with divergence θ_x, θ_y produces signal in PMT i

Question: Is it a kaon?

→ Use Bayes' Theorem:

$$P_{\theta_x, \theta_y}^i(\text{Kaon}|\text{Signal}) = \frac{P_{\theta_x, \theta_y}^i(\text{Signal}|\text{Kaon}) \cdot P_{\theta_x, \theta_y}(\text{Kaon})}{P_{\theta_x, \theta_y}^i(\text{Signal})}$$

Here:

- $P_{\theta_x, \theta_y}^i(\text{Signal}|\text{Kaon})$: Probability that Kaon at θ_x, θ_y (→ Kaonsample) produces signal in PMT i
- $P_{\theta_x, \theta_y}(\text{Kaon})$: Probability that Kaon has divergence θ_x and θ_y (→ Kaonsample)
- $P_{\theta_x, \theta_y}^i(\text{Signal})$: Probability that signal in PMT i is produced at θ_x, θ_y (→ Beamsample)

Step 3 continued

Pions and Kaons have the same divergence distribution:

$$P_{\theta_x, \theta_y}(\text{Kaon}) = P_{\theta_x, \theta_y}(\text{Pion}) = P_{\theta_x, \theta_y}(\text{Beam})$$

$\Rightarrow P_{\theta_x, \theta_y}^i(\text{Kaon}|\text{Signal})$ and $P_{\theta_x, \theta_y}^i(\text{Pion}|\text{Signal})$ have same normalization

factor $\frac{P_{\theta_x, \theta_y}(\text{Beam})}{P_{\theta_x, \theta_y}(\text{Signal})}$, thus


$$P_{\theta_x, \theta_y}^i(\text{Kaon}|\text{Signal}) \propto P_{\theta_x, \theta_y}^i(\text{Signal}|\text{Kaon})$$

$$P_{\theta_x, \theta_y}^i(\text{Pion}|\text{Signal}) \propto P_{\theta_x, \theta_y}^i(\text{Signal}|\text{Pion})$$

Also calculate


$$P_{\theta_x, \theta_y}^i(\text{Kaon}|\overline{\text{Signal}}) \text{ and } P_{\theta_x, \theta_y}^i(\text{Pion}|\overline{\text{Signal}})$$

Probability Distributions




Probability Distributions

Pions




Probability Distributions




Probability Distributions

Kaons



Probability Distributions




Additional Cut

- ▶ Statistics bad for large divergences
- ▶ Cut out events with

$$r = \sqrt{\theta_x^2 + \theta_y^2} < 200 \times 10^{-6}$$

- ▶ About 20% of all events are lost this way



Step 4: Calculate likelihoods from probabilities


- ▶ To obtain the log likelihood just add logarithms of probabilities

$$\log L(\text{Kaon}) = \sum_{i \in \text{PMT with Signal}} \log P_{\theta_x, \theta_y}^i(\text{Kaon} | \text{Signal}) + \sum_{j \in \text{PMT without Signal}} \log P_{\theta_x, \theta_y}^j(\text{Kaon} | \overline{\text{Signal}})$$


- ▶ Calculate the same for Pions.
- ▶ Compare them



Kaonsample



Pionsample



Step 5: Use likelihoods to identify particles

- ▶ Compare log likelihoods to get an ID for each CEDAR:
 - ▶ $\log L^K > \log L^\pi + \mathbf{A} \Rightarrow \text{PID } K$
 - ▶ $\log L^\pi > \log L^K + \mathbf{B} \Rightarrow \text{PID } \pi$
 - ▶ else no PID given
- ▶ Tune \mathbf{A} and \mathbf{B} due to efficiency/purity.




Step 5: Use likelihoods to identify particles

- ▶ Compare log likelihoods to get an ID for each CEDAR:
 - ▶ $\log L^K > \log L^\pi + \mathbf{A} \Rightarrow \text{PID } \mathbf{K}$
 - ▶ $\log L^\pi > \log L^K + \mathbf{B} \Rightarrow \text{PID } \pi$
 - ▶ else no PID given
- ▶ Tune \mathbf{A} and \mathbf{B} due to efficiency/purity.
- ▶ Combine CEDARs afterwards


$C_2 \setminus C_1$?	π	\mathbf{K}
?	?	π	\mathbf{K}
π	π	π	?
\mathbf{K}	\mathbf{K}	?	\mathbf{K}

Calculation of Purity

Look at




Then one knows (due to conservation of strangeness):



→ Identify \mathbf{h}'^- using the RICH and count the “wrong” particles




Select $h^- = \pi^-$ 


For CEDAR pions RICH should only see kaons ($\log_2(p(K)/p(\pi)) > 0$)
 Look at events with $p < 50$ GeV, cut out ± 0



Purity of Pion Selection




- ▶ Independent of likelihood cut
- ▶ $\approx 40\%$ better than multiplicity

Select $h^- = K^-$ 

For CEDAR pions RICH should only see pions ($\log_2 (p(K)/p(\pi)) < 0$)
 Look at events with $p < 50 \text{ GeV}$, cut out ± 0



Purity of Kaon Selection




- ▶ Independent of likelihood cut
- ▶ Compatible with multiplicity method
- ▶ Compatible with pion purity


Calculation of Efficiency

Look at expected number of particles in the beam
2.4% kaons, 97% pions

Kaons




Pions




⇒ Efficiency of around 80% for pions and kaons
Efficiency for kaons \approx 60% better than for multiplicity method

Direct Comparison with multiplicity Method




Direct Comparison with multiplicity Method



- Most of multiplicity kaons reproduced as kaons (896 of 1074, 83%)

Direct Comparison with multiplicity Method



- ▶ Most of multiplicity kaons reproduced as kaons (896 of 1074, 83%)
- ▶ 920 additional kaons from multiplicity pions

Summary

- ▶ COMPASS hadron beam consists of 97% pions and 2.4% kaons
- ▶ Pions and kaons have to be identified for analyses
- ▶ multiplicity method only identifies \approx **50%** of the kaons
 - ▶ Problems with divergent beams
- ▶ Likelihood method improves identification for divergent beams
 - ▶ Identifies 80% of the kaons with 87% purity.



Summary

- ▶ COMPASS hadron beam consists of 97% pions and 2.4% kaons
- ▶ Pions and kaons have to be identified for analyses
- ▶ multiplicity method only identifies \approx **50%** of the kaons
 - ▶ Problems with divergent beams
- ▶ Likelihood method improves identification for divergent beams
 - ▶ Identifies 80% of the kaons with 87% purity.

Outlook

- ▶ Adapt methods to 2009/2012 data taking
- ▶ Do finetuning using existing hadron analyses
- ▶ Redo $K\pi\pi$ analysis

