


Recent Results on Spectroscopy from COMPASS

Boris Grube

Physik-Department E18
Technische Universität München,
Garching, Germany


Hadron 2015

16. September 2015, Newport News, VA




Fixed-target experiment

- Two-stage spectrometer
- Large acceptance over wide kinematic range
- Electromagnetic and hadronic calorimeters
- Beam and final-state particle ID (CEDARs, RICH)



Fixed-target experiment

- Two-stage spectrometer
- Large acceptance over wide kinematic range
- Electromagnetic and hadronic calorimeters
- Beam and final-state particle ID (CEDARs, RICH)




Hadron spectroscopy

2008-09, 2012

- 190 GeV/c secondary **hadron beams**
 - h^- beam: 97 % π^- , 2 % K^- , 1 % \bar{p}
 - h^+ beam: 75 % p , 24 % π^+ , 1 % K^+
- **Various targets:** ℓ H₂, Ni, Pb, W

Fixed-target experiment

- Two-stage spectrometer
- Large acceptance over wide kinematic range
- Electromagnetic and hadronic calorimeters
- Beam and final-state particle ID (CEDARs, RICH)




Hadron spectroscopy

2008-09, 2012

- 190 GeV/c secondary **hadron beams**
 - h^- beam: 97 % π^- , 2 % K^- , 1 % \bar{p}
 - h^+ beam: 75 % p , 24 % π^+ , 1 % K^+
- **Various targets:** ℓH_2 , Ni, Pb, W

Spectroscopy program

- Explore light-meson spectrum for $m \gtrsim 2 \text{ GeV}/c^2$
- Search for states beyond the constituent quark model
- Precision measurement of known resonances



Hadron spectroscopy


2008-09, 2012

- 190 GeV/c secondary hadron beams
 - h^- beam: 97 % π^- , 2 % K^- , 1 % \bar{p}
 - h^+ beam: 75 % p , 24 % π^+ , 1 % K^+
- Various targets: ℓH_2 , Ni, Pb, W

- 1 Introduction
 - Meson production in diffractive dissociation
 - Partial-wave analysis method
- 2 PWA of diffractively produced $\pi^- \eta$ and $\pi^- \eta'$ final states
- 3 PWA of diffractively produced 3π final states
 - Observation of a new narrow axial-vector meson $a_1(1420)$
 - $J^{PC} = 1^{-+}$ spin-exotic partial wave
- 4 Conclusions and outlook


- 1 Introduction
 - Meson production in diffractive dissociation
 - Partial-wave analysis method
- 2 PWA of diffractively produced $\pi^- \eta$ and $\pi^- \eta'$ final states
- 3 PWA of diffractively produced 3π final states
 - Observation of a new narrow axial-vector meson $a_1(1420)$
 - $J^{PC} = 1^{-+}$ spin-exotic partial wave
- 4 Conclusions and outlook

Meson Production in Diffractive Dissociation



- Soft scattering of beam particle off target
 - Production of n forward-going hadrons
 - Target particle stays intact
- At $190 \text{ GeV}/c$, interaction dominated by space-like pomeron exchange
- All final-state particles are measured


Meson Production in Diffractive Dissociation




- **Exclusive measurement**

- Clean data sample
- Reduced four-momentum transfer squared $t' \equiv |t| - |t|_{\text{min}}$
 - Analyzed range:
 $0.1 < t' < 1.0 \text{ (GeV}/c)^2$

Example: $\pi^- \pi^+ \pi^-$ final state




Meson Production in Diffractive Dissociation




Example: $\pi^- \pi^+ \pi^-$ final state

- Exclusive measurement
 - Clean data sample
- Reduced four-momentum transfer squared $t' \equiv |t| - |t|_{\text{min}}$
 - Analyzed range:
 $0.1 < t' < 1.0 \text{ (GeV}/c)^2$




Meson Production in Diffractive Dissociation



- Beam particle gets excited into intermediate resonances X
- X dissociate into n -body final state
- Rich spectrum of intermediate states X

Disentanglement of all contributing X by **partial-wave analysis (PWA)**


Meson Production in Diffractive Dissociation



- Beam particle gets excited into intermediate resonances X
- X dissociate into n -body final state
- Rich spectrum of intermediate states X

Disentanglement of all contributing X by **partial-wave analysis (PWA)**


Meson Production in Diffractive Dissociation



- Beam particle gets excited into intermediate resonances X
- X dissociate into n -body final state
- Rich spectrum of intermediate states X

Disentanglement of all contributing X by **partial-wave analysis (PWA)**

Partial-Wave Analysis Method




Ansatz: Factorization of production and decay

$$\sigma(\tau; m_X) \propto \sum_{\epsilon=\pm 1} \left| \sum_i^{\text{waves}} T_i^\epsilon(m_X) A_i^\epsilon(\tau; m_X) \right|^2$$

- Transition amplitudes $T_i^\epsilon(m_X)$ contain interesting physics
- Decay amplitudes $A_i^\epsilon(\tau; m_X)$
 - Describe kinematic τ distribution of partial waves
 - Calculable using isobar model (for $n > 2$) and helicity formalism (Wigner D -functions)
- $\epsilon = \pm 1$: naturalities of exchange particle
 - 190 GeV/ c beam momentum \implies pomeron ($\epsilon = +1$) dominates

Partial-Wave Analysis Method




Ansatz: Factorization of production and decay

$$\sigma(\tau; m_X) \propto \sum_{\epsilon=\pm 1} \left| \sum_i^{\text{waves}} T_i^\epsilon(m_X) A_i^\epsilon(\tau; m_X) \right|^2$$

- Transition amplitudes $T_i^\epsilon(m_X)$ contain interesting physics
- Decay amplitudes $A_i^\epsilon(\tau; m_X)$
 - Describe **kinematic τ distribution** of partial waves
 - **Calculable** using **isobar model** (for $n > 2$) and **helicity formalism** (Wigner D -functions)
- $\epsilon = \pm 1$: **naturalities of exchange particle**
 - 190 GeV/ c beam momentum \implies **pomeron ($\epsilon = +1$) dominates**

Partial-Wave Analysis Method




Ansatz: Factorization of production and decay

$$\sigma(\tau; m_X) \propto \sum_{\epsilon=\pm 1} \left| \sum_i^{\text{waves}} T_i^\epsilon(m_X) A_i^\epsilon(\tau; m_X) \right|^2$$

- Transition amplitudes $T_i^\epsilon(m_X)$ contain interesting physics
- Decay amplitudes $A_i^\epsilon(\tau; m_X)$
 - Describe **kinematic τ distribution** of partial waves
 - **Calculable** using **isobar model** (for $n > 2$) and **helicity formalism** (Wigner D -functions)
- $\epsilon = \pm 1$: **naturalities of exchange particle**
 - 190 GeV/ c beam momentum \implies **pomeron ($\epsilon = +1$) dominates**

Partial-Wave Analysis Method



Two-stage analysis

$$\sigma(\tau; m_X) \propto \sum_{\epsilon=\pm 1} \left| \sum_i^{\text{waves}} T_i^\epsilon(m_X) A_i^\epsilon(\tau; m_X) \right|^2$$


1 Determination of m_X dependence of spin-density matrix

$$\rho_{ij}^\epsilon(m_X) = T_i^\epsilon(m_X) T_j^{\epsilon*}(m_X)$$

- Independent **maximum likelihood fits** to τ distributions in narrow m_X bins
- Take into account **detection efficiency**
- **No assumptions about resonance** content of X

2 Extraction of resonances

- χ^2 fit of resonance model to spin-density (sub)matrix



Two-stage analysis

$$\sigma(\tau; m_X) \propto \sum_{\epsilon=\pm 1} \left| \sum_i^{\text{waves}} T_i^\epsilon(m_X) A_i^\epsilon(\tau; m_X) \right|^2$$

1 Determination of m_X dependence of spin-density matrix

$$\rho_{ij}^\epsilon(m_X) = T_i^\epsilon(m_X) T_j^{\epsilon*}(m_X)$$

- Independent **maximum likelihood fits** to τ distributions in narrow m_X bins
- Take into account **detection efficiency**
- **No assumptions about resonance** content of X

2 Extraction of resonances

- χ^2 fit of **resonance model** to spin-density (sub)matrix

- 1 Introduction
 - Meson production in diffractive dissociation
 - Partial-wave analysis method
- 2 PWA of diffractively produced $\pi^- \eta$ and $\pi^- \eta'$ final states
- 3 PWA of diffractively produced 3π final states
 - Observation of a new narrow axial-vector meson $a_1(1420)$
 - $J^{PC} = 1^{-+}$ spin-exotic partial wave
- 4 Conclusions and outlook

- Odd-spin waves: **spin-exotic quantum numbers**
 - **Disputed** $J^{PC} = 1^{-+}$ resonance signals
 - $\pi_1(1400)$ in $\pi\eta$ and $\pi_1(1600)$ in $\pi\eta'$
- Comparison of $\pi\eta$ and $\pi\eta'$: information about **flavor structure**


Reconstruction from exclusive $\pi^- \pi^+ \pi^- \gamma\gamma$ final state

- $\eta \rightarrow \pi^+ \pi^- \pi^0$ with $\pi^0 \rightarrow \gamma\gamma$
- $\eta' \rightarrow \pi^+ \pi^- \eta$ with $\eta \rightarrow \gamma\gamma$

- Odd-spin waves: **spin-exotic quantum numbers**
 - **Disputed** $J^{PC} = 1^{-+}$ resonance signals
 - $\pi_1(1400)$ in $\pi\eta$ and $\pi_1(1600)$ in $\pi\eta'$
- Comparison of $\pi\eta$ and $\pi\eta'$: information about **flavor structure**

Reconstruction from exclusive $\pi^- \pi^+ \pi^- \gamma\gamma$ final state



- $\eta \rightarrow \pi^+ \pi^- \pi^0$ with $\pi^0 \rightarrow \gamma\gamma$
- $\eta' \rightarrow \pi^+ \pi^- \eta$ with $\eta \rightarrow \gamma\gamma$



- Odd-spin waves: **spin-exotic quantum numbers**
 - Disputed $J^{PC} = 1^{-+}$ resonance signals
 - $\pi_1(1400)$ in $\pi\eta$ and $\pi_1(1600)$ in $\pi\eta'$
- Comparison of $\pi\eta$ and $\pi\eta'$: information about **flavor structure**

Reconstruction from exclusive $\pi^- \pi^+ \pi^- \gamma\gamma$ final state

- $\eta \rightarrow \pi^+ \pi^- \pi^0$ with $\pi^0 \rightarrow \gamma\gamma$
- $\eta' \rightarrow \pi^+ \pi^- \eta$ with $\eta \rightarrow \gamma\gamma$



Quark-line picture for $n = (u, d)$ and pointlike resonances

- $\pi^- \eta$ and $\pi^- \eta'$ partial-wave intensities for **spin J** related by
 - Different **phase space** and barrier factors
 - **Branching fraction ratio b** of η and η' into $\pi^- \pi^+ \gamma \gamma$

$$N_J^{\pi\eta'}(m) \propto b \left[\frac{q^{\pi\eta'}(m)}{q^{\pi\eta}(m)} \right]^{2J+1} N_J^{\pi\eta}(m)$$



- $q =$ breakup momentum

Quark-line picture for $n = (u, d)$ and pointlike resonances

- $\pi^- \eta$ and $\pi^- \eta'$ partial-wave intensities for **spin J** related by
 - Different **phase space** and barrier factors
 - **Branching fraction ratio b** of η and η' into $\pi^- \pi^+ \gamma \gamma$

$$N_J^{\pi\eta'}(m) \propto b \left[\frac{q^{\pi\eta'}(m)}{q^{\pi\eta}(m)} \right]^{2J+1} N_J^{\pi\eta}(m)$$

- $q =$ breakup momentum








Quark-line picture for $n = (u, d)$ and pointlike resonances

- $\pi^- \eta$ and $\pi^- \eta'$ partial-wave intensities for **spin J** related by
 - Different **phase space** and barrier factors
 - **Branching fraction ratio b** of η and η' into $\pi^- \pi^+ \gamma \gamma$

$$N_J^{\pi\eta'}(m) \propto b \left[\frac{q^{\pi\eta'}(m)}{q^{\pi\eta}(m)} \right]^{2J+1} N_J^{\pi\eta}(m)$$




- $q =$ breakup momentum





- Similar even-spin waves
- Intermediate states couple to same final-state flavour content
- Similar physical content also in nonresonant high-mass region

$\pi^- \eta'$ final state; $\pi^- \eta$ scaled



- Resonance-model fit (Breit-Wigner)


- $\frac{N(a_2 \rightarrow \pi\eta')}{N(a_2 \rightarrow \pi\eta)} = (5 \pm 2) \%$

- First-time measurement of


$$\frac{N(a_4 \rightarrow \pi\eta')}{N(a_4 \rightarrow \pi\eta)} = (23 \pm 7) \%$$

$\pi^- \eta'$ final state; $\pi^- \eta$ scaled


Spin-exotic $J^{PC} = 1^{-+}$



2^{++}




Phase: $1^{-+} - 2^{++}$




- 1^{-+} intensities very different
- Suppression in $\pi\eta$ channel predicted for intermediate $|q\bar{q}g\rangle$ state
- Different phase motion in 1.6 GeV/c² region

$\pi^- \eta'$ final state; $\pi^- \eta$ scaled


Spin-exotic $J^{PC} = 1^{-+}$



2^{++}



Phase: $1^{-+} - 2^{++}$





- 1^{-+} resonance interpretation requires better understanding of
 - 2^{++} wave
 - Nonresonant contributions


$\pi^- \eta'$ final state; $\pi^- \eta$ scaled

Spin-exotic $J^{PC} = 1^{-+}$


2^{++}





Phase: $1^{-+} - 2^{++}$






Multi-Regge exchange, e.g.






- 1 Introduction
 - Meson production in diffractive dissociation
 - Partial-wave analysis method
- 2 PWA of diffractively produced $\pi^- \eta$ and $\pi^- \eta'$ final states
- 3 PWA of diffractively produced 3π final states
 - Observation of a new narrow axial-vector meson $a_1(1420)$
 - $J^{PC} = 1^{-+}$ spin-exotic partial wave
- 4 Conclusions and outlook






Strong $\pi^+ \pi^-$ correlations in $X^- \rightarrow \pi^- \pi^+ \pi^-$ decay




Strong $\pi^+ \pi^-$ correlations in $X^- \rightarrow \pi^- \pi^+ \pi^-$ decay



Strong $\pi^+ \pi^-$ correlations in $X^- \rightarrow \pi^- \pi^+ \pi^-$ decay




Strong $\pi^+ \pi^-$ correlations in $X^- \rightarrow \pi^- \pi^+ \pi^-$ decay




Isobar model

- X^- decays via intermediate $\pi^+ \pi^-$ resonance = "isobar"
 - $[\pi\pi]_S \quad J^{PC} = 0^{++}$
 - $\rho(770) \quad 1^{--}$
 - $f_0(980) \quad 0^{++}$
 - $f_2(1270) \quad 2^{++}$
 - $f_0(1500) \quad 0^{++}$
 - $\rho_3(1690) \quad 3^{--}$
- PWA requires precise knowledge of isobar $\rightarrow \pi^+ \pi^-$ amplitude




Isobar model

- X^- decays via intermediate $\pi^+ \pi^-$ resonance = "isobar"
 - $[\pi\pi]_S$ $J^{PC} = 0^{++}$
 - $\rho(770)$ 1^{--}
 - $f_0(980)$ 0^{++}
 - $f_2(1270)$ 2^{++}
 - $f_0(1500)$ 0^{++}
 - $\rho_3(1690)$ 3^{--}
- PWA requires precise knowledge of isobar $\rightarrow \pi^+ \pi^-$ amplitude



Isobar model

- X^- decays via **intermediate $\pi^+ \pi^-$ resonance = "isobar"**
 - $[\pi\pi]_S$ $J^{PC} = 0^{++}$
 - $\rho(770)$ 1^{--}
 - $f_0(980)$ 0^{++}
 - $f_2(1270)$ 2^{++}
 - $f_0(1500)$ 0^{++}
 - $\rho_3(1690)$ 3^{--}
- PWA requires precise knowledge of **isobar $\rightarrow \pi^+ \pi^-$ amplitude**



PWA of $\pi^- p \rightarrow (3\pi)^- p_{\text{recoil}}$

Two Data Sets

- 1 $\pi^- \pi^+ \pi^-$ (50 M events)
- 2 Crosscheck with $\pi^- \pi^0 \pi^0$ (3.5 M events)
 - Very different acceptance
 - Isobars separated by isospin
 - $I = 1$ isobars: $\pi^- \pi^0$
 - $I = 0$ isobars: $\pi^0 \pi^0$

Complex correlation of $m_{3\pi}$ and t'

- Two-dimensional PWA in bins of t' and $m_{3\pi}$
 - $\pi^- \pi^+ \pi^-$: 11 t' bins
 - $\pi^- \pi^0 \pi^0$: 8 t' bins
- Better disentanglement of resonant and nonresonant contributions

PWA of $\pi^- p \rightarrow (3\pi)^- p_{\text{recoil}}$


Two Data Sets

- 1 $\pi^- \pi^+ \pi^-$ (50 M events)
- 2 Crosscheck with $\pi^- \pi^0 \pi^0$ (3.5 M events)
 - Very different acceptance
 - Isobars separated by isospin
 - $I = 1$ isobars: $\pi^- \pi^0$
 - $I = 0$ isobars: $\pi^0 \pi^0$


Complex correlation of $m_{3\pi}$ and t'

- Two-dimensional PWA in bins of t' and $m_{3\pi}$
 - $\pi^- \pi^+ \pi^-$: 11 t' bins
 - $\pi^- \pi^0 \pi^0$: 8 t' bins
- Better disentanglement of resonant and nonresonant contributions

$$800 < m_{3\pi} < 850 \text{ MeV}/c^2$$



$$1600 < m_{3\pi} < 1650 \text{ MeV}/c^2$$



PWA of $\pi^- p \rightarrow (3\pi)^- p_{\text{recoil}}$


Two Data Sets

- 1 $\pi^- \pi^+ \pi^-$ (50 M events)
- 2 Crosscheck with $\pi^- \pi^0 \pi^0$ (3.5 M events)
 - Very different acceptance
 - Isobars separated by isospin
 - $I = 1$ isobars: $\pi^- \pi^0$
 - $I = 0$ isobars: $\pi^0 \pi^0$


Complex correlation of $m_{3\pi}$ and t'

- Two-dimensional PWA in bins of t' and $m_{3\pi}$
 - $\pi^- \pi^+ \pi^-$: 11 t' bins
 - $\pi^- \pi^0 \pi^0$: 8 t' bins
- Better disentanglement of resonant and nonresonant contributions


$0.10 < t' < 0.12 \text{ (GeV/c)}^2$





$0.44 < t' < 1.00 \text{ (GeV/c)}^2$






- $\pi^- \pi^+ \pi^-$ invariant mass spectrum
- $1^{++} 0^+ \rho(770) \pi S$:
 $a_1(1260)$
- $2^{++} 1^+ \rho(770) \pi D$:
 $a_2(1320)$
- $2^{-+} 0^+ f_2(1270) \pi S$:
 $\pi_2(1670)$







- $\pi^- \pi^+ \pi^-$ invariant mass spectrum
- $1^{++} 0^+ \rho(770) \pi S$:
 $a_1(1260)$
- $2^{++} 1^+ \rho(770) \pi D$:
 $a_2(1320)$
- $2^- 0^+ f_2(1270) \pi S$:
 $\pi_2(1670)$





- $\pi^- \pi^+ \pi^-$ invariant mass spectrum
- $1^{++} 0^+ \rho(770) \pi S$:
 $a_1(1260)$
- $2^{++} 1^+ \rho(770) \pi D$:
 $a_2(1320)$
- $2^{-+} 0^+ f_2(1270) \pi S$:
 $\pi_2(1670)$



- $\pi^- \pi^+ \pi^-$ invariant mass spectrum
- $1^{++} 0^+ \rho(770) \pi S$:
 $a_1(1260)$
- $2^{++} 1^+ \rho(770) \pi D$:
 $a_2(1320)$
- $2^{-+} 0^+ f_2(1270) \pi S$:
 $\pi_2(1670)$





- $\pi^- \pi^+ \pi^-$ invariant mass spectrum
- $1^{++} 0^+ \rho(770) \pi S$:
 $a_1(1260)$
- $2^{++} 1^+ \rho(770) \pi D$:
 $a_2(1320)$
- $2^{-+} 0^+ f_2(1270) \pi S$:
 $\pi_2(1670)$



In total 88 partial waves

- Largest wave set used so far for $\pi^- \pi^+ \pi^-$
- Spin J up to 6
- Orbital angular momentum L up to 6



PWA of $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p_{\text{recoil}}$: Selected Small Waves

[arXiv:1509.00992]

$4^{++} 1^+ \rho(770) \pi G$


- $a_4(2040)$

$0^{-+} 0^+ f_0(980) \pi S$

- $\pi(1800)$

$1^{++} 0^+ f_0(980) \pi P$

- **Unexpected peak around $1.4 \text{ GeV}/c^2$**
- Small intensity: $\approx 0.3\%$
- Similar signal in $\pi^- \pi^0 \pi^0$



PWA of $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p_{\text{recoil}}$: Selected Small Waves

[arXiv:1509.00992]

$4^{++} 1^+ \rho(770) \pi G$



- $a_4(2040)$

$0^{-+} 0^+ f_0(980) \pi S$

- $\pi(1800)$

$1^{++} 0^+ f_0(980) \pi P$

- **Unexpected peak around $1.4 \text{ GeV}/c^2$**
- Small intensity: $\approx 0.3\%$
- Similar signal in $\pi^- \pi^0 \pi^0$



PWA of $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p_{\text{recoil}}$: Selected Small Waves

[arXiv:1509.00992]

$4^{++} 1^+ \rho(770) \pi G$


- $a_4(2040)$

$0^{-+} 0^+ f_0(980) \pi S$

- $\pi(1800)$

$1^{++} 0^+ f_0(980) \pi P$

- **Unexpected peak around $1.4 \text{ GeV}/c^2$**
- Small intensity: $\approx 0.3\%$
- Similar signal in $\pi^- \pi^0 \pi^0$



PWA of $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p_{\text{recoil}}$: Selected Small Waves

$4^{++} 1^+ \rho(770) \pi G$

- $a_4(2040)$

$0^{-+} 0^+ f_0(980) \pi S$

- $\pi(1800)$

$1^{++} 0^+ f_0(980) \pi P$


- **Unexpected peak around $1.4 \text{ GeV}/c^2$**

- Small intensity: $\approx 0.3\%$

- Similar signal in $\pi^- \pi^0 \pi^0$

$\pi^- \pi^0 \pi^0$

$\pi^- \pi^+ \pi^-$ scaled for each plot





Is Peak in $1^{++} 0^+ f_0(980) \pi P$ Wave a Model Artifact?

Novel analysis method

(inspired by E791 analysis, PRD **73** (2006) 032204)

- Replace $J^{PC} = 0^{++}$ isobar parametrizations by **piece-wise constant amplitudes** in $m_{\pi^+\pi^-}$ bins
- Extract $m_{3\pi}$ dependence of $J^{PC} = 0^{++}$ isobar amplitude from data
 - Drastic reduction of model bias
 - *Caveat*: significant increase in number of fit parameters






Is Peak in $1^{++} 0^+ f_0(980) \pi P$ Wave a Model Artifact?

Novel analysis method


(inspired by E791 analysis, PRD **73** (2006) 032204)

- Replace $J^{PC} = 0^{++}$ isobar parametrizations by **piece-wise constant amplitudes** in $m_{\pi^+\pi^-}$ bins
- Extract $m_{3\pi}$ dependence of $J^{PC} = 0^{++}$ isobar amplitude from data
 - Drastic **reduction of model bias**
 - *Caveat*: significant **increase in number of fit parameters**






- Correlation of 3π intensity around $1.4 \text{ GeV}/c^2$ with $f_0(980)$
- $f_0(980)$ semicircle in Argand diagram
- Confirms that $f_0(980)\pi$ signal is *not* an artifact of isobar parametrization




- **Correlation** of 3π intensity around $1.4 \text{ GeV}/c^2$ with $f_0(980)$
- $f_0(980)$ semicircle in Argand diagram
- Confirms that $f_0(980)\pi$ signal is *not* an artifact of isobar parametrization



$\pi\pi$ S-Wave Amplitude in $J^{PC} = 1^{++} 3\pi$ Wave [arXiv:1509.00992]




- Correlation of 3π intensity around $1.4 \text{ GeV}/c^2$ with $f_0(980)$
- $f_0(980)$ semicircle in Argand diagram
- Confirms that $f_0(980)\pi$ signal is *not* an artifact of isobar parametrization



- Correlation of 3π intensity around $1.4 \text{ GeV}/c^2$ with $f_0(980)$
- $f_0(980)$ semicircle in Argand diagram
- Confirms that $f_0(980)\pi$ signal is *not* an artifact of isobar parametrization




- Correlation of 3π intensity around 1.4 GeV/c^2 with $f_0(980)$
- $f_0(980)$ semicircle in Argand diagram
- Confirms that $f_0(980)\pi$ signal is *not* an artifact of isobar parametrization




- Coupling of $\pi(1800)$ to $f_0(980)\pi$ and $f_0(1500)\pi$ decay modes


See talk 1E2 by F. Krinner (Thu, 8:55)





- Coherent sum of resonant (Breit-Wigner) and nonresonant terms



- Coherent sum of resonant (Breit-Wigner) and nonresonant terms



- 1^{++} peak consistent with Breit-Wigner resonance
- $a_1(1420)$:
 $M_0 = 1414^{+15}_{-13} \text{ MeV}/c^2$
 $\Gamma_0 = 153^{+8}_{-23} \text{ MeV}/c^2$






- 1^{++} peak consistent with Breit-Wigner resonance

- $a_1(1420)$:

$$M_0 = 1414^{+15}_{-13} \text{ MeV}/c^2$$

$$\Gamma_0 = 153^{+8}_{-23} \text{ MeV}/c^2$$






- 1^{++} peak consistent with Breit-Wigner resonance

- $a_1(1420)$:

$$M_0 = 1414_{-13}^{+15} \text{ MeV}/c^2$$

$$\Gamma_0 = 153_{-23}^{+8} \text{ MeV}/c^2$$



Nature unclear

- No quark-model states expected at $1.4 \text{ GeV}/c^2$
- Ground state $a_1(1260)$ very close and wider
- Seen only in $f_0(980)\pi$ decay mode
- Isospin partner of narrow $f_1(1420)$?
- Suspiciously close to $K\bar{K}^*$ threshold

Several proposed explanations

- **Two-quark-tetraquark** mixed state [Wang, arXiv:1401.1134]
- **Tetraquark** with mixed flavor symmetry [Chen *et al.*, PRD **91** (2015) 094022]
- Two-channel **unitarized Deck** amplitude + direct $a_1(1260)$ production [Basdevant and Berger, PRL **114** (2015) 192001 and arXiv:1501.04643]
See talk 1B4 by E. Berger (Mon, 5:30)
- Singularity (branching point) in **triangle diagram** [Mikhasenko *et al.*, PRD **91** (2015) 094015]
See talk 5A1 by B. Ketzner (Thu, 8:30)

Several proposed explanations

- Two-quark-tetraquark mixed state [Wang, arXiv:1401.1134]

- Tetraquark with mixed flavor symmetry [Chen *et al.*, PRD **91** (2015) 094022]

- Two-channel unitarized Deck amplitude + direct $a_1(1260)$ production

[Basdevant and Berger, PRL **114** (2015) 192001 and arXiv:1501.04643]

See talk 1B4 by E. Berger (Mon, 5:30)

- Singularity (branching point) in triangle diagram

[Mikhasenko *et al.*, PRD **91** (2015) 094015]

See talk 5A1 by B. Ketzner (Thu, 8:30)

Several proposed explanations

- Two-quark-tetraquark mixed state [Wang, arXiv:1401.1134]

- Tetraquark with mixed flavor symmetry

[Chen *et al.*, PRD **91** (2015) 094022]

- Two-channel unitarized Deck amplitude + direct $a_1(1260)$ production


[Basdevant and Berger, PRL **114** (2015) 192001 and arXiv:1501.04643]

See talk 1B4 by E. Berger (Mon, 5:30)

- Singularity (branching point) in triangle diagram


[Mikhasenko *et al.*, PRD **91** (2015) 094015]

See talk 5A1 by B. Ketzer (Thu, 8:30)



Spin-Exotic $J^{PC} = 1^{-+}$ Signal in $(3\pi)^-$ PWA

- Broad intensity bump
- Similar in both channels




$\pi^- \pi^0 \pi^0$

$\pi^- \pi^+ \pi^-$ scaled


Spin-Exotic $J^{PC} = 1^{-+}$ Signal in $\pi^- \pi^+ \pi^-$ PWA

Drastic Change of Mass Spectrum with t'

“Low” $t' \approx 0.1 \text{ (GeV}/c)^2$



“High” $t' \approx 0.8 \text{ (GeV}/c)^2$




- Dominant nonresonant contribution
 - Needs to be better understood in order to extract resonance content

Spin-Exotic $J^{PC} = 1^{-+}$ Signal in $\pi^- \pi^+ \pi^-$ PWA

Model for Nonresonant Component

Deck effect





- MC pseudodata generated according to model of Deck amplitude

based on ACCMOR, NPB **182** (1981) 269

- Analyzed like real data

Spin-Exotic $J^{PC} = 1^{-+}$ Signal in $\pi^{-}\pi^{+}\pi^{-}$ PWA

Deck-Model for Nonresonant Component




- Deck MC scaled to t' -summed intensity
 - Similar mass spectrum at low t'
 - Different shape at high t'

Spin-Exotic $J^{PC} = 1^{-+}$ Signal in $\pi^{-}\pi^{+}\pi^{-}$ PWA

Relative Phase w.r.t. $1^{++}0^{+}\rho(770)\pi S$ Wave

$\pi^{\pm}p \rightarrow \pi^{\mp}\pi^{\pm}\pi^{\pm}p$ (COMPASS 2008)



- Slow phase 60° motion in $1.6 \text{ GeV}/c^2$ region independent of t'

- 1 Introduction
 - Meson production in diffractive dissociation
 - Partial-wave analysis method
- 2 PWA of diffractively produced $\pi^- \eta$ and $\pi^- \eta'$ final states
- 3 PWA of diffractively produced 3π final states
 - Observation of a new narrow axial-vector meson $a_1(1420)$
 - $J^{PC} = 1^{-+}$ spin-exotic partial wave
- 4 Conclusions and outlook

Precise data on pion diffraction

- PWA reliably extracts even very small signals
 - New axial-vector state $a_1(1420)$ in $(3\pi)^-$ final states
- Novel analysis schemes:
 - PWA in bins of t'
 - Better separation of resonant and nonresonant contribution
 - Extraction of $\pi\pi$ S -wave amplitude from $\pi^-\pi^+\pi^-$ system
 - Study dependence on 3π source
 - Study rescattering effects
 - Extension to higher $\pi\pi$ waves

Nonresonant contributions play important role

- Limit extraction of **resonance parameters**
- First studies using **Deck models**
- **Extraction** of nonresonant contributions **from data**
 - Collaboration with JPAC: Veneziano amplitudes + finite-energy sum rules

Other ongoing analyses

- Pion diffraction into $\pi^- \eta \eta$, $\pi^- \pi^0 \omega$, $K \bar{K} \pi$, $K \bar{K} \pi \pi$, ...
- Kaon diffraction into $K^- \pi^+ \pi^-$
- Central-production reactions See talk 1E1 by A. Austregesilo (Thu, 8:30)
- $\pi \gamma$ scattering using Primakoff reactions on heavy targets See talk 6C4 by M. Krämer (Thu, 11:45)

Nonresonant contributions play important role

- Limit extraction of **resonance parameters**
- First studies using **Deck models**
- **Extraction** of nonresonant contributions **from data**
 - Collaboration with JPAC: Veneziano amplitudes + finite-energy sum rules

Other ongoing analyses

- **Pion diffraction** into $\pi^- \eta \eta$, $\pi^- \pi^0 \omega$, $K \bar{K} \pi$, $K \bar{K} \pi \pi$, ...
- **Kaon diffraction** into $K^- \pi^+ \pi^-$
- **Central-production** reactions See talk 1E1 by A. Austregesilo (Thu, 8:30)
- $\pi \gamma$ scattering using **Primakoff reactions** on heavy targets See talk 6C4 by M. Krämer (Thu, 11:45)