

# Recent results from COMPASS on exclusive muoproduction



Andrzej Sandacz

National Centre for Nuclear Research, Warsaw

*on behalf of the COMPASS Collaboration*



## *DSPIN-15*

XVI Workshop on High Energy Spin Physics

Dubna, Russia, September 8-12, 2015



## COMPASS GPD program activities

### ➤ Exclusive vector meson muoproduction from 2002-2010 data

with transversely polarised proton/deuteron targets ( ${}^6\text{LiD}$ ,  $\text{NH}_3$ )

no recoil detector → disadvantage for exclusive measurements

opportunity to get early results which are sensitive to GPDs E and chiral odd GPDs

  
discussed in this talk

### ➤ 'DVCS test' runs in 2008 (1.5 day) and 2009 (10 days)

40 cm  $\text{LH}_2$  target and small RPD (used for hadron spectroscopy program)

analyses of the 'DVCS test' data demonstrated feasibility to measure

exclusive  $\gamma$  (DVCS and BH) and exclusive  $\pi^0$  production at COMPASS

### ➤ GPD program of COMPASS-II (since 2012)

a part of approved new COMPASS proposal


DVCS and HEMP with polarised  $\mu^+$  and  $\mu^-$  beams at 160 GeV and

unpolarised and transversely polarised proton targets ( $\text{LH}_2$ ,  $\text{NH}_3$ )

with large recoil proton detector and large angular coverage by EM calorimetry

  
discussed in this talk

# GPDs and Hard Exclusive Meson Production



- factorisation proven only for  $\sigma_L$   
 $\sigma_T$  suppressed by  $1/Q^2$
- wave function of meson (DA)  
additional non-perturbative term
- at  $Q^2 \approx \text{few GeV}^2$  higher order pQCD  
terms important

## Chiral-even GPDs

helicity of parton unchanged

$$H^{q,g}(x, \xi, t)$$

$$\tilde{H}^{q,g}(x, \xi, t)$$

$$E^{q,g}(x, \xi, t)$$

$$\tilde{E}^{q,g}(x, \xi, t)$$

## Chiral-odd GPDs

helicity of parton changed (not probed by DVCS)

$$H_T^q(x, \xi, t)$$

$$\tilde{H}_T^q(x, \xi, t)$$

$$E_T^q(x, \xi, t)$$

$$\tilde{E}_T^q(x, \xi, t)$$

## Flavour separation for GPDs

example:

$$E_{\rho^0} = \frac{1}{\sqrt{2}} \left( \frac{2}{3} E^u + \frac{1}{3} E^d + \frac{3}{8} E^g \right)$$

$$E_{\omega} = \frac{1}{\sqrt{2}} \left( \frac{2}{3} E^u - \frac{1}{3} E^d + \frac{1}{8} E^g \right)$$

$$E_{\varphi} = -\frac{1}{3} E^s - \frac{1}{8} E^g$$

- contribution from gluons at the same order of  $\alpha_s$  as from quarks

## Spin-dependent cross section for exclusive meson leptonproduction

$$\left[ \frac{\alpha_{em}}{8\pi^3} \frac{y^2}{1-\epsilon} \frac{1-x_{Bj}}{x_{Bj}} \frac{1}{Q^2} \right]^{-1} \frac{d\sigma}{dx_{Bj} dQ^2 dt d\phi d\phi_s}$$

$$= \underbrace{\frac{1}{2}(\sigma_{++}^{++} + \sigma_{++}^{--}) + \epsilon\sigma_{00}^{++}}_{\text{}} - \epsilon \cos(2\phi) \text{Re} \sigma_{+-}^{++} - \sqrt{\epsilon(1+\epsilon)} \cos \phi \text{Re} (\sigma_{+0}^{++} + \sigma_{+0}^{--})$$

$$- P_\ell \sqrt{\epsilon(1-\epsilon)} \sin \phi \text{Im} (\sigma_{+0}^{++} + \sigma_{+0}^{--})$$

$$- S_L \left[ \epsilon \sin(2\phi) \text{Im} \sigma_{+-}^{++} + \sqrt{\epsilon(1+\epsilon)} \sin \phi \text{Im} (\sigma_{+0}^{++} - \sigma_{+0}^{--}) \right]$$


$$+ S_L P_\ell \left[ \sqrt{1-\epsilon^2} \frac{1}{2} (\sigma_{++}^{++} - \sigma_{++}^{--}) - \sqrt{\epsilon(1-\epsilon)} \cos \phi \text{Re} (\sigma_{+0}^{++} - \sigma_{+0}^{--}) \right]$$

$$- S_T \left[ \sin(\phi - \phi_S) \text{Im} (\sigma_{++}^{+-} + \epsilon\sigma_{00}^{+-}) + \frac{\epsilon}{2} \sin(\phi + \phi_S) \text{Im} \sigma_{+-}^{+-} + \frac{\epsilon}{2} \sin(3\phi - \phi_S) \text{Im} \sigma_{+-}^{-+} \right]$$

$$+ \sqrt{\epsilon(1+\epsilon)} \sin \phi_S \text{Im} \sigma_{+0}^{+-} + \sqrt{\epsilon(1+\epsilon)} \sin(2\phi - \phi_S) \text{Im} \sigma_{+0}^{-+}$$

$$+ S_T P_\ell \left[ \sqrt{1-\epsilon^2} \cos(\phi - \phi_S) \text{Re} \sigma_{++}^{+-} \right]$$

$$- \sqrt{\epsilon(1-\epsilon)} \cos \phi_S \text{Re} \sigma_{+0}^{+-} - \sqrt{\epsilon(1-\epsilon)} \cos(2\phi - \phi_S) \text{Re} \sigma_{+0}^{-+} \Big].$$



$\sigma_{mn}^{ij}$ : helicity-dependent photoabsorption cross sections and interference terms

$$\sigma_{mn}^{ij}(x_B, Q^2, t) \propto \sum (M_m^i)^* M_n^j$$

$M_m^i$ : amplitude for subprocess  $\gamma^* p \rightarrow V p'$  with photon helicity  $m$  and target proton helicity  $i$

$$\epsilon = \frac{1 - y - \frac{1}{4}y^2}{1 - y + \frac{1}{2}y^2 + \frac{1}{4}y^2}$$

$$y = 2x_{Bj} M_p / Q$$

# Azimuthal asymmetries of cross section for exclusive meson lepton production

5 single spin asymmetries

$$A_{UT}^{\sin(\varphi - \varphi_s)} = -\frac{\text{Im}(\sigma_{++}^{+-} + \epsilon \sigma_{00}^{+-})}{\sigma_0}$$

$$A_{UT}^{\sin(\varphi + \varphi_s)} = -\frac{\text{Im} \sigma_{+-}^{+-}}{\sigma_0}$$

$$A_{UT}^{\sin(2\varphi - \varphi_s)} = -\frac{\text{Im} \sigma_{+0}^{-+}}{\sigma_0}$$

$$A_{UT}^{\sin(3\varphi - \varphi_s)} = -\frac{\text{Im} \sigma_{+-}^{-+}}{\sigma_0}$$

$$A_{UT}^{\sin \varphi_s} = -\frac{\text{Im} \sigma_{+0}^{+-}}{\sigma_0}$$

3 double spin asymmetries

$$A_{LT}^{\cos(\varphi - \varphi_s)} = \frac{\text{Re} \sigma_{++}^{+-}}{\sigma_0}$$

$$A_{LT}^{\cos(2\varphi - \varphi_s)} = -\frac{\text{Re} \sigma_{+0}^{-+}}{\sigma_0}$$

$$A_{LT}^{\cos \varphi_s} = -\frac{\text{Re} \sigma_{+0}^{+-}}{\sigma_0}$$

$\sigma_0$  - 'unpolarised cross section'

$$\sigma_0 = \frac{1}{2}(\sigma_{++}^{++} + \sigma_{++}^{--}) + \epsilon \sigma_{00}^{++} = \sigma_L + \epsilon \sigma_T$$


# COMPASS setup

with polarised targets  
 NIM A 577(2007) 455

two-stage forward spectrometer **SM1 + SM2**

≈ 300 tracking detectors planes – high redundancy  
 variety of tracking detectors to cope with different  
 particle flux from  $\theta = 0$  to  $\theta \approx 200$  mrad

+ calorimetry,  $\mu$ ID, RICH




$\mu^+$  beam from SPS  
 beam polarisation  $\approx -80\%$

## Polarised Target

Two 30cm and one 60 cm long target cells [two 60cm long cells in 2002-2004] with opposite polarization

material:	$\text{NH}_3$ (protons)	$^6\text{LiD}$ (deuterons)
polarization:	$\approx 90\%$	$[\approx 50\%]$
dilution factor for exclusive $\rho^0$ production:	$\approx 25\%$	$[\approx 44\%]$

Luminosity  $5 \cdot 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$



# Exclusive $\rho^0$ production on $p^\uparrow$ and $d^\uparrow$ at COMPASS

$$\mu N \rightarrow \mu \rho^0 N$$

i.e. incoherent process

Transversely polarised **proton** target ( $\text{NH}_3$ ), 2007, 2010

Transversely polarised **deuteron** target ( ${}^6\text{LiD}$ ), 2003-2004

note: there was no RPD for these data

only two hadron tracks of opposite charge associated to the primary vertex

DIS cuts




$$\begin{aligned} 1 < Q^2 < 10 \text{ GeV}^2 \\ 0.1 < y < 0.9 \\ 0.003 < x < 0.35 \\ W > 5 \text{ GeV} \end{aligned}$$

cuts specific for exclusive  $\rho^0$  analysis

$$\begin{aligned} 0.5 < M_{\pi\pi} < 1.1 \text{ GeV} \\ -2.5 < E_{\text{miss}} < 2.5 \text{ GeV} \\ E_{\rho^0} > 15 \text{ GeV} \\ 0.05 < p_T^2 < 0.5 \text{ GeV}^2 \text{ [NH}_3\text{]} \\ 0.1 < p_T^2 < 0.5 \text{ GeV}^2 \text{ [}^6\text{LiD]} \end{aligned}$$


$$E_{\text{miss}} = (M_X^2 - M_p^2) / (2M_p)$$

— proton data (797 000 evts)  
— deuteron data (97 000 evts)






# Exclusive $\omega$ production on $p^\uparrow$ at COMPASS



Transversely polarised **proton** target ( $\text{NH}_3$ ), 2010 data

note: there was no RPD for these data







only two hadron tracks of opposite charge associated to the primary vertex  
 only two ECAL clusters time-correlated with beam and not associated to a charged particle

After selections and cuts  $\approx 19\,000$  evts

$$\begin{aligned}
 &|M_{\pi\pi\pi} - M_\omega^{\text{PDG}}| < 0.07 \text{ GeV} \\
 &-3 < E_{\text{miss}} < 3 \text{ GeV} \\
 &E_\omega > 14 \text{ GeV} \\
 &0.05 < p_T^2 < 0.5 \text{ GeV}^2
 \end{aligned}$$

$$\begin{aligned}
 &1 < Q^2 < 10 \text{ GeV}^2 \\
 &0.1 < y < 0.9 \\
 &0.003 < x < 0.35 \\
 &W > 5 \text{ GeV}
 \end{aligned}$$



dominant coherent contrib.


dominant non-exclusive bkg.



# Extraction of asymmetries and subtraction on non-exclusive background

- $\rho^0$  analysis
  - 1D (deuteron) and 2D (proton) binned maximum likelihood estimator with subtraction of background in  $(\varphi, \varphi_S)$  bins
- $\omega$  analysis
  - Unbinned maximum likelihood estimator with simultaneous fit of signal and background asymmetries

**Background rejection:**  
*For each target cell and polarization state*



shape of semi-inclusive background from MC  
*(LEPTO with COMPASS tuning + simulation of spectrometer response + reconstruction as for real data)*



MC weighted using ratio between real data and MC for wrong charge combination sample ( $h^+h^+\gamma\gamma + h^-h^-\gamma\gamma$ )

$$w(E_{miss}) = \frac{N_{RD}^{h^+h^+\gamma\gamma}(E_{miss}) + N_{RD}^{h^-h^-\gamma\gamma}(E_{miss})}{N_{MC}^{h^+h^+\gamma\gamma}(E_{miss}) + N_{MC}^{h^-h^-\gamma\gamma}(E_{miss})}$$

Normalization of MC to the real data using two component fit  
**Gaussian function (signal) + shape from MC (bkg)**

# Results on $A_{UT}^{\sin(\phi-\phi_S)}$ for exclusive $\rho^0$ production

→ NPB 865 (2012) 1



- $A_{UT}^{\sin(\phi-\phi_S)}$  for transversely polarised protons and deuterons small, compatible with 0
- for the **proton** agreement with HERMES results  
COMPASS results with statistical errors improved by **factor 3** and extended kinematic range
- for the **deuteron the first** measurement
- reasonable agreement with predictions of the GPD model of Goloskokov - Kroll  
[EPJ C59 (2009) 809]

small values expected due to approximate cancellation of contributions from  $E^u$  and  $E^d$ ,  $E^u \approx -E^d$

$$E_{\rho^0}^p \sim \frac{2}{3} E^u + \frac{1}{3} E^d + \frac{3}{8} E^g \quad \text{vs.} \quad E_{\omega}^p \sim \frac{2}{3} E^u - \frac{1}{3} E^d + \frac{3}{8} E^g \quad (\text{cf. upper-right plot})$$

# Complete set of transverse target spin asymmetries for exclusive $\rho^0$ production


- Improved method of extraction (2D)
- Simultaneous extraction of
  - 5 single spin asymmetries and
  - 3 double spin asymmetries
  - for transversely polarised **protons**

→ **PLB 731 (2014) 19**


— predictions of GPD model of Goloskokov-Kroll

● reasonable agreement with GK model (also for double spin asym.)

## Single spin asymmetries



# Azimuthal asymmetries for exclusive $\rho^0$ production on $p^\uparrow$



$$\langle x_{Bj} \rangle = 0.039, \langle Q^2 \rangle = 2.0 \text{ GeV}^2$$

$$\langle p_T^2 \rangle = 0.18 \text{ GeV}^2, \langle W \rangle = 8.1 \text{ GeV}^2$$

$M_{Vp', \gamma^*p}$  helicity amplitudes  
 $\sigma_0$  unpolarised cross section  
 $H_T(x, 0, 0) = h_1(x)$   
 $\bar{E}_T = 2\tilde{H}_T - E_T$

- asymmetries small, compatible with 0, except


$$A_{UT}^{\sin\varphi_s} = -0.019 \pm 0.008 \pm 0.003$$

- indication of  $H_T$ , 'transversity' GPD, contribution

- larger effects for some asymmetries expected for exclusive  $\omega$  production

# Azimuthal asymmetries for exclusive $\omega$ production on $p^\uparrow$

## Single spin asymmetries



$$\langle x_{Bj} \rangle = 0.049, \langle Q^2 \rangle = 2.2 \text{ GeV}^2$$


$$\langle p_T^2 \rangle = 0.17 \text{ GeV}^2, \langle W \rangle = 7.1 \text{ GeV}^2$$

- new result, to be published
- unbinned maximum likelihood method
- extraction of 8 transverse spin asymmetries

comparison to **modified** GPD model of GK

with **added  $\pi^0$  pole exchange**

EPJA 50 (2014) 9, 146



parameters constrained by HERMES SDMEs for  $\omega$   
except sign of  $\pi\omega$  form factor  
more sensitivity in azimuthal asymmetries

GK predictions for COMPASS, [private com.](#)

- no pion pole
- - - positive  $\pi\omega$  form factor
- ⋯ negative  $\pi\omega$  form factor

- 🌐 no clear conclusion from the comparison at the moment

## COMPASS-II time lines

Part of the COMPASS-II proposal approved and scheduled by CERN

- 2012: pion and kaon polarisabilities (Primakoff) + **comissioning and pilot run for DVCS**
- 2013-2014: long SPS/LHC shutdown
- 2014-2015: **Drell-Yann measurements with transversely polarised protons (NH<sub>3</sub> target)**
- 2016-2017: **stage 1 of GPD program and in parallel SIDIS (LH target)**


Measurements to be pursued at COMPASS-II > 2017 (subject to an Addendum)

- ✓ **Drell-Yann on transversely polarised protons, transversely polarised deuterons, unpolarised protons and nuclear targets**
- ✓ **stage 2 of GPD program with transversely polarised NH<sub>3</sub> target and RPD**
- ✓ SIDIS (high statistics) from transversely polarised deuteron and proton targets
- ✓ hadron program (spectroscopy in diffractive and central production, hybrids and exotics)



# What makes COMPASS unique for GPD studies

CERN SPS high energy polarised muon beam



- ✓ 100 – 190 GeV
- ✓ polarisation 80%
- ✓  $\mu^+$  and  $\mu^-$  available
  - opposite polarisation
  - $3.9 \cdot 10^8 \mu^+$  /spill
  - $I(\mu^+) \approx 2.4 I(\mu^-)$
- ✓  $L = 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$   
with 2.5 m long LH<sub>2</sub> target

## Foreseen measurements

DVCS and HEMP off unpolarised and transversely polarised protons

## Kinematic range for DVCS

$$Q^2 \rightarrow 8 \text{ GeV}^2$$


$$\sim 10^{-2} < x < \sim 10^{-1}$$

$x \rightarrow 0.27$  with extension of present calorimetry





# Interplay of DVCS and BH at 160 GeV


DVCS :



Bethe-Heitler :




$0.005 < x_{Bj} < 0.01$




BH dominates  
excellent  
reference yield

$0.01 < x_{Bj} < 0.03$



BH and DVCS at the same level  
access to DVCS amplitude  
through the interference

$x_{Bj} > 0.03$




DVCS dominates  
study of  $d\sigma^{DVCS}/dt$

Monte-Carlo  
Simulation  
for COMPASS  
set-up with  
only ECAL1+2

← non-uniform  
DVCS acceptance  
without ECAL0


# The COMPASS set-up for the GPD program



ECAL1


ECAL2

Main new equipments




2.5m-long  
Liquid H<sub>2</sub>  
Target



Target TOF System  
24 inner & outer scintillators  
1 GHz SADC readout  
goal: **310 ps** TOF resol.



ECAL0 Calorimeter  
Shashlyk modules + MAPD readout  
~ 2 × 2 m<sup>2</sup>, ~2200 ch.



# Large-angle electromagnetic calorimeter ECAL0




## Photo-diodes MAPD



$3 \times 3 \text{ mm}^2$   
Nb of pixels  $\sim 135\ 000$


## ECAL0 cell

252mm or 15 radiation length



shashlyk technology  
109 plates made of Sc 1.5 mm /Pb 0.8 mm

## Single module of 3x3 cells



56 mod. available for 2012 run  
complete ECAL0 (194 mod.)  
scheduled for 2016-2017



## Mounting of CAMERA in clean area at CERN




# Selection of single photon events

reconstructed vertex in the target volume


$Q^2 > 1 \text{ GeV}^2$ ,  $0.05 < y < 0.9$

$0.06 \text{ GeV}^2 < |t| < 0.64 \text{ GeV}^2$


only 1 photon with energy above DVCS threshold




vertex pointing




$$\Delta\varphi = \varphi_{meas}^{prot} - \varphi_{recon}^{prot}$$





$$\Delta p_{\perp} = p_{\perp,meas}^{prot} - p_{\perp,recon}^{prot}$$



$$M_X^2 = (p_{in}^{\mu} + p_{in}^p - p_{out}^{\mu} - p_{out}^p - p^{\gamma})^2$$




# Exclusive $\gamma$ production from 2012 DVCS commissioning run



- dominant BH process at small  $x_{\text{BJ}}$  clearly visible
- shape of  $\phi$  distribution reproduced well by MC
- first estimates of  $\pi^0$  background contributing at large  $x_{\text{BJ}}$
- at large  $x_{\text{BJ}}$  an excess of events wrt BH + background



# Extraction of DVCS cross section and amplitude



## Beam Charge & Spin Sum

$$S_{CS,U} \equiv d\sigma(\mu^{+\downarrow}) + d\sigma(\mu^{-\uparrow}) = 2(\underline{d\sigma^{BH}} + d\sigma^{DVCS}_{unpol}) + e_{\mu} P_{\mu} a^{BH} Im T^{DVCS}$$

$$c_0^{DVCS} + c_1^{DVCS} \cos\phi + c_2^{DVCS} \cos 2\phi$$

$$s_1^{Int} \sin\phi + s_2^{Int} \sin 2\phi$$

$$c_0^{DVCS} \rightarrow d\sigma^{DVCS}/dt$$

$$s_1^{Int} \rightarrow Im(F_1 \mathcal{H})$$

$$Im \mathcal{H}(\xi, t) = \mathbf{H}(x = \xi, \xi, t)$$

## Beam Charge & Spin Difference

$$D_{CS,U} \equiv d\sigma(\mu^{+\downarrow}) - d\sigma(\mu^{-\uparrow}) = 2(e_{\mu} a^{BH} Re T^{DVCS} + P_{\mu} d\sigma^{DVCS}_{pol})$$

$$c_0^{Int} + c_1^{Int} \cos\phi + c_2^{Int} \cos 2\phi + c_3^{Int} \cos 3\phi$$

$$s_1^{DVCS} \sin\phi$$

$$c_{0,1}^{Int} \rightarrow Re(F_1 \mathcal{H})$$


$$Re \mathcal{H}(\xi, t) = \mathcal{P} \int dx \frac{\mathbf{H}(x, \xi, t)}{x - \xi} = \mathcal{P} \int dx \frac{\mathbf{H}(x, x, t)}{x - \xi} + \mathcal{D}(t)$$



# Transverse imaging of the proton using $d\sigma^{DVCS}/dt$

integrating  $S_{CS,U}$  over  $\phi$  and subtracting BH  $\rightarrow d\sigma_{DVCS}/dt \sim \exp(-B|t|)$

COMPASS-II projections for B-slope uncertainties




'tomography':  $B(x) \Leftrightarrow \langle r_T^2 \rangle(x)$

ansatz at small  $x_B$   
 inspired by  
 Regge Phenomenology:  
 $B(x_B) = b_0 + 2 \alpha' \ln(x_0/x_B)$   
 $\alpha'$  slope of Regge traject

from 4 weeks in 2012 DVCS test

with 40 weeks in 2015-16

➤ From 2012 data expected the first measurement of B-slope for DVCS at an  $x_{Bj}$  above HERA range




# Beam Charge & Spin Difference of cross sections


$$D_{CS,U} \equiv d\sigma(\mu^{+\downarrow}) - d\sigma(\mu^{-\uparrow}) =$$

$$c_0^{Int} + c_1^{Int} \cos \phi + c_2^{Int} \cos 2\phi + c_3^{Int} \cos 3\phi + s_1^{DVCS} \sin \phi$$

$$c_{0,1}^{Int} \rightarrow \text{Re}(F_1 \mathcal{H})$$



Systematic error: 3% mostly due to luminosity measurements for  $\mu^+$  and  $\mu^-$



- Kroll, Moutarde, Sabatié
- - - KM10a
- - - KM10b

## Summary

- COMPASS has a great potential for GPD physics
  - ✓ unique polarised  $\mu^+$  and  $\mu^-$  beams
  - ✓ favourable kinematic domain ( $x_{Bj}$ )
- Large projects for new apparatus
  - ✓ 4m RPD + large angle ECAL0 (phase 1)
  - ✓ recoil proton detector incorporated into a large polarised target (phase 2)
- Investigation of GPDs with both DVCS and HEMP on unpolarised nucleons
  - ✓ t-slope of DVCS and HEMP cross section as a function of  $x_{Bj}$   
→ transverse distribution of partons
  - ✓ Beam Charge&Spin sum and difference of DVCS cross sections  
→  $\text{Re } T^{\text{DVCS}}$  and  $\text{Im } T^{\text{DVCS}}$  for the GPD H determination
  - ✓ Production of vector mesons  $\rho^0$ ,  $\omega$ ,  $\phi$  ... → flavour separation for GPD H
  - ✓ Production of  $\pi^0$  → sensitivity to GPDs  $\tilde{E}$  and  $\bar{E}_T$  ( $\equiv 2\tilde{H}_T + E_T$ )
- Transverse Target Spin Asymmetries for DVCS and hard exclusive meson production
  - GPD E and angular momentum of partons
  - also for mesons investigation of chiral-odd GPDs


Backup

# Main goals of the GPD program

- GPD a 3-dimensional image of the partonic structure of the nucleon

$$H(x, \xi=0, t) \rightarrow H(x, r_{x,y})$$

probability interpretation  
Burkardt




- Contribution to the nucleon spin puzzle


E related to the orbital angular momentum

$$2J_q = \int x (H^q(x, \xi, 0) + E^q(x, \xi, 0)) dx$$

$$\frac{1}{2} = \frac{1}{2} \Delta\Sigma + \Delta G + \langle L_z^q \rangle + \langle L_z^g \rangle$$



# Generalized Parton Distributions and DVCS




4 Generalised Parton Distributions : H, E,  $\tilde{H}$ ,  $\tilde{E}$  (chiral even)


for each quark flavour and for gluons

depend on 3 variables:  $x, \xi, t$



for DVCS gluons contribute at higher orders in  $\alpha_s$



# Role of pion exchange



- Effect known since early photoproduction experiments
- At COMPASS kinematics:
  - small for  $\rho^0$  production
  - sizable for  $\omega$  production
- Unnatural parity exchange process
  - impact on helicity-dependent observables
- Crucial for description of SDMEs for excl.  $\omega$  production
  - Goloskokov and Kroll, Eur. Phys. J. A50 (2014) 9, 146
- Sign of  $\pi\omega$  form factor not resolved from SDMEs data
  - azimuthal asymmetries more sensitive



@  $W=4.8$  GeV,  $Q^2=2.42$  GeV<sup>2</sup>


- positive  $\pi\omega$  form factor
- - - no pion pole
- ⋯ negative  $\pi\omega$  form factor

@  $W=8$  GeV,  $Q^2=2.42$  GeV<sup>2</sup>

- · - · - positive  $\pi\omega$  form factor
- ⋯ negative  $\pi\omega$  form factor



# Azimuthal dependence of exclusive photon xsec.



from Belitsky, Kirchner, Müller :  
polarized beam off unpolarized target

$$d\sigma_{(\mu p \rightarrow \mu p \gamma)} = d\sigma^{BH} + d\sigma^{DVCS}_{unpol} + P_\mu d\sigma^{DVCS}_{pol} + e_\mu a^{BH} \Re A^{DVCS} + e_\mu P_\mu a^{BH} \Im A^{DVCS}$$

$$d\sigma^{BH} = \frac{\Gamma(x_B, Q^2, t)}{P_1(\varphi)P_2(\varphi)} (c_0^{BH} + c_1^{BH} \cos \varphi + c_2^{BH} \cos 2\varphi) \leftarrow \text{Known expression}$$

$$d\sigma^{DVCS}_{unpol} = \frac{e^6}{y^2 Q^2} (c_0^{DVCS} + c_1^{DVCS} \cos \varphi + c_2^{DVCS} \cos 2\varphi)$$

$$P_\mu \times d\sigma^{DVCS}_{pol} = \frac{e^6}{y^2 Q^2} (s_1^{DVCS} \sin \varphi)$$

$$e_\mu \times a^{BH} \Re A^{DVCS} = \frac{e^6}{xy^3 + P_1(\varphi)P_2(\varphi)} (c_0^{Int} + c_1^{Int} \cos \varphi + c_2^{Int} \cos 2\varphi + c_3^{Int} \cos 3\varphi)$$

$$e_\mu P_\mu \times a^{BH} \Im A^{DVCS} = \frac{e^6}{xy^3 + P_1(\varphi)P_2(\varphi)} (s_1^{Int} \sin \varphi + s_2^{Int} \sin 2\varphi)$$

Twist-2

>>

Twist-3

Twist-2 gluon

# Study of azimuthal asymmetries from transversely polarized NH<sub>3</sub> target

$$\mathcal{D}_{CS,T} \equiv d\sigma_T(\mu^{+\downarrow}) - d\sigma_T(\mu^{-\uparrow})$$

$$\propto \text{Im}(F_2 \mathcal{H} - F_1 \mathcal{E}) \sin(\phi - \phi_S) \cos \phi + \dots$$

$$\mathcal{A}_{CS,T}^D \equiv \mathcal{D}_{CS,T} / d\sigma_0$$




$d\sigma_0$  - unpolarised, charge averaged cross section

160 GeV muon beam  
 1.2m NH<sub>3</sub> target  
 $\epsilon_{\text{global}} = 10\%$   
 with ECAL1+ ECAL2  
 40 weeks

for  $\mu p^\uparrow \rightarrow \mu \gamma p$  from NH<sub>3</sub>  
 dilution factor  $f=0.26$


$0.10 (0.14) < |t| < 0.64 \text{ GeV}^2$

## COMPASS-II proposal



# t-slope measurement for exclusive $\rho^0$ production


$$d\sigma_{\gamma N \rightarrow \rho N}/dt \sim \exp(-B|t|)$$



At large  $Q^2$  slope  $B$  sensitive mostly to the nucleon size

160 GeV muon beam  
2.5m LH<sub>2</sub> target  
 $\epsilon_{\text{global}} = 10\%$ , 280 days  
 $L = 1222\text{pb}^{-1}$

$$0.06 < |t| < 0.64 \text{ GeV}^2$$



$$\mu^2 = (Q^2 + M_V^2)/4$$

( =  $Q^2$  for DVCS )