

# Unpolarised SIDIS studies at COMPASS

Jan Matoušek

Charles University, Prague, Czechia  
on behalf of the COMPASS collaboration

18. 11. 2020

International Workshop on Hadron Structure and Spectroscopy



CHARLES UNIVERSITY  
Faculty of mathematics  
and physics





1 Introduction

2 Published measurements on  ${}^6\text{LiD}$

3 New measurements on  $\text{LH}_2$

4 Conclusion




## 1 Introduction

## 2 Published measurements on ${}^6\text{LiD}$



## 3 New measurements on $\text{LH}_2$

## 4 Conclusion



It is located at M2 beamline of CERN's SPS.

- Collaboration: 24 institutes, 13 countries.
- Fixed target, multi-purpose.
- Broad research programme:
  - **SIDIS**:  $\mu^+$  beam and L/T-polarised proton ( $\text{NH}_3$ ) or deuteron ( ${}^6\text{LiD}$ ) target (beam 160 GeV/c, 200 GeV/c in 2011)
  - **Hadron spectroscopy**: hadron beams and nuclear targets.
  - **Drell-Yan**: 190 GeV/c  $\pi^-$  beam and  $p^\uparrow$ , Al and W targets.
  - **DVCS and SIDIS**: 160 GeV/c  $\mu^\pm$  beam and liquid  $\text{H}_2$  target.



2016–2017 setup with CAMERA recoil proton detector and ECAL0 calorimeter for DVCS studies.




The cross section for producing a hadron  $h$  in DIS on unpolarised target  $\ell N \rightarrow \ell' h X$ :

[A. Bacchetta *et al.*, JHEP 0702 (2007)]

$$\begin{aligned} \frac{d\sigma}{dx dy dz d\phi_h dP_T^2} &= \frac{2\pi\alpha^2}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{2xM^2}{Q^2}\right) \left(F_{UU,T} + \varepsilon F_{UU,L}\right. \\ &\quad \left. + \sqrt{2\varepsilon(1+\varepsilon)} F_{UU}^{\cos\phi_h} \cos\phi_h + \varepsilon F_{UU}^{\cos 2\phi_h} \cos 2\phi_h + \lambda\sqrt{2\varepsilon(1-\varepsilon)} F_{LU}^{\sin\phi_h} \sin\phi_h\right) \\ &= \sigma_0 \left(1 + \varepsilon_1 A_{UU}^{\cos\phi_h} \cos\phi_h + \varepsilon_2 A_{UU}^{\cos 2\phi_h} \cos 2\phi_h + \lambda\varepsilon_3 A_{LU}^{\sin\phi_h} \sin\phi_h\right) \end{aligned}$$

- where  $x$ ,  $y$ ,  $Q^2$  are usual DIS variables,
- $\lambda$  is the beam polarisation ( $\approx 0.8$  at COMPASS),
- $\varepsilon \approx \frac{1-y}{1-y+\frac{1}{2}y^2}$ ,  $M$  nucleon mass,
- $z$  is the fraction of  $\gamma^*$  energy carried by  $h$ .
- $P_T$  is the transverse momentum of  $h$  in the  $\gamma N$  frame,  $\phi_h$  is its azimuthal angle.
- $F_{XU}^{f(\phi_h)}(x, z, P_T^2, Q^2)$  are structure functions.
- $A_{XU}^{f(\phi_h)}(x, z, P_T^2, Q^2)$  are commonly called azimuthal asymmetries.



SIDIS in the  $\gamma$ -nucleon frame.




The cross section for producing a hadron  $h$  in DIS on unpolarised target  $\ell N \rightarrow \ell' h X$ :

[A. Bacchetta *et al.*, JHEP 0702 (2007)]

$$\begin{aligned} \frac{d\sigma}{dx dy dz d\phi_h dP_T^2} &= \frac{2\pi\alpha^2}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{2xM^2}{Q^2}\right) \left(F_{UU,T} + \varepsilon F_{UU,L}\right. \\ &\quad \left. + \sqrt{2\varepsilon(1+\varepsilon)} F_{UU}^{\cos\phi_h} \cos\phi_h + \varepsilon F_{UU}^{\cos 2\phi_h} \cos 2\phi_h + \lambda\sqrt{2\varepsilon(1-\varepsilon)} F_{LU}^{\sin\phi_h} \sin\phi_h\right) \\ &= \sigma_0 \left(1 + \varepsilon_1 A_{UU}^{\cos\phi_h} \cos\phi_h + \varepsilon_2 A_{UU}^{\cos 2\phi_h} \cos 2\phi_h + \lambda\varepsilon_3 A_{LU}^{\sin\phi_h} \sin\phi_h\right) \end{aligned}$$

- where  $x$ ,  $y$ ,  $Q^2$  are usual DIS variables,
- $\lambda$  is the beam polarisation ( $\approx 0.8$  at COMPASS),
- $\varepsilon \approx \frac{1-y}{1-y+\frac{1}{2}y^2}$ ,  $M$  nucleon mass,
- $z$  is the fraction of  $\gamma^*$  energy carried by  $h$ .
- $P_T$  is the transverse momentum of  $h$  in the  $\gamma N$  frame,  $\phi_h$  is its azimuthal angle.
- $F_{XU}^{f(\phi_h)}(x, z, P_T^2, Q^2)$  are structure functions.
- $A_{XU}^{f(\phi_h)}(x, z, P_T^2, Q^2)$  are commonly called azimuthal asymmetries.



SIDIS in the  $\gamma$ -nucleon frame.



The structure functions in terms of TMD PDFs and TMD FFs, up to order  $1/Q$ :

$$F_{UU,T} = C [f_1 D_1],$$


$$F_{UU,L} = 0, \quad \text{Cahn effect}$$

$$F_{UU}^{\cos \phi_h} = \frac{2M}{Q} C \left[ -\frac{\hat{h} \cdot \mathbf{k}_T}{M} f_1 D_1 - \frac{(\hat{h} \cdot \mathbf{p}_\perp) k_T^2}{M^2 M_h} h_1^\perp H_1^\perp + \dots \right]$$

$$F_{UU}^{\cos 2\phi_h} = C \left[ -\frac{2(\hat{h} \cdot \mathbf{k}_T)(\hat{h} \cdot \mathbf{p}_\perp) - \mathbf{k}_T \cdot \mathbf{p}_\perp}{MM_h} h_1^\perp H_1^\perp \right]$$

$$F_{UU}^{\sin \phi_h} = \frac{2M}{Q} C [\dots]$$

- $f_1(x, k_T^2, Q^2)$  unpolarised TMD PDF,
- $h_1^\perp(x, k_T^2, Q^2)$  Boer–Mulders function,
- $D_1(z, p_\perp^2, Q^2)$  unpolarised TMD FF,
- $H_1^\perp(z, p_\perp^2, Q^2)$  Collins function.
- $\hat{h} = \mathbf{P}_T / P_T$ ,
- $C$  = sum over flavours and convolution over  $\mathbf{p}_\perp, \mathbf{k}_T$ ,
- ... = twist-three terms.



Observables sensitive to  $k_T$  and  $p_\perp$ :

- azimuthal asymmetries  
 $A_{UU}^{\cos \phi_h}, A_{UU}^{\cos 2\phi_h}, A_{UU}^{\sin \phi_h}$ ,
  - $k_T$  via Cahn effect,
  - Boer–Mulders function.
- $P_T$ -dependent distributions  
 $\propto F_{UU,T} = C[f_1 D_1]$ .



The structure functions in terms of TMD PDFs and TMD FFs, up to order  $1/Q$ :

$$F_{UU,T} = C [f_1 D_1],$$


$$F_{UU,L} = 0, \quad \text{Cahn effect}$$

$$F_{UU}^{\cos \phi_h} = \frac{2M}{Q} C \left[ -\frac{\hat{h} \cdot \mathbf{k}_T}{M} f_1 D_1 - \frac{(\hat{h} \cdot \mathbf{p}_\perp) k_T^2}{M^2 M_h} h_1^\perp H_1^\perp + \dots \right]$$

$$F_{UU}^{\cos 2\phi_h} = C \left[ -\frac{2(\hat{h} \cdot \mathbf{k}_T)(\hat{h} \cdot \mathbf{p}_\perp) - \mathbf{k}_T \cdot \mathbf{p}_\perp}{MM_h} h_1^\perp H_1^\perp \right]$$

$$F_{UU}^{\sin \phi_h} = \frac{2M}{Q} C [\dots]$$

- $f_1(x, k_T^2, Q^2)$  unpolarised TMD PDF,
- $h_1^\perp(x, k_T^2, Q^2)$  Boer–Mulders function,
- $D_1(z, p_\perp^2, Q^2)$  unpolarised TMD FF,
- $H_1^\perp(z, p_\perp^2, Q^2)$  Collins function.
- $\hat{h} = \mathbf{P}_T / P_T$ ,
- $C$  = sum over flavours and convolution over  $\mathbf{p}_\perp, \mathbf{k}_T$ ,
- ... = twist-three terms.



Observables sensitive to  $k_T$  and  $p_\perp$ :

- azimuthal asymmetries  
 $A_{UU}^{\cos \phi_h}, A_{UU}^{\cos 2\phi_h}, A_{UU}^{\sin \phi_h}$ ,
  - $k_T$  via Cahn effect,
  - Boer–Mulders function.
- $P_T$ -dependent distributions  
 $\propto F_{UU,T} = C[f_1 D_1]$ .





## Published unpolarised SIDIS results:

- Azimuthal asymmetries on  ${}^6\text{LiD}$  target [COMPASS, Nucl.Phys.B 886 (2014)].
- $P_T$ -dependent multiplicities on  ${}^6\text{LiD}$  target [COMPASS, Phys.Rev.D97 (2018)]
- Background to the asymmetries from decays of exclusive vector mesons [COMPASS, Nucl.Phys.B 956 (2020)].

## Ongoing analysis presented in this talk:

- 2016–2017 data taken with 2.5 m long  $\text{LH}_2$  target.
- Primary goal: DVCS measurement, but useful for SIDIS as well.
- Advantages:
  - pure proton target,
  - alternating  $\mu^\pm$  beam with balanced statistics (stability tests for systematics),
  - MC development in synergy with DVCS analysis.
- Part of the data (about 11 %) used for preliminary results, released in August 2020.

## Future:

- 2021 run with  ${}^6\text{LiD}$  target (transversely polarised).



## Published unpolarised SIDIS results:

- Azimuthal asymmetries on  ${}^6\text{LiD}$  target [COMPASS, Nucl.Phys.B 886 (2014)].
- $P_T$ -dependent multiplicities on  ${}^6\text{LiD}$  target [COMPASS, Phys.Rev.D97 (2018)]
- Background to the asymmetries from decays of exclusive vector mesons [COMPASS, Nucl.Phys.B 956 (2020)].

## Ongoing analysis presented in this talk:

- 2016–2017 data taken with 2.5 m long  $\text{LH}_2$  target.
- Primary goal: DVCS measurement, but useful for SIDIS as well.
- Advantages:
  - pure proton target,
  - alternating  $\mu^\pm$  beam with balanced statistics (stability tests for systematics),
  - MC development in synergy with DVCS analysis.
- Part of the data (about 11 %) used for preliminary results, released in August 2020.

## Future:

- 2021 run with  ${}^6\text{LiD}$  target (transversely polarised).



## Published unpolarised SIDIS results:

- Azimuthal asymmetries on  ${}^6\text{LiD}$  target [COMPASS, Nucl.Phys.B 886 (2014)].
- $P_T$ -dependent multiplicities on  ${}^6\text{LiD}$  target [COMPASS, Phys.Rev.D97 (2018)]
- Background to the asymmetries from decays of exclusive vector mesons [COMPASS, Nucl.Phys.B 956 (2020)].

## Ongoing analysis presented in this talk:

- 2016–2017 data taken with 2.5 m long  $\text{LH}_2$  target.
- Primary goal: DVCS measurement, but useful for SIDIS as well.
- Advantages:
  - pure proton target,
  - alternating  $\mu^\pm$  beam with balanced statistics (stability tests for systematics),
  - MC development in synergy with DVCS analysis.
- Part of the data (about 11 %) used for preliminary results, released in August 2020.

## Future:

- 2021 run with  ${}^6\text{LiD}$  target (transversely polarised).



1 Introduction




2 Published measurements on  ${}^6\text{LiD}$

3 New measurements on  $\text{LH}_2$

4 Conclusion




- [COMPASS, Nucl.Phys.B 886 (2014)]
- Isoscalar target, effectively deuteron.
- Unidentified charged hadrons studied.
- 1D analysis  
(bins in  $x$ ,  $z$  and  $P_T$  separately).
- 3D analysis (3D grid of bins).
- Strong kinematic dependence of the  $\cos \phi_h$  and  $\cos 2\phi_h$  asymmetries.
- At the time, some features were not understood (e.g. positive  $A_{UU}^{\cos \phi_h}$ )
- Exclusive vector meson contribution has been proved important later.






- [COMPASS, Nucl.Phys.B 886 (2014)]
- Isoscalar target, effectively deuteron.
- Unidentified charged hadrons studied.
- 1D analysis  
(bins in  $x$ ,  $z$  and  $P_T$  separately).
- 3D analysis (3D grid of bins).
- Strong kinematic dependence of the  $\cos \phi_h$  and  $\cos 2\phi_h$  asymmetries.
- At the time, some features were not understood (e.g. positive  $A_{UU}^{\cos \phi_h}$ )
- Exclusive vector meson contribution has been proved important later.







- [COMPASS, Phys.Rev.D97 (2018)]
- **4D analysis**  
(bins in  $x$ ,  $Q^2$ ,  $z$  and  $P_T^2$ )
- Unidentified charged hadrons studied.
- QED radiative effects taken into account.
- Contribution of the decay of exclusive vector mesons
  - Contamination estimated from HEPGEN MC generator [A. Sandacz & P. Sznajder, arXiv:1207.0333].
  - Subtracted in each bin.
  - $\rho^0$ : small  $P_T$ , large  $z$ , small  $Q^2$ .
  - $\phi$ : tiny  $P_T$ , medium  $z$ , small  $Q^2$ .






- [COMPASS, Phys.Rev.D97 (2018)]
- 4D analysis (bins in  $x$ ,  $Q^2$ ,  $z$  and  $P_T^2$ )
- Unidentified charged hadrons studied.
- QED radiative effects taken into account.
- Contribution of the decay of exclusive vector mesons
  - Contamination estimated from HEPGEN MC generator [A.Sandacz & P.Sznajder, arXiv:1207.0333].
  - Subtracted in each bin.
  - $\rho^0$ : small  $P_T$ , large  $z$ , small  $Q^2$ .
  - $\phi$ : tiny  $P_T$ , medium  $z$ , small  $Q^2$ .



$1 - \rho^0$  contamination fraction.






Diffractively produced  $\rho^0 \rightarrow \pi^+\pi^-$ , creating a background to SIDIS.








- [COMPASS, Nucl.Phys.B 956 (2020)].
- The exclusive VMs inherit  $\gamma^*$  polarisation.
- The decay hadrons obtain large azimuthal modulations. Especially in  $\cos\phi_h$ .
- They were measured in the data selecting
  - only  $\mu'h^+h^-$ ,
  - $z_1 + z_2 > 0.95$ .
- The contamination fraction from HEPGEN.
- Subtraction at the asymmetry level.


Total  $z$  for  $h^+h^-$ . $\phi_h$ - $z$  correlation.




- [COMPASS, Nucl.Phys.B 956 (2020)].
- The exclusive VMs inherit  $\gamma^*$  polarisation.
- The decay hadrons obtain large azimuthal modulations. Especially in  $\cos\phi_h$ .
- They were measured in the data selecting
  - only  $\mu'h^+h^-$ ,
  - $z_1 + z_2 > 0.95$ .
- The contamination fraction from HEPGEN.
- Subtraction at the asymmetry level.


Total  $z$  for  $h^+h^-$ .

Before (empty) and after (full) subtraction.  
 $0.1 < P_T/(\text{GeV}/c) < 0.3$ .



The contamination fraction:  $3D(P_T, z, x)$  representation.





After exclusive VM decay contribution is subtracted, rather good agreement with an MC model based on Cahn effect and string fragmentation [A. Kerbizi *et al.*, Phys.Rev.D 97 (2018)] can be reached [COMPASS, Nucl.Phys.B 956 (2020)].





1 Introduction

2 Published measurements on  ${}^6\text{LiD}$

3 New measurements on  $\text{LH}_2$

4 Conclusion





The  $x$  and  $Q^2$  range covered.

## DIS event selection


- $Q^2 > 1 \text{ (GeV}/c)^2$ ,
- $W > 5 \text{ GeV}/c^2$ ,
- $0.003 < x < 0.13$ ,
- $0.2 < y < 0.9$ ,
- $\theta_\gamma < 60 \text{ mrad}$ ,
- Exclusive VM decay cut:  
if only  $\mu'h^+h^-$  outgoing,  
 $z_1 + z_2 = z_t < 0.95$ .

## Hadron selection

- $0.1 < z < 0.85$ ,
- $0.1 < P_T/(\text{GeV}/c) < 1.73$ .



The  $x$  and  $Q^2$  range covered.





Selected range with moderate acceptance corrections.

## DIS event selection


- $Q^2 > 1 \text{ (GeV/c)}^2$ ,
- $W > 5 \text{ GeV/c}^2$ ,
- $0.003 < x < 0.13$ ,
- $0.2 < y < 0.9$ ,
- $\theta_\gamma < 60 \text{ mrad}$ ,
- Exclusive VM decay cut:  
if only  $\mu'h^+h^-$  outgoing,  
 $z_1 + z_2 = z_t < 0.95$ .

## Hadron selection

- $0.1 < z < 0.85$ ,
- $0.1 < P_T / (\text{GeV/c}) < 1.73$ .



The  $x$  and  $Q^2$  range covered.




Selected range with moderate acceptance corrections.

## DIS event selection



- $Q^2 > 1$  (GeV/c)<sup>2</sup>,
- $W > 5$  GeV/c<sup>2</sup>,
- $0.003 < x < 0.13$ ,
- $0.2 < y < 0.9$ ,
- $\theta_\gamma < 60$  mrad,
- Exclusive VM decay cut: if only  $\mu'h^+h^-$  outgoing,  $z_1 + z_2 = z_t < 0.95$ .

## Hadron selection

- $0.1 < z < 0.85$ ,
- $0.1 < P_T / (\text{GeV}/c) < 1.73$ .



[COMPASS, Nucl.Phys.B 956 (2020)]




$Q^2$  and  $x$  bins for the  $P_T$ -dependent distributions.


## Binning

- Based on the published results.
- Asymmetries:
  - 1D in  $x$ ,  $z$  and  $P_T$ .
  - 3D in  $x$ ,  $z$  and  $P_T$
- $P_T$ -dependent distributions
  - 4D in  $x$ ,  $Q^2$ ,  $z$  and  $P_T^2$ .
  - Larger bins w.r.t the publication (2 bins in every variable merged).







- Different approach w.r.t published d asymmetries.
- ‘Visible’ exclusive  $h^+h^-$  removed in event selection.
  - About 80 % of the decays are ‘visible’.
- ‘Invisible’ decays (only one h observed)
  - HEPGEN MC generator with azimuthal modulations.
  - Normalised to the data using  $E_{\text{miss}}$  distribution of the ‘visible’ decays.
  - Subtracted in every bin (including  $\phi_h$  bins).





The number of signal events in the peak after SIDIS (from LEPTO) background subtraction is used to normalise HEPGEN.




The VM-contamination fraction.




The azimuthal modulations of hadrons from the 'visible' VM decays. The 'invisible' ones have very similar modulations.





The impact of the VM-subtraction ('visible' and 'invisible') on the  $P_T$ -dependent distributions.



- Acceptance correction
  - LEPTO generator, full Geant simulation of COMPASS.
- QED radiative effects – not yet taken into account
  - Plan to use DJANGO generator [DJANGO6] (→ evaluate impact on hadronic variables as well)
- 1D results
  - Strong kinematic dependences, differences between  $h^\pm$ ,
  - qualitative agreement with published deuteron results [COMPASS, Nucl.Phys.B 886 (2014)].





# New measurements on LH<sub>2</sub>: Results for the asymmetries



Qualitative agreement with published deuteron results [COMPASS, Nucl.Phys.B 956 (2020)].






- The distributions are normalised to the first bin.
- Gaussian model for  $f_1$  and  $D_1$ :





$$\frac{d^2 N}{dz dP_T} \propto \exp\left(-\frac{P_T^2}{\langle P_T^2 \rangle}\right)$$

$$\langle P_T^2 \rangle = z^2 \langle k_T^2 \rangle + \langle p_\perp^2 \rangle.$$

- Deviation from the simple exponential visible at  $P_T > 1$  (GeV/c)<sup>2</sup>.





# New measurements on LH<sub>2</sub>: Results for the $P_T$ -distributions





$$\frac{d^2N}{dzdP_T} \propto \exp\left(-\frac{P_T^2}{\langle P_T^2 \rangle}\right)$$



Exponential fit in  $P_T < 1$  (GeV/c)<sup>2</sup> range.

- Deviations from the linear trend expected from the simple Gaussian model



$$\langle P_T^2 \rangle = z^2 \langle k_T^2 \rangle + \langle p_{\perp}^2 \rangle.$$

- Possible dependence of  $\langle p_{\perp}^2 \rangle$  on  $z$  or of  $\langle k_T^2 \rangle$  on  $x$ .






$$\frac{d^2N}{dzdP_T} \propto \exp\left(-\frac{P_T^2}{\langle P_T^2 \rangle}\right)$$



Exponential fit in  $P_T < 1$  (GeV/c)<sup>2</sup> range.



The fitted  $\langle P_T^2 \rangle$  versus  $z^2$  in the  $x$  and  $Q^2$  bins.

- Deviations from the linear trend expected from the simple Gaussian model

$$\langle P_T^2 \rangle = z^2 \langle k_T^2 \rangle + \langle p_{\perp}^2 \rangle.$$

- Possible dependence of  $\langle p_{\perp}^2 \rangle$  on  $z$  or of  $\langle k_T^2 \rangle$  on  $x$ .



1 Introduction

2 Published measurements on  ${}^6\text{LiD}$

3 New measurements on  $\text{LH}_2$

4 Conclusion



## Interesting observables in unpolarised SIDIS

- **Azimuthal asymmetries:** sensitive to  $k_T$  (via Cahn effect) and to Boer–Mulders function.
- **$P_T$ -dependent distributions:** sensitive to  $k_T$  and  $p_\perp$  dependence of  $f_1$  and  $D_1$ .
- Contamination from decays of exclusive VMs plays an important role in both measurements.

## COMPASS measurements

- Published results on  ${}^6\text{LiD}$  target: [COMPASS, Nucl.Phys.B 886 (2014)], [COMPASS, Phys.Rev.D97 (2018)], [COMPASS, Nucl.Phys.B 956 (2020)].
- New preliminary results (August 2020) on liquid  $\text{H}_2$  target.
  - 11 % of the statistics,
  - More robust method for exclusive VM subtraction.
  - Alternating  $\mu^\pm$  beam – systematic check.
  - Qualitative agreement with deuteron, rich kinematic dependences.
  - More results will come.
- Scheduled 2021 measurement with (transversely polarised)  ${}^6\text{LiD}$  target.

These measurements provide important input to general understanding of the transverse-momentum-dependent structure of the nucleon and of the fragmentation process.



## Interesting observables in unpolarised SIDIS




- **Azimuthal asymmetries:** sensitive to  $k_T$  (via Cahn effect) and to Boer–Mulders function.
- **$P_T$ -dependent distributions:** sensitive to  $k_T$  and  $p_\perp$  dependence of  $f_1$  and  $D_1$ .
- Contamination from decays of exclusive VMs plays an important role in both measurements.

## COMPASS measurements

- Published results on  ${}^6\text{LiD}$  target: [COMPASS, Nucl.Phys.B 886 (2014)], [COMPASS, Phys.Rev.D97 (2018)], [COMPASS, Nucl.Phys.B 956 (2020)].
- New preliminary results (August 2020) on liquid  $\text{H}_2$  target.
  - 11 % of the statistics,
  - More robust method for exclusive VM subtraction.
  - Alternating  $\mu^\pm$  beam – systematic check.
  - Qualitative agreement with deuteron, rich kinematic dependences.
  - More results will come.
- Scheduled 2021 measurement with (transversely polarised)  ${}^6\text{LiD}$  target.

These measurements provide important input to general understanding of the transverse-momentum-dependent structure of the nucleon and of the fragmentation process.

Thank you for your attention!



Normalised kinematic distributions: real data, LEPTO, HEPGEN  $\rho^0$  and HEPGEN  $\phi$ .

