


## $J/\psi$ pair production in $\pi N$ collisions at COMPASS



**Gridin Andrei (JINR)**  
*On behalf of the COMPASS collaboration*  
**[andrei.gridin@cern.ch](mailto:andrei.gridin@cern.ch)**  
**25.03.2020**



# First evidence of double $J/\psi$ production

**The NA3** double  $J/\psi$  results:

- $\pi^-$  (150, 280 GeV) and  $p$  (400 GeV) beams;
- N.B. kinematic distributions are not corrected for the acceptance;
- interpreted using single parton scattering mechanism ( $q\bar{q} \rightarrow 2J/\psi$  and  $gg \rightarrow 2J/\psi$ );
- interpreted using intrinsic charm hypothesis ( $|d\bar{u}c\bar{c}c\bar{c}\rangle$  Fock component of pion materialization).


**R.E. Ecclestone, D.M. Scott, Phys. Lett. B. V. 120. 1983**  
**B. Humpert, P. Mery, Phys. Lett. B. V. 124. 1983**

**R Vogt, S.J. Brodsky, Phys. Lett. B, v349: 569-575, 1995**

**J.Badier et al (NA3)**

**Phys. Lett. B, v114, No6,1982,**

**Phys. Lett. B, v158, No1,1985**




# Intrinsic charm of hadron

BHPS model: **S.J. Brodsky, Phys. Lett. B 93, 451 (1980)**

- The existence of non-perturbative (intrinsic) Fock component in a hadron with  $c$ -quarks is postulated:

$$|p\rangle = a_0 |uud\rangle + a_1 |uudg\rangle + \underline{a_2 |uudc\bar{c}\rangle} + \dots$$

- In perturbative QCD the extrinsic charm component in hadrons arises from gluon splitting.
- Intrinsic charm contribution is generated non-perturbatively via  $gg \rightarrow Q\bar{Q}$ ;



**V.A. Bednyakov, G.I. Lykasov  
Phys. Lett. B, 728, 602 (2014)**


# $T_{4c}$ -tetraquarks

Y. Iwasaki, Prog. Theor. Phys. V.54, p492 (1975)

- first time  $T_{4c}$  states were predicted in 1975;
- many theoretical models ( $[cc][\bar{c}\bar{c}]$  model, Drell-Yan type mechanism, etc) exist;
- $M_{T_{4c}} \approx 6 - 7 \text{ GeV}/c^2$ ;
- no experimental observations of  $T_{4c}$  till 2020;


$N^{2S+1}\ell_J$	$M_{T_{4c}}$	$J^{PC}$
<b>Diquark</b>		
$1^3S_1$	3133.4	$1^+$
<b>Tetraquark</b>		
$1^1S_0$	5969.4	$0^{++}$
$1^3S_1$	6020.9	$1^{+-}$
$1^5S_2$	6115.4	$2^{++}$
$1^1P_1$	6577.1	$1^{--}$
$1^3P_0$	6480.4	$0^{-+}$
$1^3P_1$	6577.4	$1^{-+}$
$1^3P_2$	6609.9	$2^{-+}$
$1^5P_1$	6495.4	$1^{--}$
$1^5P_2$	6600.2	$2^{--}$
$1^5P_3$	6641.2	$3^{--}$
$2^1S_0$	6663.3	$0^{++}$
$2^3S_1$	6674.5	$1^{+-}$
$2^5S_2$	6698.1	$2^{++}$
$2^1P_1$	6944.1	$1^{--}$
$2^3P_0$	6866.5	$0^{-+}$
$2^3P_1$	6943.9	$1^{-+}$
$2^3P_2$	6970.4	$2^{-+}$
$2^5P_1$	6875.6	$1^{--}$
$2^5P_2$	6962.1	$2^{--}$
$2^5P_3$	6996.7	$3^{--}$

V.R.Debastiani, F.S.Navarra  
Chinese Phys. C 43 013105



$[cc][\bar{c}\bar{c}]$  model

Bing-An Li, Keh-Fei Liu  
Phys. Rev. D 29, Vol 3, 1984



Drell-Yan type mechanism

# LHCb double $J/\psi$ results

The LHCb reported the narrow X(6900) structure in the double  $J/\psi$  mass spectrum using proton-proton collision data at  $\sqrt{s} = 7, 8$  and 13 TeV.

**Model I - X(6900) resonance (without interference):**

$$m[X(6900)] = 6905 \pm 11 \pm 7 \text{ MeV}/c^2$$

$$\Gamma[X(6900)] = 80 \pm 19 \pm 33 \text{ MeV}$$

**5.1  $\sigma$**



**Model II - X(6900) + interference between broad structure (6.2-6.8) and SPS:**

$$m[X(6900)] = 6886 \pm 11 \pm 11 \text{ MeV}/c^2$$

$$\Gamma[X(6900)] = 168 \pm 33 \pm 69 \text{ MeV}$$

**6.4  $\sigma$**

Sci. Bull., V65, №23, p1983-1993, 2020





# COMPASS experiment at CERN

COmmon MUon PProton Apparatus for Structure and Spectroscopy



## Phase 1:

- Nucleon Spin Structure (2002-2011)
- Hadron Spectroscopy (2008-2009)


## Phase 2:

- Primakoff (2012)
- DVCS, SIDIS (2012, 2016, 2017)
- Drell-Yan (2015, 2018)

CERN's Accelerator Complex




# COMPASS Drell-Yan setup



## Unique hadron beam in DY runs :

- hadron beam composition: 96.80%  $\pi^-$ , 2.40%  $\bar{K}$ , 0.80%  $\bar{p}$ ;
- beam momentum :  $190 \pm 3$  GeV/c;
- intensity: up to  $7 \times 10^7$  hadrons / sec;







# COMPASS Drell-Yan setup

## Polarized target:

- two 55 cm long cells filled with  $\text{NH}_3$  immersed in LHe used in particular in polarized DY studies.


## Nuclear targets (Al and W):

- used to remove hadrons originating from target interactions or beam;
- used as an additional nuclear targets:
  - aluminum ( $A \sim 27$ ): 7cm length;
  - tungsten (beam plug, 120 cm,  $A \sim 184$ ): first 10 cm used for the physics analyses.






# DY and $J/\psi$ studies at COMPASS



- Large statistics of single  $J/\psi$  event collected
- Mass resolution:  $\sigma_{J/\psi} = 0.181 \text{ GeV}/c^2$
- A shoulder from  $\psi(2S)$  is visible

# Double $J/\psi$ data at COMPASS



2015: ~4 months of data taking;  
2018: ~5 months of data taking;


NH<sub>3</sub> target: 25 events

- used for the analysis

Al target: 4 events


W target: 21 events

- large background contamination
- used only for cross-section estimation.



# Differential cross-sections

## COMPASS results:




## The NA3 results:

- Results were interpreted using double IC of pion hypothesis.



R Vogt, S.J. Brodsky  
 Phys.Lett.B349:569-575,1995

- N.B. Double  $J/\psi$  kinematic distributions were published without acceptance correction.



# Double $J/\psi$ mass spectrum


Sci. Bull., V65, N°23, p1983-1993, 2020



The COMPASS double  $J/\psi$  mass spectrum does not contain any evident signal from  $T_{4c}$  states.



# Double $J/\psi$ production mechanisms



## SPS curve:

- HELAC-Onia generator:  
**H.S.Shao, Comput. Phys. Commun., Vol. 198, p. 238-259, 2016;**
- Color Singlet  $J/\psi$  production model.


## IC curve:

- predictions for COMPASS from  
**Phys.Part.Nucl.Lett. Vol17, No6, 2020.**

## The SPS + Intrinsic Charm fit:

- the double parton scattering (DPS) is not considered in the fit;
- the DPS contribution at  $\sqrt{s} = 18.9$  GeV is less than 8% (**arXiv:1909.06195 [hep-ph]**);
- the data are consistent with pure SPS hypothesis.

# Double $J/\psi$ cross-section measurement



Main sources of systematics:

- uncertainty of  $\sigma_{J/\psi}$
- background estimation
- acceptance of double  $J/\psi$
- acceptance of single  $J/\psi$
- uncertainty of the number of single  $J/\psi$

$$\frac{\sigma_{2J/\psi}}{\sigma_{J/\psi}} \Big|_{x_F > 0} = (1.1 \pm 0.3_{stat} \pm 0.2_{syst}) \cdot 10^{-4} (NH3)$$

$$\sigma_{2J/\psi}^{NH3} \Big|_{x_F > 0} = 8.8 \pm 2.2_{stat} \pm 2.4_{syst} \frac{pb}{nucleon}$$

$$\sigma_{2J/\psi}^W \Big|_{x_F > 0} = 3.4 \pm 4.3_{stat} \pm 5.8_{syst} \frac{pb}{nucleon}$$

$$\sigma_{2J/\psi}^{Al} \Big|_{x_F > 0} = 14.3 \pm 7.7_{stat} \pm 4.5_{syst} \frac{pb}{nucleon}$$

COMPASS results do not contradict to NA3 values.

No A-dependence of  $\sigma_{2J/\psi}$  was found.

The measured by the NA3

$$\sigma_{J/\psi} = 4.9 \pm 0.77 \frac{nb}{nucleon}$$

was used for the estimation of  $\sigma_{2J/\psi}$

# Summary

1. Double  $J/\psi$  hadroproduction is a tool:
  - to study the intrinsic charm component of hadrons
  - to search for bound  $T_{4c}$  states.
2. The COMPASS collaboration:
  - has searched for double  $J/\psi$  events produced in  $\text{NH}_3$ , Al and W targets
  - has estimated double  $J/\psi$  production cross-section.
3. The COMPASS data are consistent with SPS production mechanism.
4. No evidence of presence of  $T_{4c}$  states in the double  $J/\psi$  mass spectrum.


**BACKUP**



# DPS/SPS ratio and generated SPS MC

J.-P. Lansberg, H.-S. Shao

Nucl. Phys. B 900 (2015) 273



- The DPS contribution is expected to be low at  $\sqrt{s} = 18.9$  GeV.

- The generated distributions for double  $J/\psi$  MC (SPS sample, HELAC-Onia generator):

