




# Meson Spectroscopy at COMPASS

Philipp Haas for the COMPASS Collaboration

19.03.2024 – PAW'24


# Motivation

- Mesons in the Constituent Quark Model:  $|q\bar{q}\rangle$  states




# Motivation

- Mesons in the Constituent Quark Model:  $|q\bar{q}\rangle$  states
- Many predicted  $|q\bar{q}\rangle$  states unobserved/need confirmation




# Motivation

- Mesons in the Constituent Quark Model:  $|q\bar{q}\rangle$  states
- Many predicted  $|q\bar{q}\rangle$  states unobserved/need confirmation



- QCD allows meson configurations beyond  $|q\bar{q}\rangle$  - so-called exotics





# Motivation

- Mesons in the Constituent Quark Model:  $|q\bar{q}\rangle$  states

- Many predicted  $|q\bar{q}\rangle$  states unobserved/need confirmation

- QCD allows meson configurations beyond  $|q\bar{q}\rangle$  - so-called exotics




- Meson spectroscopy:

⇒ Understand non-perturbative QCD

⇒ Input to test SM predictions with experimental data

# Experimental Setup


- Located at CERN SPS
- 190 GeV/c negative hadron beam
- **Non-strange** light meson spectroscopy:  
 $\pi^- p$  scattering



[COMPASS](#)  
[Nucl. Instrum. Methods. 779 \[2015\] 69](#)

# Non-Strange Light-Meson Spectroscopy at COMPASS

- Diffractive scattering of high-energy pion beam
- Excited non-strange meson resonance  $X^-$
- Decay to  $N$  hadron final state



# Non-Strange Light-Meson Spectroscopy at COMPASS

- Diffractive scattering of high-energy pion beam
- Excited non-strange meson resonance  $X^-$
- Decay to  $N$  hadron final state


Analyzed channels:

- $\pi^- \pi^- \pi^+ / \pi^- \pi^0 \pi^0$
- $\eta \pi^- / \eta' \pi^-$
- $\omega \pi^- \pi^0$




# Partial-Wave Analysis – Method

Data  
Diffractively produced events




# Partial-Wave Analysis – Method




# Partial Waves

- Excited meson  $X^-$  with quantum numbers  $0^-0^+$




# Partial Waves

- Excited meson  $X^-$  with quantum numbers  $0^-0^+$
- Isobar model:  $X^- \rightarrow \omega \rho(770)$ 
  - Unstable intermediate state/isobar  $\rho(770)$




# Partial Waves

- Excited meson  $X^-$  with quantum numbers  $0^-0^+$
- Isobar model:  $X^- \rightarrow \omega \rho(770)$ 
  - Unstable intermediate state/isobar  $\rho(770)$
  - $LS = P1$  coupling between  $\omega$  and  $\rho(770)$




# Partial Waves

- Excited meson  $X^-$  with quantum numbers  $0^-0^+$
- Isobar model:  $X^- \rightarrow \omega \rho(770)$ 
  - Unstable intermediate state/isobar  $\rho(770)$
  - $LS = P1$  coupling between  $\omega$  and  $\rho(770)$
- $\rho(770) \rightarrow \pi^- \pi^0$ 
  - Second  $ls = P1$  coupling
- $i = 0^-0^+ [\rho(770)P] \omega P1$



# Partial Waves

- Excited meson  $X^-$  with quantum numbers  $J^P M^E$
- Isobar model:  $X^- \rightarrow \omega \xi^-$ 
  - Unstable intermediate state/**isobar**  $\xi^-$
  - $LS$  coupling between  $\omega$  and  $\xi^-$
- $\xi^- \rightarrow \pi^- \pi^0$ 
  - Second  $ls$  coupling
- $i = J^P M^E [\xi l] \omega LS$




# Partial-Wave Decomposition

Model measured intensity

- by a coherent sum over different partial waves




$$i = J^P M^\epsilon [\xi l] \omega LS$$

- in narrow  $(m_X, t')$  cells
- without assumptions about resonance content of  $X^-$  in partial waves











# Partial-Wave Analysis – Method



# Partial-Wave Analysis – Method



# Partial-Wave Analysis – Method




# Spin-exotic $1^{-+}$ at COMPASS

- Non-Strange light mesons: certain  $J^{PC}$  not possible for  $|q\bar{q}\rangle$  - spin-exotic
- Lattice QCD: lightest hybrid meson is spin-exotic  $1^{-+}$  state

# Spin-exotic $1^{-+}$ at COMPASS

$$\pi^{-}\pi^{-}\pi^{+}$$




Resonance parameters:

$m_0 \text{ (MeV}/c^2)$	$\Gamma \text{ (MeV}/c^2)$
$1600^{+110}_{-60}$	$580^{+100}_{-230}$

[M. Aghasyan et al. \(COMPASS\)  
Phys. Rev. D 98, 092003 \[2018\]](#)

# Spin-exotic $1^{-+}$ at COMPASS

$$\pi^{-}\pi^{-}\pi^{+}$$



Resonance parameters:

$m_0$ (MeV/c <sup>2</sup> )	$\Gamma$ (MeV/c <sup>2</sup> )
$1600^{+110}_{-60}$	$580^{+100}_{-230}$

[M. Aghasyan et al. \(COMPASS\) Phys. Rev. D 98, 092003 \[2018\]](#)


$$\eta\pi^{-}$$

$$\pi_1(1400)?$$

$$\eta'\pi^{-}$$

$$\pi_1(1600)?$$

Coupled-channel analysis of COMPASS data by JPAC:  
One pole is sufficient to describe both partial waves




Pole parameters:

$m_0$ (MeV/c <sup>2</sup> )	$\Gamma$ (MeV/c <sup>2</sup> )
$1564 \pm 24 \pm 86$	$492 \pm 54 \pm 102$

[A. Rodas et al. \(JPAC\), Phys. Rev. D 122, 042002 \[2019\]](#)


Extended Analysis incl. Crystal Barrel:

[B. Kopf et al., Eur. Phys. J.C 81, 1056 \[2021\]](#)



# Spin-exotic $1^{-+}$ at COMPASS

$$\pi^{-}\pi^{-}\pi^{+}$$



Resonance parameters:


$m_0$ (MeV/c <sup>2</sup> )	$\Gamma$ (MeV/c <sup>2</sup> )
$1600^{+110}_{-60}$	$580^{+100}_{-230}$

[M. Aghasyan et al. \(COMPASS\) Phys. Rev. D 98, 092003 \[2018\]](#)

$$\eta\pi^{-}$$


$$\pi_1(1400)?$$

Coupled-channel analysis of COMPASS data by JPAC:  
One pole is sufficient to describe both partial waves



$$\eta'\pi^{-}$$

$$\pi_1(1600)?$$



Pole parameters:

$m_0$ (MeV/c <sup>2</sup> )	$\Gamma$ (MeV/c <sup>2</sup> )
$1564 \pm 24 \pm 86$	$492 \pm 54 \pm 102$


[A. Rodas et al. \(JPAC\), Phys. Rev. D 122, 042002 \[2019\]](#)

Extended Analysis incl. Crystal Barrel:

[B. Kopf et al., Eur. Phys. J.C 81, 1056 \[2021\]](#)

$$\omega(782)\pi^{-}\pi^{+}$$

- Resonance-like signal in  $1^{-+}$
- Resonance-model fit in progress




# Non-Strange Light-Meson Spectroscopy at COMPASS

Analyzed channels:

- $\pi^- \pi^- \pi^+ / \pi^- \pi^0 \pi^0$
- $\eta \pi^- / \eta' \pi^-$
- $\omega \pi^- \pi^0$

Spin-exotic  $1^{-+}$  signals at COMPASS




⇒ Non-strange light-meson spectrum well studied by COMPASS




# Strange-Meson Spectrum

- Many missing states predicted by the Constituent Quark Model
- Most experimental results published 30+ years ago
- No established exotic strange meson (except  $K_0^*(700)/\kappa$ )



# Experimental Setup

- Located at CERN SPS
- 190 GeV/c negative hadron beam
- **Strange-meson spectroscopy**  
 $K^- p$  scattering




[COMPASS](#)  
[Nucl. Instrum. Methods. 779 \[2015\] 69](#)

# Experimental Setup – Particle ID

- Located at CERN SPS
- 190 GeV/c negative hadron beam
- Strange-meson spectroscopy



$K^- p$  scattering



[COMPASS](#)  
[Nucl. Instrum. Methods. 779 \[2015\] 69](#)

# Strange-Meson Spectroscopy in $K^- \pi^- \pi^+$

- World's largest  $K^- \pi^- \pi^+$  data set
  - 720k diffractive events
- Well known states are dominant
  - ⇒ Less well-known states not visible in  $m_{K\pi\pi}$  spectrum
  - ⇒ Partial-wave analysis reveals rich spectrum




# Limited final state PID of $K^- \pi^- \pi^+$

- Significant backgrounds from  $\pi^- \pi^- \pi^+$ ,  $K^- K^- K^+$ , ...  
⇒ Partial-wave model includes background contributions
- Final state PID only works for  $p \lesssim 50 \text{ GeV}/c$   
⇒ Artifacts in certain partial waves for  $m_{K\pi\pi} \lesssim 1.5 \text{ GeV}/c^2$


# Resonance-Model Fit of $K^- \pi^- \pi^+$

- Model  $m_X$  dependence of partial waves as **sum of contributions**:




# Resonance-Model Fit of $K^- \pi^- \pi^+$

- Model  $m_X$  dependence of partial waves as **sum of contributions**:
  - Breit-Wigner amplitudes of  $K\pi\pi$  resonances



# Resonance-Model Fit of $K^- \pi^- \pi^+$


- Model  $m_X$  dependence of partial waves as **sum of contributions**:
  - Breit-Wigner amplitudes of  $K\pi\pi$  resonances
  - Coherent **non-resonant background**






# Resonance-Model Fit of $K^- \pi^- \pi^+$

- Model  $m_X$  dependence of partial waves as **sum of contributions**:
  - Breit-Wigner amplitudes of  $K\pi\pi$  resonances
  - Coherent **non-resonant background**
  - Incoherent  $\pi^- \pi^- \pi^+$  **background** modeled by COMPASS  $\pi^- \pi^- \pi^+$  analysis




# Resonance-Model Fit of $K^- \pi^- \pi^+$

- Model  $m_X$  dependence of partial waves as **sum of contributions**:
  - Breit-Wigner amplitudes of  $K\pi\pi$  resonances
  - Coherent **non-resonant background**
  - Incoherent  $\pi^- \pi^- \pi^+$  **background** modeled by COMPASS  $\pi^- \pi^- \pi^+$  analysis
  - Incoherent **effective background** for other background processes





# Strange Mesons with $J^P = 0^-$

- $K(1460)$  peak at about  $1.4 \text{ GeV}/c^2$ 
  - Established state
  - Increased background below  $1.5 \text{ GeV}/c^2$   
 $\Rightarrow$  Fixed Breit-Wigner resonance
- $K(1630)$  peak at about  $1.7 \text{ GeV}/c^2$ 
  - $8.3\sigma$  statistical significance
- $K(1830)$  peak at about  $2.0 \text{ GeV}/c^2$ 
  - $5.4\sigma$  statistical significance




# Strange Mesons with $J^P = 0^-$


- Quark model predicts 2 excited  $0^-$  states
- Indications for 3 states in a single analysis
  - ⇒ Supernumerary state  $K(1630)$
  - ⇒ Possible candidate for exotic strange meson



# Strange Mesons




# Strange Mesons



- Most comprehensive analysis of  $K^- \pi^- \pi^+$ 
  - 11 states extracted from COMPASS data


# Strange Mesons



- Most comprehensive analysis of  $K^- \pi^- \pi^+$ 
  - 11 states extracted from COMPASS data

⇒ Strange light-meson spectrum is studied by COMPASS

# Strange Mesons



- Most comprehensive analysis of  $K^- \pi^- \pi^+$ 
  - 11 states extracted from COMPASS data

⇒ Strange light-meson spectrum is studied by COMPASS  
But limited due to final state PID



# Conclusion & Outlook

- COMPASS contributes to the knowledge of the light meson spectrum by studying many different final states in great detail
  - Exotic  $\pi_1(1600)$  observed in multiple final states
- Most comprehensive analysis of the  $K^- \pi^- \pi^+$  final state
  - Measured resonance parameters of 11 states
  - Exotic strange-meson candidate: Supernumerary state in  $J^P = 0^-$
- Active analysis of other final states:  $f_1 \pi^-$ ,  $K_S K^-$ ,  $K_S K_S \pi^-$ ,  $K_S \pi^-$ ,  $\Lambda \bar{p}$

# Conclusion & Outlook

- COMPASS contributes to the knowledge of the light meson spectrum by studying many different final states in great detail
  - Exotic  $\pi_1(1600)$  observed in multiple final states
- Most comprehensive analysis of the  $K^- \pi^- \pi^+$  final state
  - Measured resonance parameters of 11 states
  - Exotic strange-meson candidate: Supernumerary state in  $J^P = 0^-$
- Active analysis of other final states:  $f_1 \pi^-$ ,  $K_S K^-$ ,  $K_S K_S \pi^-$ ,  $K_S \pi^-$ ,  $\Lambda \bar{p}$
- AMBER phase-II: Strange-meson spectroscopy
  - ⇒ Promising experiment to probe strange-meson spectrum with even higher precision

B. Seitz: Status of the AMBER Phase-2 Proposal Preparation (Tue, 14:30)

# Backup

# Limited final state PID of $K^- \pi^- \pi^+$

- Significant backgrounds from  $\pi^- \pi^- \pi^+$ ,  $K^- K^- K^+$ , ...  
⇒ Partial-wave model includes background contributions
- Limited acceptance  
⇒ Artifacts in certain partial waves for  $m_{K\pi\pi} \lesssim 1.5 \text{ GeV}$

