Transversity signals in two hadron correlation at COMPASS

Rainer Joosten
on behalf of the COMPASS collaboration

Helmholtz Institut für Strahlen- und Kernphysik
Nußallee 14-16, 53115 Bonn, Germany
E-mail: Rainer.Joosten@cern.ch

Abstract.
Measurement of two hadron production introducing the chiral odd interference fragmentation function $H_1^{\sphericalangle}$ is considered a new probe of the transverse spin distribution $\Delta_T q(x)$. COMPASS is a fixed target experiment on the SPS M2 beamline at CERN. Its target can be polarised both longitudinally and transversely with respect to the polarised 160 GeV/c μ^+ beam. In 2002, 2003, and 2004, 20% of the beam-time was spent in the transverse configuration on a ^{6}LiD target, allowing the measurement of transversity effects. First results of the analysis of two hadron production will be reported.

Keywords: Transverse Spin Physics, Interference Fragmentation Functions, Transversity
PACS: 14.20.Dh, 13.60.Hb

The cross-section for deep inelastic scattering off spin 1/2 hadrons can be parametrised, in leading order, in terms of three quark distribution functions: the spin averaged distribution $q(x)$, the helicity distribution $\Delta q(x)$, and the transverse spin distribution $\Delta_T q(x)$. This last distribution function, referred to as transversity, is chiral-odd and can only be measured in combination with another chiral-odd function. So far, attempts were made to measure $\Delta_T q(x)$ in combination with the Collins fragmentation-function $\Delta D_a^h(z, p_T^h)$, requiring the partial detection of the hadronic products (semi-inclusive measurement) [1][2]. Another promising probe suggested is the measurement of two hadron production introducing the chiral odd interference fragmentation function $H_1^{\sphericalangle}(z, M_h^2)$. The properties of interference fragmentation functions are described in Refs. [3][4][5][6][7][8]. At leading twist, the fragmentation function of a transversely polarised quark q into a pair h of two hadrons h_1 and h_2 can be written as:

$$D_q^h(z, M_h^2) + H_1^{\sphericalangle}(z, M_h^2) sin(\phi_{RS}) \tag{1}$$

with $\phi_{RS} = \phi_R - \phi_{S'} = \phi_R + \phi_S - \pi$, where $\phi_{S'}$ is the azimuthal angle of the struck quark spin, ϕ_S is the azimuthal angle of the initial quark spin and $\phi_{S'} = \pi - \phi_S$. Here, ϕ_R is the angle between the scattering plane of the incoming $\mathbf{l}$ and outgoing $\mathbf{l'}$ lepton and the plane spanned by the virtual photon momentum $\mathbf{q}$ and the component $\mathbf{R}_T$ of the relative hadron momentum $\mathbf{R} = \frac{1}{2}(\mathbf{P}_1 - \mathbf{P}_2)$ which is perpendicular to the summed hadron momentum $\mathbf{P}_h = \mathbf{P}_1 + \mathbf{P}_2$. Following the *Trento conventions* the angles are defined [9] to be:

$$\cos\phi_R = \frac{(\hat{\mathbf{q}} \times \mathbf{l})}{|\hat{\mathbf{q}} \times \mathbf{l}|} \cdot \frac{(\hat{\mathbf{q}} \times \mathbf{R_T})}{|\hat{\mathbf{q}} \times \mathbf{R_T}|}, \qquad \sin\phi_R = \frac{(\mathbf{l} \times \mathbf{R_T}) \cdot \hat{\mathbf{q}}}{|\hat{\mathbf{q}} \times \mathbf{l}|\,|\hat{\mathbf{q}} \times \mathbf{R_T}|}, \tag{2}$$

While, $z = z_1 + z_2 = (E(h_1) + E(h_2))/(E_l - E_{l'})$ is the fraction of the transfered energy carried by the two hadrons (E_l being the incoming and $E_{l'}$ the scattered lepton energy), and M_h^2 is their invariant mass squared.

An asymmetry is expected in the azimuthal angle of the hadron plane depending on ϕ_{RS}. This asymmetry, which is connected to the $\Delta_T q(x)$ as well as $H_1^{\sphericalangle}(z, M_h^2)$, has not been measured so far on a transversely polarised deuteron target. The measured raw asymmetry $A_{UT}^{sin\phi_{RS}}$ is connected to the physical asymmetry $A_{\phi_{RS}}$ by

$$\frac{A_{UT}^{sin\phi_{RS}}}{D_{NN} f P} = A_{\phi_{RS}} = \frac{\Sigma_i e_i \Delta_T q_i(x) H_1^{\sphericalangle h}(z, M_h^2)}{\Sigma_i e_i q_i(x) D_i^h(z, M_h^2)} \tag{3}$$

where $f (\approx 0.4)$ is the dilution factor, $P (\approx 0.45)$ the target polarisation and $D_{NN} = (1-y)/(1-y+y^2/2)$ the depolarisation factor. Here, $y = (E_l - E_{l'})/E_l$ is the fraction of the incoming lepton energy transfered to the hadronic system. While one model predicts a strong dependence of the fragmentation function $H_1^{\sphericalangle}(z, M_h^2)$ on the invariant mass of the two-hadron system in the region of the ρ-mass (Ref. [5]) due to an interference term in two pion production, another model [6] does not see this dependence.

The COMPASS experiment [10][11] uses the high intensity 160 GeV secondary μ^+-beam in the CERN SPS M2 beamline. This beam is naturally longitudinally polarised with a polarisation of ≈-76%. The polarised target consists of two target cells, each 60 cm long, filled with ^{6}LiD, which are individually polarised using separate RF-cavities to allow simultaneous data taking on target cells of opposite polarisation. The target can be polarised longitudinally or transversely with respect to the beam axis.

The data discussed here was taken in 2002 and 2003 with a transversely polarised target. The sample consists of three independent data taking periods, where each period was split in two subperiods with opposite spin orientation in the individual target cells.

The event selection is analogous to the analysis of the Collins and Sivers asymmetries [1][2]. The primary vertex, with identified incoming and scattered muon, is required to be in either of the two target cells. At least two hadrons are required to origin from the same vertex. The separation of muons and hadrons is primarily done by cutting on the amount of traversed material in the spectrometer and on the energy loss in the two hadronic calorimeters. Moreover, the kinematic cuts $Q^2 > 1\,(GeV/c)^2$, $W > 5\,GeV/c^2$ and $0.1 < y < 0.9$ were applied to ensure a deep-inelastic scattering sample above nuclear resonances and within the COMPASS trigger acceptance. The final data sample had average values for $x = 0.035$, $y = 0.33$, and $Q^2 = 2.4\,(GeV/c)^2$. The mean hadron multiplicity of the events selected by these kinematic cuts is 1.9 hadrons/event.

Based on this sample, hadron pairs are selected by choosing all combinations of positive (h_1) and negative (h_2) hadrons fulfilling the requirements $z_1 > 0.1$ and $z_2 > 0.1$ to reject the target fragmentation region as well as $z = z_1 + z_2 < 0.9$ to suppress the contamination with exclusively produced ρ-mesons. The resulting sample contains $2.8 \cdot 10^6$ hadron combinations. From the data, for each target cell and polarisation, the property

$$N(\phi_{RS}) = N_0 \cdot (1 + A_{UT}^{sin\phi_{RS}} \cdot sin\phi_{RS}) \cdot F_{acc}(\phi_{RS}) \tag{4}$$

can be derived, where $F_{acc}(\phi_{RS})$ is the (unknown) angle dependant acceptance function of the detector. However, by comparing the subperiods with opposite target spin,

this acceptance function cancels, resulting in

$$A_{UT}^{sin\phi_{RS}} \cdot sin\phi_{RS} = \frac{N^{\Uparrow}(\phi_{RS}) - RN^{\Downarrow}(\phi_{RS}+\pi)}{N^{\Uparrow}(\phi_{RS}) + RN^{\Downarrow}(\phi_{RS}+\pi)} \quad (5)$$

where $R = N_{tot}^{\Uparrow}/N_{tot}^{\Downarrow}$ is the ratio of the events with opposite target polarisation. (For details please refer to [2]). This procedure is repeated for both target cells and all periods. Finally, the weighted mean of the results is calculated.

Figure 1 shows the preliminary results from the COMPASS 2002 and 2003 data. The plots show (from top to bottom) the asymmetries vs. the invariant mass M_h of the hadron pair, Bjorken-x and z respectively. The observed asymmetries are very small and no significant signal can be observed. Especially the asymmetry vs. M_h does not at all show a strong dependence on the hadron invariant mass. Within statistical errors, all measured aysmmetries are compatible with zero.

Including the data of the COMPASS 2004 run on a ^{6}LiD target will double the total statistics. Using hadron identification based on the RICH information will clean the hadron sample. Some investigations are ongoing, concerning the selection of hadrons. Preliminary results on ordering by p_t have already been produced. According to [3] [4], an ordering based on z might enhance the signal. Additionally, a complementary measurement on a proton target is planned for the 2006 run.



Figure 1: Asymmetries $A_{\phi_{RS}}$ for the 2002 and 2003 data vs. invariant mass of the hadron pair (top), Bjorken- x (center) and $z = z_1 + z_2$ (bottom).

REFERENCES

1. A. Bressan (COMPASS), these proceedings.
2. V.Yu. Alexakhin *et al.* [COMPASS Collaboration], *Phys. Rev. Lett.* **94**, 202002 (2005).
3. J.R. Collins, S.F. Heppelmann and G.A. Ladinsky,*Nucl. Phys.* **B420**, 565 (1994).
4. X. Artru and J.C. Collins, *Z. Phys.* **C69**, 277 (1996).
5. R.L. Jaffe, X. Jin and J. Tang, *Phys. Rev. Lett.* **80**, 1166 (1998).
6. M. Radici, R. Jakob and A. Bianconi, *Phys. Rev.* **D65** 074031 (2002).
7. A. Bianconi *et al.*, *Phys. Rev.* **D62**, 034008 (2000).
8. A. Bacchetta, M. Radici, *Phys. Rev.* **D69**, 074026 (2004).
9. A. Bacchetta, M. Radici, Proceeding of DIS 2004, hep-ph/0407345 (2004).
10. G. Baum *et al.* [COMPASS Collaboration], CERN-SPSLC-96-14.
11. G.K. Mallot, Nucl. Instrum. Meth. A **518**, 121 (2004).