

Studies of TMDs at COMPASS

H. Wollny on behalf of the COMPASS Collaboration

*Physikalisches Institut, Albert-Ludwigs University of Freiburg,
79104 Freiburg, Germany
Heiner.Wollny@cern.ch

Transverse spin and transverse momentum distribution functions of the constituents of the nucleon are a crucial input for a complete description of the nucleon. COMPASS measured such for longitudinally and transversely polarized deuterons and protons. In the following we will focus on recent results from the 2007 transverse proton data and on the results for unpolarized deuterons.

Keywords: SIDIS, polarized target, Transversity, TMD, Collins, Sivers

1. Introduction

At leading order and integrating over transverse quark momenta, three parton distribution functions (PDFs) are needed for a complete description of the nucleon. The quark density $q(x)$, the helicity distribution $\Delta q(x)$ and the transversity distribution $\Delta_T q(x)$. The first PDF, $q(x)$, describes the probability to scatter off a quark inside the nucleon carrying the momentum fraction x . The helicity PDF, $\Delta q(x)$, is defined for a longitudinal polarized proton as the difference of the probabilities that the struck quark carries momentum fraction x and its spin is parallel or anti-parallel to the spin of the parent nucleon. The transversity PDF, $\Delta_T q(x)$, is analogously defined to $\Delta q(x)$, however for a transversely polarized nucleon. Because $\Delta_T q(x)$ is chiral-odd it cannot be measured in inclusive DIS. However it can be accessed in semi-inclusive deep-inelastic scattering in combination with a chiral-odd fragmentation function. Three different channels to address the transversity distribution, have been analysed, both for deuteron and proton targets. They will be discussed in Sec. 3.

If one considers intrinsic transverse momenta of the quarks inside the nucleon, several further PDFs so called transverse momentum dependent distribution functions (TMDs) contribute to the SIDIS cross-section, which will be discussed in Sec. 4.

2. The COMPASS Experiment

COMPASS is a fixed target experiment situated at the M2 beam line of the SPS accelerator at CERN. It has a wide physics program dedicated to the study of the nucleon spin structure and of hadron spectroscopy. In the years 2002, 2003, 2004 and 2006 data with a polarized ${}^6\text{LiD}$ target were taken. In the years 2002-2004 approximately 20% of the data taking was dedicated to transverse target polarization. In the year 2007 COMPASS took data with a NH_3 target, equally shared between longitudinal and transverse target polarization. After the years 2008 and 2009, dedicated to baryon and meson spectroscopy measurements with hadron beams, in 2010 COMPASS continues with muon beam and transversely polarized protons.

The target consists of several cells along the beam direction, which are oppositely polarized. The polarization is periodically reversed to reduce systematic effects due to the different acceptance of the cells. The detector consists of two open field dipole magnets, allowing the detection of particles with ± 180 mrad. Particle identification is done with a Ring Imaging Cerenkov detector, two hadron calorimeters and muon filters. For a detailed description we refer to reference [1].

3. Transversity

Collins Asymmetry: The chiral-odd transversity distribution can be measured in SIDIS involving the chiral-odd Collins fragmentation function $\Delta_T^0 D_q^h$ [2], which describes the fragmentation of transversely polarized quarks into unpolarized hadrons. According to Collins the fragmentation leads to an asymmetry A_{Coll} of the number of produced hadrons in $\sin(\phi_h + \phi_S - \pi)$. Here ϕ_h is the azimuthal angle, measured around the direction of the virtual photon, of the hadron with respect to the scattering plane and ϕ_S is the azimuthal angle between the spin of the initial quark and the scattering plane [3-5]. In first order the measured asymmetry A_{Coll} is proportional to a convolution over intrinsic quark transverse momenta of the transversity distribution and the Collins fragmentation function measured at Belle [6]. Because of the convolution, assumptions about the transverse momentum dependence of the distribution and the fragmentation functions have to be made to extract transversity [7]. In addition the Q^2 evolution of the Collins fragmentation function from Belle to COMPASS energies is not known, introducing a further uncertainty.

Dihadron Interference: Another possibility to access the transversity distribution is to measure it in SIDIS in combination with the polarized dihadron interference fragmentation function (DiFF) [8–11]. Here, the fragmentation of a transversely polarized quark into two unpolarized hadrons leads to an azimuthal modulation in $\sin(\phi_R + \phi_S - \pi)$ in the number of produced hadron pairs. Where ϕ_S is defined as above and ϕ_R is the azimuthal angle, measured around the direction of the virtual photon, between the vector $\vec{R} = (z_2\vec{P}_1 - z_1\vec{P}_2)/(z_1 + z_2)$ and the scattering plane. Where P_1 and P_2 are the momenta of the two hadrons and z_1 and z_2 are their fractional energies. For oppositely charged hadron pairs P_1 is per definition the hadron with positive charge, otherwise P_1 is the hadron which has the largest energy fraction. Here, the measured asymmetry A_{RS} is proportional to a product of the transversity distribution and the polarized DiFF, recently measured at Belle [12]. Hence, no assumptions about transverse momentum dependences of distribution and fragmentation functions are needed, to extract transversity. In addition, the Q^2 evolution of the polarized DiFF from Belle energies to COMPASS energies is known [13]. Hence, the extraction of transversity with this channel is from theoretical point of view cleaner than the extraction from the Collins asymmetries.

Lambda Polarization: A third channel to measure transversity is to study Λ and $\bar{\Lambda}$ polarization in SIDIS with a transversely polarized target [14]. The polarization P_T^Λ of the produced $\Lambda/\bar{\Lambda}$ -Hyperons, measured via the parity violating decay into $\Lambda \rightarrow p\pi^-$ and $\bar{\Lambda} \rightarrow \bar{p}\pi^+$, respectively, is used as a polarimeter of the initial transverse spin of the fragmenting quark. It is in first order proportional to a product of transversity and the fragmentation function $\Delta_T D_q^\Lambda$, which describes the fragmentation of a transversely polarized quark into a transversely polarized Λ -Hyperon.

4. Transverse Momentum Dependent Distribution Functions, TMDs

Taking into account intrinsic transverse momenta of the quarks in total eight TMDs are needed for a complete description at leading-twist [15]. For unpolarized nucleons the Cahn effect, the Boer-Mulders TMD and perturbative QCD effects, like gluon radiation, contribute to $\cos\phi_h$ and $\cos 2\phi_h$ modulations in the SIDIS cross section of single hadrons [16–18]. For a transversely polarized nucleon the Sivers TMD $\Delta_0^T q$ [19] is related to the quark angular orbital momentum inside a transversely polarized nucleon and is therefore of special interest since this could be the crucial piece to

solve the nucleon spin puzzle. The Sivers effect leads to an azimuthal modulation of the number of produced hadrons in $\sin(\phi_h - \phi_S)$, where ϕ_h and ϕ_S are defined as for the Collins asymmetry discussed in Sec. 3. The asymmetry A_{Siv} is proportional to a convolution over intrinsic quark transverse momenta of the Sivers function and the well-known unpolarized fragmentation function D_q^h .

Recent results of the Sivers function measured with protons and results of $\cos\phi_h$ and $\cos 2\phi_h$ modulations for unpolarized deuterons will be reviewed in this article. For the six remaining azimuthal asymmetries measured with transversely polarized deuterons it is referred to [20]. Detailed results of azimuthal asymmetries measured with longitudinally polarized deuterons can be found in [21].

5. Event selection

Kinematic cuts on the squared four momentum transfer $Q^2 > 1 (\text{GeV}/c)^2$, on the fractional energy transfer of the muon $0.1 < y < 0.9$ and the hadronic invariant mass $W > 5 \text{ GeV}/c^2$ are applied to select DIS events. The selection of charged hadrons differs slightly for the various analyses. However, for the single hadron analyses in general a fractional energy $z > 0.2$ of the observed final hadron is required to select hadrons in the current fragmentation region. In addition a transverse momentum $p_T^h > 0.1 \text{ GeV}/c$ of the hadron with respect to the virtual photon is required to ensure a good definition of ϕ_h . Muons are rejected in demanding a minimal energy deposit in the hadronic calorimeters.

For the hadron pair analysis $z_{1,2} > 0.1$ is required for each hadron and for the sum $z_1 + z_2 < 0.9$ to reject exclusively produced ρ^0 -mesons. In order to have a good definition of ϕ_R a cut on $R_T > 70 \text{ MeV}/c$, the transverse component of \vec{R} with respect to the virtual photon direction, is applied.

6. Results

6.1. Transversity

The Collins asymmetries A_{Coll}^p have been evaluated for charged hadrons in bins of Bjorken x , energy fraction z and transverse momentum p_T^h . The results for the full 2007 proton statistics [22] are shown in Fig. 1. For positive hadrons the asymmetry is negative and for negative hadrons it is positive. For both charges the size of the asymmetry increases with x and is compatible in their strengths. The values agree both in magnitude and in sign with the previous measurements of HERMES [23], which have been performed

at a considerably lower electron beam momentum of $27.5 \text{ GeV}/c$. From the transversely polarized deuteron data Collins asymmetries for unidentified charged hadrons, charged pions and charged kaons have been extracted [3–5]. All asymmetries found to be small and compatible with zero within the statistical errors. Hence for deuteron the transversity distribution must be small or even vanishing due to isospin symmetry, because the proton results and the Belle results [6], confirmed the existence of a non-zero Collins fragmentation function.

The results for the transverse 2007 proton data of the Dihadron asymme-




Fig. 1. Collins asymmetries for positive and negative hadrons as a function of x , z and p_T . The horizontal bands indicate the systematical errors.

tries A_{RS}^p for oppositely charged hadron pairs [24] are shown as a function of x , z and M_{inv} in Fig. 2. A strong asymmetry is observed in the valence x -region. In the invariant mass one observes a strong signal around the ρ^0 -mass at around $0.77 \text{ GeV}/c^2$ and the asymmetry is negative over the whole mass range. Compared to the Collins asymmetries, the size of A_{RS}^p is approximately a factor of two larger, emphasizing the good analyzing power of the dihadron interference fragmentation function to measure transversity. From the transversely polarized deuteron data COMPASS extracted dihadron asymmetries for unidentified h^+h^- -pairs and identified $\pi^+\pi^-$, K^+K^- , π^+K^- and $K^+\pi^-$ -pairs [25–27]. In addition also different charge combinations have been analysed, considering the two most energetic hadrons in each event. All asymmetries found to be small and compatible with zero within the statistical errors.

The polarization of Λ and $\bar{\Lambda}$ -Hyperons measured with transversely polarized ${}^6\text{LiD}$ and NH_3 targets, respectively, have been evaluated in bins of x and z [28,29]. For both targets the evaluated polarizations of Λ and $\bar{\Lambda}$ -Hyperons are small and compatible with zero within the errors.




Fig. 2. Dihadron asymmetries A_{RS}^p for oppositely charged hadron pairs as a function of x , z and M_{inv} . The horizontal band indicates the systematical error.

6.2. *Sivers-Function*

The Sivers asymmetries for charged hadrons from the transverse 2007 proton data [22] are shown, as a function of x , z and p_T^h , in Fig. 3. Beside the shown systematical errors, for positive hadrons an additional systematical error of ± 0.01 is assigned, reflecting a 0.02 difference in the mean values of the asymmetries extracted in the first and the second half of data taking used for this analysis. The asymmetry for positive hadrons is significantly positive over almost the complete x -range. The asymmetry for negative hadrons is small and compatible with zero within the statistical errors. Both results show the same trend as the previous measurement of HERMES [30]. However, the magnitude of the asymmetry for positive hadrons is smaller than the one of HERMES. This might be explained by a possible dependence on W , the invariant mass of the photon-nucleon system. Studies indicate, that the asymmetry seem to be only present at small W , where HERMES measures and go to zero with increasing W . The Sivers asymmetries for unidentified charged hadrons, charged pions and kaons extracted from transversely polarized ${}^6\text{LiD}$ [3–5] were all found to be small and compatible with zero within the statistical errors.

6.3. *Azimuthal Asymmetries for Unpolarized Deuterons*

An unpolarized deuteron sample is obtained by combining data samples with opposite target polarization taken in the year 2004. The asymmetries of $\cos \phi_h$ and $\cos 2\phi_h$ modulations for charged hadrons as a function of x , z and p_T are shown in Fig. 4. Large asymmetries up to 20% are found for the $\cos \phi_h$ modulation and asymmetries in the order of 5% for the $\cos 2\phi_h$ modulation. The trend of the asymmetries for positive and negative hadrons




Fig. 3. Sivers asymmetries for positive and negative hadrons as a function of x , z and p_T . The horizontal bands indicate the systematical errors.

are similar, however, the magnitudes of the two differs significantly.




Fig. 4. Unpolarized asymmetries for charged hadrons as a function of x , z and p_T . The kinematical factors are: $\epsilon_1 = (2-y)\sqrt{1-y}/(1+(1-y)^2)$ and $\epsilon_2 = (1-y)/(1+(1-y)^2)$.

7. Summary and Outlook

Recent results of COMPASS measurements related to transversity and transverse momentum dependent distribution functions have been presented. In the year 2010 and 2011 COMPASS will continue its measurements with transversely and longitudinally polarized protons, which will significantly reduce the statistical errors.

References

1. P. Abbon *et al.*, *Nucl. Instrum. Meth.* **A577**, 455 (2007).
2. J. C. Collins, *Nucl. Phys.* **B396**, 161 (1993).
3. V. Y. Alexakhin *et al.*, *Phys. Rev. Lett.* **94**, p. 202002 (2005).
4. E. S. Ageev *et al.*, *Nucl. Phys.* **B765**, 31 (2007).
5. M. Alekseev *et al.*, *Phys. Lett.* **B673**, 127 (2009).
6. R. Seidl *et al.*, *Phys. Rev.* **D78**, p. 032011 (2008).
7. M. Anselmino *et al.*, *Nucl. Phys. Proc. Suppl.* **191**, 98 (2009).
8. A. V. Efremov *et al.*, *Phys. Lett.* **B284**, 394 (1992).
9. X. Artru and J. C. Collins, *Z. Phys.* **C69**, 277 (1996).
10. M. Radici, R. Jakob and A. Bianconi, *Phys. Rev.* **D65**, p. 074031 (2002).
11. A. Bacchetta and M. Radici, *Phys. Rev.* **D69**, p. 074026 (2004).
12. A. Vossen *et al.*, First Measurement of the Interference Fragmentation Function in e^+e^- at Belle, in *Proc. of XIII Advanced Research Workshop on High Energy Spin Physics, Dubna, Russia*, Sept. 2009.
13. F. A. Ceccopieri, M. Radici and A. Bacchetta, *Phys. Lett.* **B650**, 81 (2007).
14. X. Artru and M. Mekhfi, *Nucl. Phys.* **A532**, 351 (1991).
15. P. J. Mulders and R. D. Tangerman, *Nucl. Phys.* **B461**, 197 (1996).
16. R. N. Cahn, *Phys. Lett.* **B78**, p. 269 (1978).
17. D. Boer and P. J. Mulders, *Phys. Rev.* **D57**, 5780 (1998).
18. H. Georgi and H. D. Politzer, *Phys. Rev. Lett.* **40**, p. 3 (1978).
19. D. W. Sivers, *Phys. Rev.* **D41**, p. 83 (1990).
20. A. Kotzinian, Beyond Collins and Sivers: further..., in *Proc. of 15th Int. Workshop on DIS and Related Subjects, Munich, Germany*, April 2007.
21. I. A. Savin, Azimuthal asymmetries in production of charged hadrons by high energy muons on polarized deuterium targets, in *Proc. of XIII Advanced Research Workshop on High Energy Spin Physics, Dubna, Russia*, Sept. 2009.
22. H. Fischer, Review of Spin Physics in DIS, a personal selection, in *Proc. of 18th Int. Workshop on DIS and Related Subjects, Florence, Italy*, April 2010.
23. A. Airapetian *et al.*, *arXiv:1006.4221* (2010).
24. H. Wollny, Transversity Signal in two Hadron Pair Production in COMPASS, in *Proc. of 17th Int. Workshop on DIS and Related Topics, Madrid, Spain*, April 2009.
25. R. Joosten, *AIP Conf. Proc.* **792**, 957 (2005).
26. R. Joosten, *AIP Conf. Proc.* **915**, 646 (2007).
27. A. Vossen, Measurement of Transverse Spin Effects at COMPASS, in *Proc. of 42nd Rencontres de Moriond on QCD and Hadronic Interactions, La Thuile, Italy*, March 2007.
28. A. Ferrero, *AIP Conf. Proc.* **915**, 436 (2007).
29. D. Kang, Longitudinal and transverse Lambda polarization at COMPASS, in *Proc. of 18th Int. Workshop on DIS and Related Subjects, Florence, Italy*, April 2010.
30. A. Airapetian *et al.*, *Phys. Rev. Lett.* **103**, p. 152002 (2009).