

Review of COMPASS results

Barbara Badelek, University of Warsaw
– On behalf of COMPASS –

Low x meeting

Ischia Island, September 9 – 13, 2009

Outline

- 1 Collaboration and programmes
- 2 History and plans
- 3 Detector
- 4 Acceptance
- 5 Parton distribution functions and observables
- 6 Inclusive measurements
- 7 Flavour separation of helicity distributions
- 8 Measurements on the transversely polarised target
- 9 Nucleon spin structure
- 10 COMPASS “near” plans (2010 – 2012): muon run
- 11 COMPASS “medium” plans (\gtrsim 2012): DVCS measurements
- 12 Outlook

Outline

- 1 Collaboration and programmes
- 2 History and plans
- 3 Detector
- 4 Acceptance
- 5 Parton distribution functions and observables
- 6 Inclusive measurements
- 7 Flavour separation of helicity distributions
- 8 Measurements on the transversely polarised target
- 9 Nucleon spin structure
- 10 COMPASS “near” plans (2010 – 2012): muon run
- 11 COMPASS “medium” plans (\gtrsim 2012): DVCS measurements
- 12 Outlook

CO_{mmon} MUon and PROton APParatus for STructure and SPECTroscopy



NA58, at the CERN SPS
 ~ 250 physicists
 ~ 30 institutes

Muon programme	Hadron programme
Spin dependent structure function g_1 Gluon polarisation in the nucleon Quark polarisation distributions Transversity Vector meson production Λ polarisation	Primakoff effect, π and K polarisabilities Exotic states, glueballs (Double) charmed mesons Multiquark states
Future: DVCS	Future: Drell-Yan production

Outline

- 1 Collaboration and programmes
- 2 History and plans**
- 3 Detector
- 4 Acceptance
- 5 Parton distribution functions and observables
- 6 Inclusive measurements
- 7 Flavour separation of helicity distributions
- 8 Measurements on the transversely polarised target
- 9 Nucleon spin structure
- 10 COMPASS “near” plans (2010 – 2012): muon run
- 11 COMPASS “medium” plans (\gtrsim 2012): DVCS measurements
- 12 Outlook

History: 2002 – 2009 and beyond

- 2002 160 GeV polarised μ beam & ${}^6\text{LiD}$ long./transv. polaris.
- 2003 *idem*
- 2004 *idem*
- 2004 pilot hadron run
- 2005 no SPS beam (several upgrades: target, RICH)
- 2006 160 GeV polarised μ beam & ${}^6\text{LiD}$ long. polarisation
- 2007 160 GeV polarised μ beam & NH_3 transv./long. polaris.
- 2008 190 GeV pion beam; diffractive and central production
- 2009 190 GeV pion, kaon, proton beams
- \gtrsim 2010 Proposal Addendum: 1+1 year muon run
with transv. and longit. polarised proton target
- \gtrsim 2012 Letter-of-Intent: DVCS and Drell–Yan

Outline


- 1 Collaboration and programmes
- 2 History and plans
- 3 **Detector**
- 4 Acceptance
- 5 Parton distribution functions and observables
- 6 Inclusive measurements
- 7 Flavour separation of helicity distributions
- 8 Measurements on the transversely polarised target
- 9 Nucleon spin structure
- 10 COMPASS “near” plans (2010 – 2012): muon run
- 11 COMPASS “medium” plans (\gtrsim 2012): DVCS measurements
- 12 Outlook

COMPASS Spectrometer



(muon run)

Nucl. Instr. Meth. A577 (2007) 455

Two stages, ~ 350 planes
 Calorimetry
 Particle identification
 Large, solid state polarised target




COMPASS polarised targets




- * Two (three in 2006, 2007) target cells, oppositely polarised
- * Polarisation reversed every 8 h (less frequent in 2006, 2007)
- * Material: solid ^6LiD (NH_3 in 2007)
- * Polarisation: $\sim 50\%$ ($\sim 90\%$ in 2007), by the Dynamical Nuclear Polarisation
- * Dilution: $f \sim 0.4$ (~ 0.15 in 2007)
- * Polar acceptance: ~ 70 mrad (~ 180 mrad in 2006, 2007)

COMPASS RICH



Before upgrade: white distribution
After upgrade: yellow distribution



Outline

- 1 Collaboration and programmes
- 2 History and plans
- 3 Detector
- 4 Acceptance**
- 5 Parton distribution functions and observables
- 6 Inclusive measurements
- 7 Flavour separation of helicity distributions
- 8 Measurements on the transversely polarised target
- 9 Nucleon spin structure
- 10 COMPASS “near” plans (2010 – 2012): muon run
- 11 COMPASS “medium” plans (\gtrsim 2012): DVCS measurements
- 12 Outlook

Acceptance of electroproduction experiments




Figure from: N. D'Hose, Villars 2004


COMPASS

Outline


- 1 Collaboration and programmes
- 2 History and plans
- 3 Detector
- 4 Acceptance
- 5 Parton distribution functions and observables**
- 6 Inclusive measurements
- 7 Flavour separation of helicity distributions
- 8 Measurements on the transversely polarised target
- 9 Nucleon spin structure
- 10 COMPASS “near” plans (2010 – 2012): muon run
- 11 COMPASS “medium” plans (\gtrsim 2012): DVCS measurements
- 12 Outlook

Partonic structure of the nucleon; distribution functions


Three species of **twist-two** quark distributions in QCD (after integrating over the quark intrinsic k_t):



Quark momentum DF;
well known (unpolarised DIS → $F_{1,2}(x)$).



Difference in DF of quarks with spin parallel or antiparallel to the nucleon's spin;
known (polarised DIS → $g_1(x)$).



Difference in DF of quarks with spin parallel or antiparallel to the nucleon's spin in the transversely polarised nucleon;
unknown (polarised DIS → Collins asymmetry).

In the nonrelativistic approach $\Delta_T q(x)$ identical with $\Delta q(x)$.

$\Delta_T q(x)$ are C-odd and chiral-odd; may only be measured with another chiral-odd partner, e.g. fragmentation function.

If the k_t taken into account ⇒ 8 TMD appear; one, f_{1T}^\perp accessible through "Sivers asymmetry".

Observables in the polarised μ -p(d) scattering

- A direct observable, μ -p(d) cross section asymmetry $A_{\text{meas}}(x, Q^2)$, inclusive asymmetry $A_1(x, Q^2)$ and longitudinal spin-dependent structure function, $g_1(x, Q^2)$, are related as:

$$A_{\text{meas}} = \frac{1}{fP_T P_B} \left(\frac{N^{\leftarrow} - N^{\rightarrow}}{N^{\leftarrow} + N^{\rightarrow}} \right) = D(A_1 + \eta A_2) \approx D \frac{\sum_q e_q^2 \Delta q(x, Q^2)}{\sum_q e_q^2 q(x, Q^2)} = D \frac{g_1^d(x, Q^2)}{F_1^d(x, Q^2)}$$

(in the COMPASS kinematics η is small, $|\eta A_2| \ll |A_1|$)

$$f \sim 0.4, \quad P_T \sim 0.5, \quad P_B \sim -0.8, \quad \Delta q = q^+ - q^-, \quad q = q^+ + q^-$$

Important: $g_1^d = g_1^N (1 - \frac{3}{2} \omega_D) = \frac{g_1^p + g_1^n}{2} (1 - \frac{3}{2} \omega_D); \quad \omega_D = 0.05 \pm 0.01$

- Similarly, the semi-inclusive asymmetry A_1^h :



$$A_1^h(x, z, Q^2) \approx D \frac{\sum_q e_q^2 \Delta q(x, Q^2) D_q^h(z, Q^2)}{\sum_q e_q^2 q(x, Q^2) D_q^h(z, Q^2)}$$

Outline

- 1 Collaboration and programmes
- 2 History and plans
- 3 Detector
- 4 Acceptance
- 5 Parton distribution functions and observables
- 6 Inclusive measurements**
- 7 Flavour separation of helicity distributions
- 8 Measurements on the transversely polarised target
- 9 Nucleon spin structure
- 10 COMPASS “near” plans (2010 – 2012): muon run
- 11 COMPASS “medium” plans (\gtrsim 2012): DVCS measurements
- 12 Outlook

$g_1^d(x)$ in the nonperturbative ($Q^2 < 1 \text{ (GeV/c)}^2$) region



V.Yu. Alexakhin (COMPASS) *et al.* Phys. Lett. B **647** (2007) 330



- Order of magnitude improvement over the statistical precision of the SMC.
- Interplay between perturbative and nonperturbative mechanisms.
- Spin effects in g_1^d at low x and Q^2 absent ?

$g_1(x)$ for proton and deuteron, $Q^2 > 1$ (GeV/c) 2



NEW: proton data 2007; full deuteron statistics



Good agreement
between the experiments

$g_1(x)$ for proton and deuteron, $Q^2 > 1$ (GeV/c^2)...cont'd


NEW: proton data 2007; full deuteron statistics



COMPASS measurements at high Q^2 important for the QCD analysis!

$g_1(x)$ for proton and deuteron, $Q^2 > 1$ (GeV/c) 2 ...cont'd

- From the new proton data: $g_1^{NS} = g_1^p - g_1^n$; its first moment, $\Gamma_1^{NS}(Q^2) = \frac{1}{6} \left| \frac{g_A}{g_V} \right| C^{NS}(Q^2)$ (fundamental Bjorken sum rule)
- From QCD NLO fit to g_1^{NS} (COMPASS data only): $\frac{g_A}{g_V} = 1.30 \pm 0.07 \pm 0.10$ ($g_A/g_V = \Delta u - \Delta d = 1.260 \pm 0.003$ from the β decay of the neutron).
Test and confirmation of the Bjorken sum rule.



Outline



- 1 Collaboration and programmes
- 2 History and plans
- 3 Detector
- 4 Acceptance
- 5 Parton distribution functions and observables
- 6 Inclusive measurements
- 7 Flavour separation of helicity distributions
- 8 Measurements on the transversely polarised target
- 9 Nucleon spin structure
- 10 COMPASS “near” plans (2010 – 2012): muon run
- 11 COMPASS “medium” plans (\gtrsim 2012): DVCS measurements
- 12 Outlook

Flavour separation of helicity distributions (@ LO)

M. Alekseev *et al.* (COMPASS), Phys. Lett. **B660** (2008) 458.

- Difference asymmetry: $A^{h^+ - h^-} : A_d^{\pi^+ - \pi^-}(x) = A_d^{K^+ - K^-}(x) = \frac{\Delta u_v(x) + \Delta d_v(x)}{u_v(x) + d_v(x)}$
- At LO, the fragmentation functions drop out


$$Q^2 = 10 \text{ (GeV/c)}^2 \quad (\text{SIDIS} + \text{DIS})$$



$$\int_{0.006}^{0.7} (\Delta u_v + \Delta d_v) dx = 0.40 \pm 0.07 \pm 0.05$$

- Unmeasured regions contribute negligibly.
- Non-symmetric sea preferred ?

Flavour separation of helicity distributions (@ LO),....




M. Alekseev et al. (COMPASS), CERN-PH-EP/2009-008,
hep-ex 0905.2828, to appear in Phys. Lett.

$$\bullet \frac{\Delta s}{s} = A_1^d + \left(A_1^{K^+ + K^-} \right) \frac{Q/s + \alpha}{\alpha - 0.8}$$

$$\bullet \alpha = \frac{2R_{UF} + 2R_{SF}}{3R_{UF} + 2}, \quad Q = u + \bar{u} + d + \bar{d}$$



$$\bullet R_{UF} = \frac{\int D_d^{K^+}(z) dz}{\int D_u^{K^+}(z) dz}, \quad R_{SF} = \frac{\int D_s^{K^+}(z) dz}{\int D_u^{K^+}(z) dz}$$



SIDIS: $\Delta s(x)$ compatible with zero !

Flavour separation of helicity distributions (@ LO),....

NEW: proton data 2007 full deuteron statistics




Polarisation distribution of gluons, $\Delta g(x)$

- Scaling violation of g_1 (QCD fits, world data).
 - Direct measurements – via the cross section asymmetry for the photon–gluon fusion (PGF) with subsequent fragmentation into:
 - charm mesons, $q \equiv c$, (max. @ low Q^2 , perturbative low statistics, few theoretical assumptions;

$$A_{\text{meas}} = p_B \, p_T \, f \, a_{LL} \, \frac{\sigma_{PGF}}{\sigma_{PGF} + \sigma_{BGD}} \, \frac{\Delta g}{g} + A_{BGD}$$

- a pair of hadrons of large p_T , $q \equiv u, d, s$, separately for low- and high Q^2 (perturbative scale: e.g. p_T): high statistics, several quantities from MC.


$$A_{\text{meas}} = p_B \cdot p_T \cdot f \left[R_{PGF} \cdot a_{LL}^{PGF} \cdot \frac{\Delta g}{g} + R_{LO} \cdot D \cdot A_1^{LO} + R_{QCDC} \cdot a_{LL}^{QCDC} \cdot A_1^{LO} \right]$$



COMPASS QCD analysis of inclusive g_1^d

V.Yu. Alexakhin (COMPASS) et al. Phys Lett B647 (2007) 8


- Two programs: DGLAP evolution of structure functions and evolutions of moments
- NLO \overline{MS} scheme
- World data: 9 experiments, 230 data points (43 from COMPASS)
- Two solutions, $\Delta G > 0$ and $\Delta G < 0$ describe data equally well.



Quark polarisation from COMPASS data only (@ $Q^2 = 3 \text{ (GeV/c)}^2$):



$$a_0 = 0.35 \pm 0.03(\text{stat.}) \pm 0.05(\text{syst.}) \text{ and gluon polarisation: } |\Delta G| \approx 0.2 - 0.3$$

Direct Δg measurements; open charm production




$$D^0 \rightarrow K + \pi,$$

$$D^* \rightarrow D^0 + \pi_S \rightarrow K + \pi + \pi_S$$



- Choose $D^0 \rightarrow K\pi$ ($\text{BR} \sim 4\%$); pions and kaons identified by RICH.
- Combinatorial background significantly reduced for the $D^* \rightarrow D^0 + \pi_S \rightarrow K + \pi + \pi_S$.
- Charm in the nucleon neglected.
- A weighting method used to optimise the $\Delta g(x)$ extraction
- Recently added: $D^0 \rightarrow K\pi(\pi^0)$, D^* decays with K below RICH threshold of 9 GeV

Summary of the gluon polarisation measurements



At $x_g \sim 0.1$, $\Delta g/g$ is compatible with zero! Qualitative agreement with RHIC results.

Outline

- 1 Collaboration and programmes
- 2 History and plans
- 3 Detector
- 4 Acceptance
- 5 Parton distribution functions and observables
- 6 Inclusive measurements
- 7 Flavour separation of helicity distributions
- 8 Measurements on the transversely polarised target**
- 9 Nucleon spin structure
- 10 COMPASS “near” plans (2010 – 2012): muon run
- 11 COMPASS “medium” plans (\gtrsim 2012): DVCS measurements
- 12 Outlook

Properties of transversity

Properties of $\Delta_T q(x)$:

- is chiral-odd \implies hadron(s) in final state needed to be observed
- simple QCD evolution since no gluons involved
- related to GPD
- sum rule for transverse spin
- first moment gives “tensor charge” (now being studied on the lattice)

Asymmetry measured e.g. via the Collin's asymmetry (asymmetry in the distribution of hadrons):

$$N_h^\pm(\phi_c) = N_h^0 [1 \pm p_T D_{NN} A_{Coll} \sin \phi_c]$$

which in turn gives at LO:

$$A_{Coll} \sim \frac{\sum_q e_q^2 \cdot \Delta_T q \cdot \Delta_T^0 D_q^h}{\sum_q e_q^2 \cdot q \cdot D_q^h}$$




But transverse fragmentation functions $\Delta_T^0 D_q^h$ needed to extract $\Delta_T q(x)$ from the Collin's assymmetry! Recently those FF measured by BELLE.

Properties of the Sivers process: it is related to L_q in the proton. Fundamental !

Results for the transverse asymmetries

NEW data for the proton target; full 2007 statistics

COMPASS 2007 proton data



- Collins 1-h asymmetries for proton large at $x \gtrsim 0.1$, consistent with HERMES
- Sivers 1-h asymmetries for proton compatible with 0, contrary to HERMES
- 2-h asymmetry for proton large in the valence region; HERMES sees less.

- COMPASS deuteron data: both Collins and Sivers asymmetries very small.
These data + Hermes + Belle: $\Rightarrow \Delta_T u + \Delta_T d \sim 0$
- First $\Delta_T q$ global analyses performed (M. Anselmino *et al.*).

Outline

- 1 Collaboration and programmes
- 2 History and plans
- 3 Detector
- 4 Acceptance
- 5 Parton distribution functions and observables
- 6 Inclusive measurements
- 7 Flavour separation of helicity distributions
- 8 Measurements on the transversely polarised target
- 9 Nucleon spin structure**
- 10 COMPASS “near” plans (2010 – 2012): muon run
- 11 COMPASS “medium” plans (\gtrsim 2012): DVCS measurements
- 12 Outlook

Nucleon spin decomposition

$$\frac{1}{2} = \frac{1}{2}\Delta\Sigma + \Delta G + L$$

Are we approaching the solution of the “proton spin puzzle”?

- Restoration of $\Delta\Sigma=0.6$ via the axial anomaly improbable.

COMPASS @ 3 GeV²: $a_0 = 0.35 \pm 0.03 \pm 0.05$

As a consequence of the “axial anomaly” the measured quantity is:

$$a_0(Q^2) = \Delta\Sigma^{AB} - \left(\frac{3\alpha_s}{2\pi}\right)\Delta G(Q^2)$$

and the “spin crisis” can be solved ($\Delta\Sigma \sim 0.6$) if $\Delta G \sim 2.2$ (and $L \sim -2$) at $Q^2 = 3$ GeV².


- Global, consistent NLO analysis of ΔG needed.
- Independent measurement of L necessary
(\Rightarrow DVCS, lattice QCD?).
- All candidates are contributing about equally to the nucleon spin?

Outline

- 1 Collaboration and programmes
- 2 History and plans
- 3 Detector
- 4 Acceptance
- 5 Parton distribution functions and observables
- 6 Inclusive measurements
- 7 Flavour separation of helicity distributions
- 8 Measurements on the transversely polarised target
- 9 Nucleon spin structure
- 10 COMPASS "near" plans (2010 – 2012): muon run**
- 11 COMPASS "medium" plans (\gtrsim 2012): DVCS measurements
- 12 Outlook

Plans for the 2010–2012: muon run

Addendum to the COMPASS proposal (CERN-SPSC-2009-025)
for 1+1 year μ run on protons, \perp and \parallel polarised.



COMPASS data
(+ and - hadrons)
on protons

↔ projected precision


(blue points - HERMES)

Outline


- 1 Collaboration and programmes
- 2 History and plans
- 3 Detector
- 4 Acceptance
- 5 Parton distribution functions and observables
- 6 Inclusive measurements
- 7 Flavour separation of helicity distributions
- 8 Measurements on the transversely polarised target
- 9 Nucleon spin structure
- 10 COMPASS "near" plans (2010 – 2012): muon run
- 11 COMPASS "medium" plans (\gtrsim 2012): DVCS measurements**
- 12 Outlook

3D picturing of the proton via GPD


D. Mueller, X. Ji, A. Radyushkin, A. Belitsky, ...
 M. Burkardt, ... Interpretation in impact parameter space



Proton form factors,
 transverse charge &
 current densities




Correlated quark momentum
 and helicity distributions in
 transverse space - **GPDs**



Structure functions,
 quark **longitudinal**
 momentum & helicity
 distributions


Access GPD through the DVCS mechanism




- Four GPDs ($H, E, \tilde{H}, \tilde{E}$) for each flavour and for gluons
- Factorisation proven for σ_L only
- All depend on 3 variables: x, ξ, t ; DIS @ $\xi = t = 0$
- H, \tilde{H} conserve nucleon helicity
 E, \tilde{E} flip nucleon helicity
- H, E refer to unpolarised distributions

- H, E accessed in vector meson production *via* A_{UT} asymmetries
- \tilde{H}, \tilde{E} accessed in pseudoscalar meson production *via* A_{UT} asymmetries
- All 4 accessed in DVCS (γ production) in $A_C, A_{LU}, A_{UT}, A_{UL}$
- Integrals of $H, E, \tilde{H}, \tilde{E}$ over x give Dirac-, Pauli-, axial vector- and pseudoscalar vector form factors resp.
- **Important:** $J_z^q = \frac{1}{2} \int dx x [H^q(x, \xi, t=0) + E^q(x, \xi, t=0)] = \frac{1}{2} \Delta \Sigma + L_z^q$ (X. Ji)

DVCS ($\mu^- p \rightarrow \mu^+ p \gamma$) kinematics




COMPASS DVCS programme



- μ^\pm beams available with opposite polarisations
- Energies: 100/190 GeV
- LH2 target, 2.5 m long
 $L = 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$
- Lumi limits Q^2 to $\sim 8 (\text{GeV}/c)^2$
- ENC@FAIR, $E_e=3 \text{ GeV}$, $E_p=15 \text{ GeV}$ is equivalent to $E_\mu=100 \text{ GeV}$

DVCS and background



$$d\sigma(\mu p \rightarrow \mu p \gamma) \sim |T_{DVCS}|^2 + |T_{BH}|^2 + \text{interference}$$

$$d\sigma = d\sigma^{BH} + d\sigma_{unpol}^{DVCS} + P_\mu d\sigma_{pol}^{DVCS} + e_\mu a^{BH} \text{Re} T^{DVCS} + e_\mu P_\mu a^{BH} \text{Im} T^{DVCS}$$

Observables:


$$D_{U,CS} \equiv d\sigma(\mu^{+\downarrow}) - d\sigma(\mu^{-\uparrow}) \implies \text{Re}(F_1 H)$$

$$S_{U,CS} \equiv d\sigma(\mu^{+\downarrow}) + d\sigma(\mu^{-\uparrow}) \implies \text{Im}(F_1 H)$$


$$d\sigma(\phi, \phi_s) - d\sigma(\phi, \phi_s + \pi) \implies \text{Im}(F_2 H - F_1 E)$$

DVCS and background, ...c.d.

At $E=160 \text{ GeV}$, $Q^2=2 \text{ GeV}^2$ $|t|=0.1 \text{ GeV}^2$




$x=0.01$




BH dominates
(reference)

$x=0.04$



BH and DVCS at the same level
DVCS boosted by interference
 $\text{Re } T^{\text{DVCS}}$ or $\text{Im } T^{\text{DVCS}}$


$x=0.1$



DVCS dominates
study of $d\sigma^{\text{DVCS}}/dt$
(not possible at JLab)

Transverse imaging of the nucleon

LOI CERN – SPSC – 2009-003



- $S_{U,CS}$ integrated over ϕ , BH substr.
 $\Rightarrow d\sigma^{DVCS}/dt \sim \exp(-B|t|)$
- $B(x) = b_0 + 2\alpha' \ln(x_0/x)$
- α' depends on component (models)
- For flavour f :
 $B^f(x) \sim 1/2 \langle (b_\perp^f)^2 \rangle(x)$

GPD at COMPASS (proposal); two phases

PHASE 1

- In \sim 2012.
- To constrain H and to get $d\sigma/dt \rightarrow$ transverse imaging of the nucleon.
- Measured: $D_{U,CS}$ and $S_{U,CS}$; $\mu^{+\downarrow}, \mu^{-\uparrow}$ beams + unpolarised, long LH_2 (\equiv proton) target.
- To be designed and built: ~ 2.5 m LH target and ~ 4 m Recoil Proton Detector.

PHASE 2

- In \sim 2014.
- To constrain E .
- Measured: $d\sigma(\phi, \phi_s) - d\sigma(\phi, \phi_s + \pi)$; μ^+ beam and transversely polarised NH_3 (\equiv proton) target.
- To be designed and built: polarised transverse target and associated RPD.

Outline

- 1 Collaboration and programmes
- 2 History and plans
- 3 Detector
- 4 Acceptance
- 5 Parton distribution functions and observables
- 6 Inclusive measurements
- 7 Flavour separation of helicity distributions
- 8 Measurements on the transversely polarised target
- 9 Nucleon spin structure
- 10 COMPASS “near” plans (2010 – 2012): muon run
- 11 COMPASS “medium” plans (\gtrsim 2012): DVCS measurements
- 12 Outlook

Outlook

COMPASS takes data since 2002 and is the only large fixed-target experiment @ CERN now. Energy larger than HERMES and physics processes different than that of RHICspin.

- Muon programme on proton and deuteron:

- results of spin dependent structure function $g_1(x, Q^2)$;
- polarisation distributions of quarks (valence, sea), Δq , and gluons, Δg ;
- measurements of transversity, $\Delta_T q$, and of the Sivers process;
- several other measurements: exclusive ρ production, Λ polarisation, azimuthal asymmetries;

- Hadron programme:


- data taking in 2008 with 190 GeV π and in 2009 with 190 GeV π , K and p on liquid H_2 target;
- Search for exotic mesons and glueballs, diffractive dissociation and central production.

- Future:


- Proposal addendum for the μ run on the \perp and $||$ polarised protons (2010–2012);
- Letter-of-Intent for the DVCS and Drell-Yan running $\gtrsim 2012$.

SPARE

LSS06 QCD analysis including COMPASS g_1^d



$\Delta G > 0$



Can we ever tell the sign of ΔG ? ...


Figures from E. Leader, DIS2008




LSS06 QCD analysis including COMPASS g_1^d ...cont'd

... except at an ep collider ?

LSS06, $\Delta G > 0$



LSS06, $\Delta G < 0$




Figures from E. Leader, DIS2008

Direct $\Delta G/G$ measurements; high p_T hadrons @ $Q^2 < 1 \text{ GeV}^2$

E.S. Ageev (COMPASS) *et al.* Phys. Lett. B 633 (2006) 25

Resolved photons



Link to DIS and elastic form factors

DIS at $\xi=t=0$

$$H^q(x,0,0) = q(x)$$


$$\tilde{H}^q(x,0,0) = \Delta q(x)$$

Form factors (sum rules)

$$\int dx \sum_q [H^q(x, \xi, t)] = F_1(t) \text{ Dirac f.f.}$$

$$\int dx \sum_q [E^q(x, \xi, t)] = F_2(t) \text{ Pauli f.f.}$$

$$\int_{-1}^1 dx \tilde{H}^q(x, \xi, t) = G_{A,q}(t), \quad \int_{-1}^1 dx \tilde{E}^q(x, \xi, t) = G_{P,q}(t)$$



Angular Momentum Sum Rule

$$J^q = \frac{1}{2} - J^G = \frac{1}{2} \int_{-1}^1 dx [H^q(x, \xi, 0) + E^q(x, \xi, 0)]$$

X. Ji, Phy.Rev.Lett.78,610(1997)

Slide from V.D. Volker, LANL 2007