

# COMPASS-II: a Facility to study QCD



a fixe target experiment  
at the CERN SPS

~ 250 physicists  
from 24 Institutions  
of 13 Countries

**C**OMMON  
**M**UON and  
**P**ROTON  
**A**PPARATUS for  
**S**TRUCTURE and  
**S**PECTROSCOPY



*Nicole d'Hose, CEA-Saclay, for the COMPASS Collaboration*

*At The fifth workshop on hadron physics in China and Opportunities in US*

*Huangshan, China, July 2, 2013*

# COMPASS-II: a Facility to study QCD



**COMMON**  
**MUON** and  
**PROTON**  
**APPARATUS** for  
**STRUCTURE** and  
**SPECTROSCOPY**

## Long Term Plans for at least 5 years (starting in 2012)

- ✓ Primakoff **with  $\pi$ , K beam** → Test of Chiral Perturb. Theory
- ✓ Drell-Yan **with  $\pi$  beams** → Transverse Momentum Dependent PDFs
- ✓ DVCS & HEMP **with  $\mu$  beams** → Transv. Position Dependent GPDs
- ✓ SIDIS (with GPD prog.) → Strange PDF and Transv. Mom. Dep. PDFs

2012


*LHC shutdown*

2015

2016-17


(+1month in 2012)

**COMPASS:** Versatile facility to study QCD  
with hadron ( $\pi^\pm$ ,  $K^\pm$ ,  $p$  ...) and lepton (polarized  $\mu^\pm$ ) beams  
of  $\sim 200$  GeV for hadron spectroscopy and  
hadron structure studies using SIDIS, DY, DVCS, DVMP...



# The COMPASS experiment at CERN

NIM A 577 (2007) 455




Two stage magnetic spectrometer for **large angular & momentum acceptance**

Variety of tracking detectors to cope with all particles from  $\theta = 0$  to  $\theta \approx 200\text{mrad}$

Particle identification with:

- Ring Imaging Cerenkov Counter
- Electromagnetic and Hadronic calorimeters
- Hadron absorbers

**Targets:** polarized  ${}^6\text{LiD}$ ,  $\text{NH}_3$  (consecutive cells of  $\neq$  polarisation) but also  $\text{LH}_2$ , Ni




## Pion Polarisabilities and Chiral predictions

The pion: Goldstone boson (spontaneous breaking of chiral symmetry)  
lightest quark-gluon bound state system

→ understanding its internal structure is a fundamental challenge

The polarisabilities give the deformation of the pion shape by an EM field



$$\alpha_{\pi} > 0$$

$$S=0 \text{ diamagnetic contr. } \beta_{\pi} < 0$$

2-loop ChPT prediction:

$$\alpha_{\pi} + \beta_{\pi} = (0.2 \pm 0.1) 10^{-4} \text{ fm}^3$$

$$\alpha_{\pi} - \beta_{\pi} = (5.7 \pm 1.0) 10^{-4} \text{ fm}^3$$

≠ methods:


experiments:  $\alpha_{\pi} - \beta_{\pi}$  from 4 to 14  $\cdot 10^{-4} \text{ fm}^3$

$\gamma p \rightarrow \gamma \pi^+ n$  MAINZ exper. EPJA23 (2005) (≠ ChPT pred.)

$\gamma \gamma \rightarrow \pi^+ \pi^-$  (after Mark-II) future GLUEX experiment at JLab

$\pi^- \gamma \rightarrow \pi^- \gamma$  (after Serpukov) COMPASS experiment (the most direct measurement)

# Primakoff experiments with $\pi$ , K or inverse Compton Scattering on $\pi$ , K



The chiral perturbation theory (ChPT) predicts the low-energy behavior of the cross section with  $s$  varying from threshold ( $m_\pi^2$ ) to a few  $m_\pi^2$

$$\frac{d\sigma_{\pi\gamma}}{d\Omega_{\text{cm}}} = \left[ \frac{d\sigma_{\pi\gamma}}{d\Omega_{\text{cm}}} \right]_{\text{point-like}} + C \cdot \frac{(s - m_\pi^2)}{s^2} \cdot \mathcal{P}(\alpha_\pi, \beta_\pi)$$

**Deviation due to  $\pi$  polarisabilities**

*the point-like cross section is measured with the muon beam*

# Pion Polarisabilities measurement

$$\frac{d\sigma_{\pi\gamma}}{d\Omega_{\text{cm}}} = \left[ \frac{d\sigma_{\pi\gamma}}{d\Omega_{\text{cm}}} \right]_{\text{pt}} + C \cdot \frac{(s - m_\pi^2)}{s^2} \cdot \left( (1 - \cos\theta_{\text{cm}})^2 (\alpha_\pi - \beta_\pi) + (1 + \cos\theta_{\text{cm}})^2 (\alpha_\pi + \beta_\pi) \right) \frac{s^2}{m_\pi^4} + (1 - \cos\theta_{\text{cm}})^3 (\alpha_2 - \beta_2) \frac{(s - m_\pi^2)^2}{24s}$$

at backward      or      forward angle

2009 data

$$\alpha_\pi - \beta_\pi \text{ (in } 10^{-4} \text{ fm}^3\text{)} = 3.7 \pm 1.4_{\text{stat}} \pm 1.6_{\text{syst}}$$

**AGREEMENT WITH CHPT**  
**STILL PRELIMINARY**

2012 data, 3 components


$$(\alpha_\pi - \beta_\pi)$$

$$(\alpha_\pi + \beta_\pi)$$


$$(\alpha_2 - \beta_2)$$

can be measured with  
an accuracy of 10%


and kaon polarizabilities



## Deep Inelastic Scattering



While unpolarised light quark PDF well constrained, strange quark distributions are not so well known




# Semi-Inclusive Deep Inelastic Scattering

- Semi-Inclusive DIS measurements
- with polarized targets (2002-2011)
  - with a pure proton target (with GPD program)

Use of RICH detector and Calorimeters

Charge separation and identification  $K^+$ ,  $K^-$ ,  $K^0$ ,  $\pi^+$ ,  $\pi^-$ ,  $\pi^0$ ,  $\Lambda$ ...

Major progress as compared to previous experiments  
to strange PDFs:  $s(x)$  and  $\Delta s(x)$



Hadron multiplicities at LO

$$\frac{dN^h(x, z, Q^2)}{dN^{DIS}} = \frac{\sum_q e_q^2 q(x, Q^2) D_q^h(z, Q^2)}{\sum_q e_q^2 q(x, Q^2)}$$

PDF  
depend on  $x$


quark Fragmentation Function  
depend on  $z$  (fraction of energy  
of the outgoing hadron)

Final goal: extensive measurements  $(x, z, Q^2, p_T^h)$

to provide inputs to NLO global analysis for both PDF and FF


# SIDIS and multiplicities

Charged  $\pi^+$  and  $\pi^-$  multiplicities vs  $z$  in  $(x,y)$  bins




# SIDIS and multiplicities

## Charged $K^+$ and $K^-$ multiplicities vs $z$ in $(x,y)$ bins



# Projection for the strange PDF $s(x)$


LO analysis from COMPASS data alone integrated over  $z$



Projection for 1 week with 2.5m  $\text{LH}_2$  target  $\rightarrow$  high statistics

# SIDIS and azimuthal asymmetries


Asymmetries in the azimuthal angle  $\phi_h$  of the outgoing hadron around the virtual photon can reveal **quark transverse spin** and **quark transverse momentum ( $k_T$ )** effects beyond the collinear approximation



At leading twist, not only  $f_1(x, k_T)$ ,  $g_{1L}(x, k_T)$ ,  $h_1(x, k_T)$  but also 5 other **Transverse Momentum Dependent PDF (TMD ( $x, k_T$ ))** which do not survive after integration on  $k_T$

## 2 examples of TMDs


The **Boer-Mulders** function



correlates the quark  $k_T$  and the quark transverse spin (unpol N)

*Chiral-odd and T-odd*

The **Sivers** function




correlates the quark  $k_T$  and the nucleon spin (transv. Pol. N)

*Chiral-even and T-odd*

# Sivers asymmetry on transv. pol. proton

$$A_{Siv}^p \sin(\phi_h - \phi_S) \approx \frac{\sum_q e_q^2 f_{1T}^{\perp q} \otimes D_{1q}^h}{\sum_q e_q^2 f_1^q \otimes D_{1q}^h}$$



In region of overlap, agreement with HERMES, but smaller strength  
*to be done soon: multidimensional analysis ( $x, z, Q^2, p_T^h$ )*


# Boer-Mulders and Cahn effects on unpol. deuteron

$$A^{\cos(2\phi_h)}$$

Vincenzo Barone,<sup>1,2</sup> Stefano Melis,<sup>1,2</sup> and Alexei Prokudin<sup>3</sup>

COMPASS Deuteron

PHYSICAL REVIEW D **81**, 114026 (2010)




a multidimensional analysis seems to indicate a strong  $z$  dependence

→ many data collected and still to be collected in SIDIS with GPD program

# After SIDIS, polarized Drell-Yan to study TMDs

Drell-Yan  $\pi^- p^\uparrow \rightarrow \mu^+ \mu^- X$



Cross sections:

In SIDIS: convolution of a TMD with a fragmentation function

In DY: convolution of 2 TMDs

$$\sigma^{DY} \propto f_{\bar{u}|\pi^-} \otimes f'_{u|p}$$

→ complementary information and universality test

# The polarized Drell-Yan process in $\pi^- p$

 $d\sigma^{DY}$ 

$$\begin{aligned} &\propto \left(1 + \int d^2k_{1T} d^2k_{2T} \mathcal{W}(k_{1T}, k_{2T}) \bar{h}_1^\perp(x_1, k_{1T}^2) \otimes h_1^\perp(x_2, k_{2T}^2) \cos 2\phi\right) \\ &+ |S_T| \left( \int d^2k_{1T} d^2k_{2T} \mathcal{X}(k_{1T}, k_{2T}) \bar{f}_1(x_1, k_{1T}^2) \otimes f_{1T}^\perp(x_2, k_{2T}^2) \sin \phi_S \right. \\ &+ \int d^2k_{1T} d^2k_{2T} \mathcal{Y}(k_{1T}, k_{2T}) \bar{h}_1^\perp(x_1, k_{1T}^2) \otimes h_{1T}^\perp(x_2, k_{2T}^2) \sin(2\phi + \phi_S) \\ &\left. + \int d^2k_{1T} d^2k_{2T} \mathcal{Z}(k_{1T}, k_{2T}) \bar{h}_1^\perp(x_1, k_{1T}^2) \otimes h_1(x_2, k_{2T}^2) \sin(2\phi - \phi_S) \right) \end{aligned}$$


→ Access to TMDs for incoming pion  $\otimes$  target nucleon  
TMD as Transversity, Sivers, Boer-Mulders, pretzelosity

Collins-Soper frame (of virtual photon)

$\theta, \phi$  lepton plane wrt hadron plane

target rest frame

$\phi_S$  target transverse spin vector /virtual photon





# Experimental check of the change of sign of TMDs confronting Drell-Yan and SIDIS results

T-odd character of the Boer-Mulders and Sivers functions

In order not to be forced to vanish by time-reversal invariance the SSA requires an interaction phase generated by a rescattering of the struck parton in the field of the hadron remnant

*Time reversal*



these functions are process dependent, they change sign to provide the gauge invariance

$$h_1^\perp(\text{SIDIS}) = -h_1^\perp(\text{DY})$$

Boer-Mulders

Sivers


$$f_{1T}^\perp(\text{SIDIS}) = -f_{1T}^\perp(\text{DY})$$

**NEED EXPERIMENTAL VERIFICATION**  
**SIGN + AMPLITUDE + SHAPE**  
**TEST OF CONSISTENCY OF THE APPROACH**


# Why DY $\pi^\pm p^\uparrow$ is very favourable at COMPASS?

$\sigma^{DY}$  dominated by the annihilation of a valence anti-quark from the pion and a valence quark from the polarised proton

$$\sigma^{DY} \propto f_{\bar{u}|\pi^-} \otimes f_{u|p}$$



$x_F$  acceptance plot



$x_p$  vs  $x_\pi$  scatter plot: in black all generated events, in blue events in acceptance


large acceptance of COMPASS in the valence quark region for p and  $\pi$  where SSA are expected to be larger

Competitive experiments at

RHIC (STAR, PHENIX) collider  $p^\uparrow p$   
 Fermilab fixed target  $p^\uparrow \Rightarrow H, pH^\uparrow \Rightarrow$   
 J-PARC fixed target  $pp^\uparrow, \pi p^\uparrow$   
 FAIR (PAX) collider  $\bar{p}^\uparrow p^\uparrow$   
 NICA collider  $p^\uparrow p^\uparrow, d^\uparrow d^\uparrow$


COMPASS has the chance to be the first experiment to collect single polarized DY

# Q<sup>2</sup> vs x phase space at COMPASS



The phase spaces of the two processes overlap at COMPASS  
→ Consistent extraction of TMD DPFs in the same region



# DY $\pi^- + p^\uparrow \rightarrow \mu^+ \mu^- + X$ and COMPASS set-up



## Key elements for a small cross section investigation at high luminosity

1. high intensity pion beam  $10^8 \pi^-$  per second on a thick target ( $\sim 1$  interaction length)
2. a hadron absorber to stop secondary particles and a beam plug to stop the non-interacting beam
3. rearrangement of the target area to place the absorber
  - a new muon trigger in the first stage spectrometer (60% of the DY acceptance)
  - a vertex detector (SciFi) to improve the cell separation
4. RICH1, Calorimetry – also important to reduce the background

# Results from DY tests in 2007-8-9 and 2012



Distributions of primary vertexes along beam axis

→ a vertex detector has been proposed


Recent test done in the condition of the future measurement with the hadron absorber. During the short data taking, the J/ψ peak and DY events were observed as expected and the two cells were distinguished.

- Target temperature OK
- Detector occupancies OK
- Radioprotection limits respected
- Agreement with simulations

# Predictions for Drell-Yan at COMPASS

$$A_T^{\sin \phi_S}(x_a, x_b) = \frac{2}{f |S_T|} \frac{\int d\phi_S d\phi \frac{dN(x_a, x_b, \phi, \phi_S)}{d\phi d\phi_S} \sin \phi_S}{N(x_a, x_b)}$$

Sivers asymmetry in the safe dimuon mass region  $4 < M_{\mu+\mu^-} < 9$  GeV



## 2 years of data

190 GeV pion beam

$6 \cdot 10^8 \pi/\text{spill}$  (of 9.6s)

1.1 m transv. pol.  $\text{NH}_3$  target

Lumi =  $1.2 \cdot 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$

## Red solid and dot-dashed line

Anselmino et al., PRD79 (2009)

## Black solid and dashed:

Efremov et al., PLB612 (2005)

## Black dot-dashed:

Collins et al., PRD73 (2006)

## Squares:

Bianconi et al., PRD73 (2006)

## Green short-dashed:



Bacchetta et al., PRD78 (2008)

# Predictions for Drell-Yan at COMPASS

$$4. \leq M_{\mu\mu} \leq 9. \text{ GeV}/c^2$$



Sivers

M. Anselmino  
et al, Phys.  
Rev. D 79,  
054010 (2009)





Boer-  
Mulders  
B. Zhang et al,  
Phys. Rev. D  
77, 054011  
(2008)

BM  $\otimes$   
pretz.



BM  $\otimes$   
transv.  
A. N. Sissakian,  
Phys. Part.  
Nucl. 41,  
64-100 (2010)


The first ever polarised Drell-Yan experiment sensitive to TMDs



Beyond collinear approximation -> Trans. Position ( $b_{\perp}$ ) Dependent GPD in Excl. React.  
 -> as Trans. Momentum ( $k_{\perp}$ ) Dep. PDF or TMD in SIDIS & DY

**3D tomography**

# Kinematic domain ( $Q^2$ , $x_B$ ) for GPDs



## COMPASS unique for GPDs

CERN High energy muon beam

✓ 100 - 190 GeV

✓  $\mu^{\downarrow}$  and  $\mu^{\uparrow}$  available

✓ 80% Polarisation  
with opposite polarization

✓  $4.6 \cdot 10^8 \mu^+$

➔ Lumi =  $10^{32} \text{ cm}^{-2} \text{ s}^{-1}$   
with 2.5m LH2 target


- Explore the intermediate  $x_{Bj}$  region
- Uncovered region between ZEUS+H1 & HERMES + Jlab before new colliders may be available

**It's time to show the impact of COMPASS**

**=> goal of the 2012 DVCS pilot run**


# Upgrades of the COMPASS spectrometer


DVCS :  $\mu p \rightarrow \mu' p \gamma$



New equipments:

- 2.5m LH2 target
- 4m ToF Barrel CAMERA
- ECALO





ECAL2

ECAL1

ECAL0

CAMERA recoil proton detector  
surrounding the 2.5m long  
LH2 target

$\mu^\pm$




18-10-2012

## **Constraints on the GPD H**


**with recoil proton detection and hydrogen target**

- ❖ **Very first tests in 2008-9**
- ❖ **1 month in november 2012**
- ❖ **2 years 2016-17**

# Contributions of DVCS and BH at $E_\mu = 160$ GeV



$$d\sigma \propto |T^{DVCS}|^2 + |T^{BH}|^2 + \text{Interference Term}$$



Monte-Carlo Simulation for COMPASS set-up with only ECAL1+2




← Missing DVCS acceptance without ECAL0

**BH dominates**  
 excellent reference yield

**study of Interference**  
 →  $\text{Re } T^{DVCS}$   
 or  $\text{Im } T^{DVCS}$

**DVCS dominates**  
 study of  $d\sigma^{DVCS}/dt$   
 → Transverse Imaging

# 2009 DVCS test run (10 days, short RPD+target)



$$\epsilon_{\mu p \rightarrow \mu' \gamma p} \approx 35\%$$

$\times (0.8)^4$  for SPS + COMPASS avail. + trigger eff + dead time

$$\epsilon_{\text{global}} \approx 0.14 \quad \text{confirmed } \epsilon_{\text{global}} = 0.1$$

as assumed for COMPASS II predictions

**54 evts**  $\approx$  20 BH

+ **22 DVCS**

+ about **12  $\gamma$**  from  $\pi^0$

# Deeply Virtual Compton Scattering

$$d\sigma_{(\mu p \rightarrow \mu p \gamma)} = d\sigma^{BH} + d\sigma^{DVCS}_{unpol} + \cancel{P_\mu d\sigma^{DVCS}_{pol}} \\ + \cancel{e_\mu a^{BH} \text{Re} A^{DVCS}} + e_\mu P_\mu a^{BH} \text{Im} A^{DVCS}$$


Phase 1: DVCS experiment to study the transverse imaging

with  $\mu^{+\downarrow}, \mu^{-\uparrow}$  beam + unpolarized 2.5m long LH2 (proton) target

$$S_{CS,U} \equiv d\sigma(\mu^{+\downarrow}) + d\sigma(\mu^{-\uparrow}) \propto d\sigma^{BH} + d\sigma^{DVCS}_{unpol} + K.s_1^{Int} \sin \phi$$


Using  $S_{CS,U}$  and BH subtraction  
and integration over  $\phi$

$$d\sigma^{DVCS}/dt \sim \exp(-B|t|)$$




# Transverse imaging at COMPASS

$$d\sigma^{DVCS} / dt \sim \exp(-B|t|)$$




**2 years of data**  
 160 GeV muon beam  
 2.5m  $\text{LH}_2$  target  
 $\epsilon_{\text{global}} = 10\%$

ansatz at small  $x_B$   
 inspired by  
 Regge Phenomenology:  
 $B(x_B) = b_0 + 2 \alpha' \ln(x_0/x_B)$   
 $\alpha'$  slope of Regge trajet



# Transverse imaging at COMPASS

$$d\sigma^{DVCS} / dt \sim \exp(-B |t|)$$



**DVCS test in 2012**

With 2 weeks  
 Using the 4m long RPD  
 + the 2.5m long LH2 target

1/20 of the complete  
 statistics

**2012:** we can determine one mean value of B  
 in the COMPASS kinematic range

# Deeply Virtual Compton Scattering


$$d\sigma_{(\mu p \rightarrow \mu p \gamma)} = \cancel{d\sigma^{BH}} + \cancel{d\sigma^{DVCS}_{unpol}} + P_{\mu} d\sigma^{DVCS}_{pol} \\ + e_{\mu} a^{BH} \mathcal{R}e A^{DVCS} + e_{\mu} P_{\mu} \cancel{a^{BH} \mathcal{I}m A^{DVCS}}$$

## Phase 1: DVCS experiment to constrain GPD H

with  $\mu^{+\downarrow}, \mu^{-\uparrow}$  beam + unpolarized 2.5m long LH2 (proton) target

$$\mathcal{D}_{CS,U} \equiv d\sigma(\mu^{+\downarrow}) - d\sigma(\mu^{-\uparrow}) \propto c_0^{Int} + c_1^{Int} \cos \phi \quad \text{and} \quad c_{0,1}^{Int} \sim \mathcal{R}e(F_1 \mathcal{H})$$

$$\mathcal{S}_{CS,U} \equiv d\sigma(\mu^{+\downarrow}) + d\sigma(\mu^{-\uparrow}) \propto d\sigma^{BH} + c_0^{DVCS} + K.s_1^{Int} \sin \phi \quad \text{and} \quad s_1^{Int} \sim \mathcal{I}m(F_1 \mathcal{H})$$



$$\xi \sim x_B / (2 - x_B)$$

Note: dominance of **H** at COMPASS kinematics

$$\mathcal{I}m \mathcal{H}(\xi, t) = \mathbf{H}(x = \xi, \xi, t)$$

$$\mathcal{R}e \mathcal{H}(\xi, t) = \mathcal{P} \int dx \frac{\mathbf{H}(x, \xi, t)}{x - \xi} = \mathcal{P} \int dx \frac{\mathbf{H}(x, x, t) + \mathbf{D}(t)}{x - \xi}$$

$\mathcal{R}e$  part of the *Compton Form Factors* linked to the  $\mathcal{D}$  term

# Deeply Virtual Compton Scattering

$$d\sigma_{(\mu p \rightarrow \mu p \gamma)} = \cancel{d\sigma^{BH}} + \cancel{d\sigma^{DVCS}_{unpol}} + P_{\mu} d\sigma^{DVCS}_{pol} \\ + e_{\mu} a^{BH} \mathcal{R}e A^{DVCS} + e_{\mu} P_{\mu} \cancel{a^{BH} \mathcal{I}m A^{DVCS}}$$

## Phase 1: DVCS experiment to constrain GPD H

with  $\mu^{+\downarrow}, \mu^{-\uparrow}$  beam + unpolarized 2.5m long LH2 (proton) target

$$\mathcal{D}_{CS,U} \equiv d\sigma(\mu^{+\downarrow}) - d\sigma(\mu^{-\uparrow}) \propto c_0^{Int} + c_1^{Int} \cos \phi \quad \text{and} \quad c_{0,1}^{Int} \sim \mathcal{R}e(F_1 \mathcal{H}) \\ \mathcal{S}_{CS,U} \equiv d\sigma(\mu^{+\downarrow}) + d\sigma(\mu^{-\uparrow}) \propto d\sigma^{BH} + c_0^{DVCS} + K.s_1^{Int} \sin \phi \quad \text{and} \quad s_1^{Int} \sim \mathcal{I}m(F_1 \mathcal{H})$$







Angular decomposition of **sum** and **diff** of the **DVCS cross section** will provide unambiguous way to separate the  $\mathcal{R}e$  and  $\mathcal{I}m$  of the *Compton Form Factors* from higher twist contributions

$$D_{CS,U} \equiv d\sigma(\mu^{+\downarrow}) - d\sigma(\mu^{-\uparrow}) \propto c_0^{Int} + c_1^{Int} \cos \phi \quad \text{and} \quad c_{0,1}^{Int} \sim \text{Re}(F_1 \mathcal{H})$$

$A_{CS,U}^{\cos\phi}$  related to  $c_1^{Int}$

Predictions with  
**VGG** and **D. Mueller**

$\text{Re}(F_1 \mathcal{H}) > 0$  at H1  
< 0 at HERMES/JLab  
Value of  $x_B$  for the node?



**COMPASS** 2 years of data  $E_\mu = 160 \text{ GeV}$   $1 < Q^2 < 8 \text{ GeV}^2$  With ECAL2 + ECAL1 + ECAL0

HERMES 10 years of data


JLab

# With transv. polarized target **Constraints on other GPDs**

$\sigma$

## Chiral-even GPDs

$$H \rightarrow q$$




$A_{UT}^{\sin(\phi - \phi_s)}$

$$E \leftrightarrow f_{1T}^\perp$$

Sivers correlates

quark  $k_T$  and nucleon spin (transv. pol. N)





## Chiral-odd GPDs

$A_{UT}^{\sin(\phi_s)}$

$$H_T \leftrightarrow h_1$$

Transversity correlates


quark spin and nucleon spin (transv. pol. N)



$$\bar{E}_T = 2\tilde{H}_T + E_T \leftrightarrow h_1^\perp$$

Boer-Mulders correlates

quark  $k_T$  and quark transverse spin (unpol N)





# DVCS –Transv. Polar. Target

COMPASS-II (future addendum) : with  $\mu^{+\downarrow}$ ,  $\mu^{-\uparrow}$  beam and transversely polarized NH3 (proton)

$$\mathcal{D}_{CS,T} \equiv d\sigma_T(\mu^{+\downarrow}) - d\sigma_T(\mu^{-\uparrow})$$

$$\propto \text{Im}(F_2 \mathcal{H} - F_1 \mathcal{E}) \sin(\phi - \phi_S) \cos \phi$$




**2 years of data**

160 GeV muon beam

1.2 m polarised NH<sub>3</sub> target

$\epsilon_{\text{global}} = 10\%$

$A_{CS,T}^{\sin(\phi - \phi_S) \cos \phi}$



# Summary for GPD @ COMPASS

## GPDs investigated with Hard Exclusive Photon and Meson Production

COMPASS-II 2016-17: with LH<sub>2</sub> target + RPD (phase 1)  $\mu^{+\downarrow}$ ,  $\mu^{-\uparrow}$  160 GeV

- ✓ the t-slope of the DVCS and HEMP cross section  
→ transverse distribution of partons
- ✓ the Beam Charge and Spin Sum and Difference  
→  $\mathcal{R}e T^{DVCS}$  and  $\mathcal{I}m T^{DVCS}$  for the GPD H determination
- ✓ Vector Meson  $\rho^0, \rho^+, \omega, \Phi$
- ✓ Pseudo-saclar  $\pi^0$

Using the 2007-10 data: transv. polarized NH<sub>3</sub> target without RPD

In a future addendum > 2017: transv. polarised NH<sub>3</sub> target with RPD (phase 2)

- ✓ the Transverse Target Spin Asymm  
→ GPD E and chiral-odd (transverse) GPDs

**For the next 10 years, before any collider is available,  
and complementary to Jlab 12 GeV,  
COMPASS@CERN can be a major player in QCD physics  
using its unique high energy (~200 GeV) hadron  
and polarised positive and negative muon beams**



# Boer-Mulders and Cahn effects on unpol. proton

$$\frac{d\sigma}{dx dy d\phi_h} = \frac{\alpha^2}{xy Q^2} \frac{1 + (1-y)^2}{2} \left[ F_{UU} + \varepsilon_1 \cos \phi_h F_{UU}^{\cos \phi_h} + \varepsilon_2 \cos 2\phi_h F_{UU}^{\cos 2\phi_h} + \lambda_\mu \varepsilon_3 \sin \phi_h F_{LU}^{\sin \phi_h} \right]$$

Cahn effect  $\rightarrow$  info on  $\langle k_T \rangle$

Boer-Mulders TMD  $\otimes$  Collins FF + Cahn effect

# Transverse imaging at COMPASS

$$d\sigma^{DVCS}/dt \sim \exp(-B|t|)$$

$$B(x_B) = \frac{1}{2} \langle r_{\perp}^2(x_B) \rangle$$

distance between the active quark and the center of momentum of spectators

## Transverse size of the nucleon


mainly dominated by  $H(x, \xi=x, t)$

$$\text{related to } \frac{1}{2} \langle b_{\perp}^2(x_B) \rangle$$


distance between the active quark and the center of momentum of the nucleon

## Impact Parameter Representation

$$q(x, b_{\perp}) \leftrightarrow H(x, \xi=0, t)$$



Note  $0.65 \text{ fm} = \sqrt{2/3} \times 0.8 \text{ fm}$



# Deeply Virtual Compton Scattering

$$d\sigma_{(\mu p \rightarrow \mu p \gamma)} = \cancel{d\sigma^{BH}} + \cancel{d\sigma^{DVCS}_{unpol}} + P_{\mu} d\sigma^{DVCS}_{pol} \\ + e_{\mu} a^{BH} \mathcal{R}e A^{DVCS} + e_{\mu} P_{\mu} \cancel{a^{BH} \mathcal{I}m A^{DVCS}}$$

## Phase 1: DVCS experiment to constrain GPD H

with  $\mu^{+\downarrow}, \mu^{-\uparrow}$  beam + unpolarized 2.5m long LH2 (proton) target

$$\mathcal{D}_{CS,U} \equiv d\sigma(\mu^{+\downarrow}) - d\sigma(\mu^{-\uparrow}) \propto c_0^{Int} + c_1^{Int} \cos \phi \quad \text{and} \quad c_{0,1}^{Int} \sim \mathcal{R}e(F_1 \mathcal{H}) \\ \mathcal{S}_{CS,U} \equiv d\sigma(\mu^{+\downarrow}) + d\sigma(\mu^{-\uparrow}) \propto d\sigma^{BH} + c_0^{DVCS} + K.s_1^{Int} \sin \phi \quad \text{and} \quad s_1^{Int} \sim \mathcal{I}m(F_1 \mathcal{H})$$

Angular decomposition of **sum** and **diff** of the **DVCS cross section** will provide unambiguous way to separate the  $\mathcal{R}e$  and  $\mathcal{I}m$  of the *Compton Form Factors* from higher twist contributions


# Deeply Virtual Compton Scattering

$$d\sigma_{(\mu p \rightarrow \mu p \gamma)} = \cancel{d\sigma^{BH}} + \cancel{d\sigma^{DVCS}_{unpol}} + P_{\mu} d\sigma^{DVCS}_{pol} \\ + e_{\mu} a^{BH} \mathcal{R}e A^{DVCS} + e_{\mu} P_{\mu} \cancel{a^{BH} \mathcal{I}m A^{DVCS}}$$

## Phase 1: DVCS experiment to constrain GPD H

with  $\mu^{+\downarrow}, \mu^{-\uparrow}$  beam + unpolarized 2.5m long LH2 (proton) target

$$D_{CS,U} \equiv d\sigma(\mu^{+\downarrow}) - d\sigma(\mu^{-\uparrow}) \propto c_0^{Int} + c_1^{Int} \cos \phi \quad \text{and} \quad c_{0,1}^{Int} \sim \mathcal{R}e(F_1 \mathcal{H}) \\ S_{CS,U} \equiv d\sigma(\mu^{+\downarrow}) + d\sigma(\mu^{-\uparrow}) \propto d\sigma^{BH} + c_0^{DVCS} + K.s_1^{Int} \sin \phi \quad \text{and} \quad s_1^{Int} \sim \mathcal{I}m(F_1 \mathcal{H})$$



$$\xi \sim x_B / (2 - x_B)$$

Note: dominance of **H** at COMPASS kinematics



$$\triangleright \mathcal{I}m \mathcal{H}(\xi, t) = \mathbf{H}(x = \xi, \xi, t)$$

$$\triangleright \mathcal{R}e \mathcal{H}(\xi, t) = \mathcal{P} \int dx \mathbf{H}(x, \xi, t) / (x - \xi)$$

Related with a dispersion relation + Dterm

# Beam Charge and Spin Difference (using $\mathcal{D}_{CS,U}$ )

## Comparison to different models




**2 years of data**  
 160 GeV muon beam  
 2.5m LH<sub>2</sub> target  
 $\epsilon_{\text{global}} = 10\%$

Note: Kroll, Moutarde, Sabatié predictions are of the same order of magnitude than Mueller predictions

High precision beam flux and acceptance determination  
 Systematic error bands assuming a 3% charge-dependent effect between  $\mu^+$  and  $\mu^-$  (control with inclusive evts, BH...)


# Selection of Exclusive $\rho^0$ Production: $\mu p \rightarrow \mu' \rho^0 p$ without RPD



$1 < Q^2 < 10 \text{ GeV}^2$     $0.1 < y < 0.9$     $W > 4 \text{ GeV}$     $E_{\rho^0} > 15 \text{ GeV}$

1- Assuming both hadrons are  $\pi$   
 $0.5 < M_{\pi\pi} < 1.1 \text{ GeV}$


*To maximize the purity of the sample of  $\rho^0$  /  
non resonant  $\pi^+\pi^-$*



2- Suppression of incoherent production on quasi-free protons in  $\text{NH}_3$  polarized target  
+ Suppression of SIDIS background

$0.05 < p_t^2 < 0.5 \text{ GeV}^2$

*Contamination of about a 5% coherent production*



3- Exclusivity of the reaction

$$E_{\text{miss}} = \frac{M_X^2 - M_P^2}{2 \cdot M_P} = E_{\gamma^*} - E_{\rho^0} + t / (2 \cdot M_P)$$


$-2.5 < E_{\text{miss}} < 2.5 \text{ GeV}$

*Diffractive dissociation contamination  $\sim 14\%$   
No attempt to remove it (motivated by HERA)*

$\rightarrow$  correction for SIDIS background (5 to 40%)  
in each bin ( $x_{\text{Bj}}$ ,  $Q^2$ ,  $p_T^2$ , cell and polar. State)

# Hard Exclusive Vector Meson Production

$$A_{UT}(\rho^0_L) \propto \sqrt{|-t'|} \operatorname{Im}(\mathcal{E}^* \mathcal{H}) / |\mathcal{H}|^2$$



$$E_{\rho^0} \propto \frac{2}{3} E^u + \frac{1}{3} E^d + \frac{3}{8} E^g$$

$$E_{\omega} \propto \frac{2}{3} E^u - \frac{1}{3} E^d + \frac{1}{8} E^g$$

$$E_{\rho^+} \propto E^u - E^d - \frac{3}{8} E^g$$

Cancellation between gluon and sea contributions

$$\kappa^q = \int e^q(x) dx$$

$$\rightarrow E^{uval} \sim -E^{dval}$$

$A_{UT}(\rho^0)$  very small

$A_{UT}(\omega)$  and  $A_{UT}(\rho^+)$  should be more promising  
analysis on going for  $\omega$ ,  $\rho^+$ ,  $\phi$  and  $\gamma$


Bins in  $\Phi - \Phi_s$

asymmetry extraction

using a **1D** binned maximum likelihood fit  
after subtracting the SIDIS background

# Exclusive $\rho^0$ production on transverse polar. target

without Recoil Detection



COMPASS (NPB 865 1- July 2012)

and predictions by

Goloskokov & Kroll, EPJ C59 (2009)

# NEW ANALYSIS

Bins in  $\Phi$  and  $\Phi_s$

asymmetry extraction

using a **2D** binned maximum likelihood fit


After subtracting the SIDIS background

transv. pol. Protons

NH3 target 2007-2010


$A_{UT}$

NEW RESULTS



transv. pol. Protons

NH3 target 2007-2010

 $A_{LT}$ 


# Recoil Proton Detector CAMERA

ToF between 2 rings of scintillators  $\sigma(\text{ToF}) < 300\text{ps}$

$0.06 < t < 0.8 \text{ GeV}/c^2$


1 GHz digitization of the PMT signal  
to cope with high rate

GANDALF boards → First level trigger




upstream side

3.90m




downstream side




# calibration of CAMERA

Energy lost in Ring B





Energy lost in Ring B




# ECAL0 to enlarge the angular coverage

ECAL0 made of 200 modules ( $12 \times 12 \text{ cm}^2$ ) of 9 cells read by 9 MAPDs


56 Modules are available for the 2012 setup  
They are already calibrated (24 Oct 2012)



Invariant  $\gamma\gamma$  mass spectra  
for  $\pi^0$  production using pion beam



# The hadron absorber



Structure of the hadron absorber:

- 120cm tungsten beam plug
- aluminium conical part
- 200cm alumina ( $\text{Al}_2\text{O}_3$ )
- Stainless steel shielding sandwiches

+ absorber surrounded by  
2m of iron-free concrete on each side

