



Exclusive Measurements at COMPASS

Po-Ju Lin

CEA, Université Paris-Saclay

on behalf of the COMPASS collaboration


PANIC 2021

September 05, 2021

COMPASS Experiment



Versatile facility with hadron (π^\pm , K^\pm , p ...) & lepton (polarized μ^\pm) beams of energy from 100 to 200 GeV



LHC

COMPASS


M2

SPS

~220 physicists from 25 Institutes of 13 Countries

COmmun
Muon and
Proton
Apparatus for
Structure and
Spectroscopy

COMPASS Setup for Hard Exclusive Measurements




Muon Beams

- μ^+ & μ^- with opposite polarisation
- About $\pm 80\%$ polarisation
- Momentum: 160 GeV/c

Two-stage, large angle, and wide momentum range spectrometer. PID including hadron absorbers, RICH, HCALs, ECALs, and muon filters.


❖ NIM A 577 (2007) & NIM A 779 (2015) 69

COMPASS Setup for Hard Exclusive Measurements



~50m


Exclusive Muoproduction




Two-stage, large angle, and wide momentum range spectrometer. PID including hadron absorbers, RICH, HCALs, ECALs, and muon filters.

❖ NIM A 577 (2007) & NIM A 779 (2015) 69

COMPASS Setup for Hard Exclusive Measurements




Exclusive Muoproduction




Two-stage, large angle, and wide momentum range spectrometer. PID including hadron absorbers, RICH, HCALs, ECALs, and muon filters.

❖ NIM A 577 (2007) & NIM A 779 (2015) 69

COMPASS Setup for Hard Exclusive Measurements



Exclusive Muoproduction



Two-stage, large angle, and wide momentum range spectrometer. PID including hadron absorbers, RICH, HCALs, ECALs, and muon filters.


❖ NIM A 577 (2007) & NIM A 779 (2015) 69

- 2012 pilot run with 4-week data taking
- 2016-17 dedicated run. 2 x 6 months.


Deeply Virtual Compton Scattering



COMPASS Coverage for DVCS



Deeply Virtual Compton Scattering



$$\text{DVCS: } l + p \rightarrow l' + p' + \gamma$$


To experimentally access the information about Generalized Parton Distributions (GPDs), DVCS is regarded as the golden channel and its interference with the well-understood Bethe-Heitler process gives access to more info.

The variables measured in the experiment:


$$E_l, Q^2, x_{Bj} \sim 2\xi / (1+\xi),$$

$$t \text{ (or } \theta_{\gamma^*\gamma}) \text{ and } \phi \text{ (} \ell\ell' \text{ plane}/\gamma\gamma^* \text{ plane)}$$

**BH
Process**



Deeply Virtual Compton Scattering



- The GPDs depend on the following variables:
 - x : average longitudinal momentum fraction
 - ξ : longitudinal momentum fraction difference
 - t : four momentum transfer
(correlated to b_{\perp} via Fourier transform)
 - Q^2 : virtuality of γ^*

Sensible to 4 GPDs, with LH_2 target and small x_B coverage
 → focuses on H at COMPASS

CFP (yellow dashed arrow) points to \mathcal{H} . GPD (blue dashed arrow) points to \mathcal{H} .




$$\mathcal{H} = \int_{-1}^{+1} dx \frac{\mathbf{H}(x, \xi, t)}{x - \xi + i\epsilon} \quad \text{with } t, \xi \text{ fixed}$$

$$= \mathcal{P} \int_{-1}^{+1} dx \frac{\mathbf{H}(x, \xi, t)}{x - \xi} - i\pi \mathbf{H}(x = \pm \xi, \xi, t)$$

REAL part Imaginary part

$$\text{Re } \mathcal{H}(\xi, t) = \mathcal{P} \int dx \frac{\text{Im } \mathcal{H}(x, t)}{x - \xi} + d(t)$$

Transverse Imaging and Pressure Dist.



$$\begin{aligned}
 \text{CFP} & \rightarrow \mathcal{H} = \int_{-1}^{+1} dx \frac{\mathbf{H}(x, \xi, t)}{x - \xi + i\epsilon} \\
 \text{GPD} & \rightarrow \mathcal{H} = \int_{-1}^{+1} dx \frac{\mathbf{H}(x, \xi, t)}{x - \xi + i\epsilon} = \mathcal{P} \int_{-1}^{+1} dx \frac{\mathbf{H}(x, \xi, t)}{x - \xi} - i\pi \mathbf{H}(x = \pm \xi, \xi, t) \\
 \text{REAL part} & \rightarrow \text{Re } \mathcal{H}(\xi, t) = \mathcal{P} \int dx \frac{\text{Im } \mathcal{H}(x, t)}{x - \xi} + d(t) \\
 \text{Imaginary part} & \rightarrow \text{Im } \mathcal{H}(x, t) = \dots
 \end{aligned}$$

➤ Main background of **exclusive single photon events**: π^0 decay


➤ **Visible** (both γ detected) – subtracted

A high-energy DVCS photon candidate is combined with all detected photons with energies lower than the DVCS threshold: (4,5) GeV in Ecal (0,1) respectively




➤ **Invisible** (one γ lost) – estimated by MC

- **Semi-inclusive LEPTO 6.1**
- **Exclusive HEPGEN π^0** (GK model)

The sum of LEPTO and HEPGEN contributions is normalized to the π^0 peak in $M_{\gamma\gamma}$ of the real data



COMPASS 2016 Preliminary Results



➤ Beam charge-spin sum

$$S_{CS,U}(\phi) \equiv d\sigma(\mu^{+\leftarrow}) + d\sigma(\mu^{-\rightarrow}) = 2[d\sigma^{BH} + d\sigma_{unpol}^{DVCS} + \text{Im } I]$$

$$= 2[d\sigma^{BH} + c_0^{DVCS} + c_1^{DVCS} \cos \phi + c_2^{DVCS} \cos 2\phi + s_1^I \sin \phi + s_2^I \sin 2\phi]$$


$$c_0^{DVCS} \underset{\text{small } x_{Bj}}{\propto} 4(\mathcal{H}\mathcal{H}^* + \tilde{\mathcal{H}}\tilde{\mathcal{H}}^*) + \frac{t}{M^2} \mathcal{E}\mathcal{E}^* \rightarrow 4 (\text{Im } \mathcal{H})^2$$

model dependent

Transverse extension of partons – 2012 data




$$d\sigma^{DVCS}/d|t| \propto e^{-B|t|}$$



$$\langle r_{\perp}^2(x_{Bj}) \rangle \approx 2B(x_{Bj}) \text{ At small } x_{Bj}$$

- COMPASS: $\langle Q^2 \rangle = 1.8 \text{ (GeV/c)}^2$
- ◆ ZEUS: $\langle Q^2 \rangle = 3.2 \text{ (GeV/c)}^2$
- ▲ H1: $\langle Q^2 \rangle = 4.0 \text{ (GeV/c)}^2$
- ▼ H1: $\langle Q^2 \rangle = 8.0 \text{ (GeV/c)}^2$
- H1: $\langle Q^2 \rangle = 10. \text{ (GeV/c)}^2$



- The transverse-size evolution as a function of x_{Bj}
- ➔ Expect at least 3 x_{Bj} bins from 2016-17 data


$$\sqrt{\langle r_{\perp}^2 \rangle} = (0.58 \pm 0.04_{\text{stat}} \pm 0.01_{\text{sys}} \pm 0.04_{\text{model}}) \text{ fm}$$

With $\langle x_{Bj} \rangle = 0.056$

GPDs in Hard Exclusive Meson Production



Quark contribution




4 chiral-even GPDs: helicity of parton unchanged

$\mathbf{H}^q(x, \xi, t)$	$\mathbf{E}^q(x, \xi, t)$	For Vector Meson
$\tilde{\mathbf{H}}^q(x, \xi, t)$	$\tilde{\mathbf{E}}^q(x, \xi, t)$	For Pseudo-Scalar Meson

+ 4 chiral-odd or transversity GPDs: helicity of parton changed
(not possible in DVCS)

$\mathbf{H}_T^q(x, \xi, t)$	$\mathbf{E}_T^q(x, \xi, t)$	$\bar{\mathbf{E}}_T^q = 2 \tilde{\mathbf{H}}_T^q + \mathbf{E}_T^q$
$\tilde{\mathbf{H}}_T^q(x, \xi, t)$	$\tilde{\mathbf{E}}_T^q(x, \xi, t)$	

Gluon contribution at the same order in α_s




- Universality of GPDs, quark flavor filter
- Ability to probe the chiral-odd GPDs.
- Additional non-perturbative term from meson wave function
- In addition to nuclear structure, provide insights into reaction mechanism


2012 Exclusive π^0 Prod. on Unpolarized Proton




$$\frac{d^2\sigma}{dt d\phi_\pi} = \frac{1}{2\pi} \left[\left(\frac{d\sigma_T}{dt} + \epsilon \frac{d\sigma_L}{dt} \right) + \epsilon \cos 2\phi_\pi \frac{d\sigma_{TT}}{dt} + \sqrt{2\epsilon(1+\epsilon)} \cos \phi_\pi \frac{d\sigma_{LT}}{dt} \right]$$



A dip at small t would indicate a large impact of \bar{E}_T



Exclusive ω Production on Unpolarized Proton



Experimental angular distributions

$$\mathcal{W}^{U+L}(\Phi, \phi, \cos \Theta) = \mathcal{W}^U(\Phi, \phi, \cos \Theta) + P_b \mathcal{W}^L(\Phi, \phi, \cos \Theta)$$

15 unpolarized SDMEs in W^U and 8 polarized in W^L

$$\begin{aligned} \mathcal{W}^U(\Phi, \phi, \cos \Theta) = & \frac{3}{8\pi^2} \left[\frac{1}{2}(1 - r_{00}^{04}) + \frac{1}{2}(3r_{00}^{04} - 1) \cos^2 \Theta - \sqrt{2}\text{Re}\{r_{10}^{04}\} \sin 2\Theta \cos \phi - r_{1-1}^{04} \sin^2 \Theta \cos 2\phi \right. \\ & - \epsilon \cos 2\Phi \left(r_{11}^1 \sin^2 \Theta + r_{00}^1 \cos^2 \Theta - \sqrt{2}\text{Re}\{r_{10}^1\} \sin 2\Theta \cos \phi - r_{1-1}^1 \sin^2 \Theta \cos 2\phi \right) \\ & - \epsilon \sin 2\Phi \left(\sqrt{2}\text{Im}\{r_{10}^2\} \sin 2\Theta \sin \phi + \text{Im}\{r_{1-1}^2\} \sin^2 \Theta \sin 2\phi \right) \\ & + \sqrt{2\epsilon(1 + \epsilon)} \cos \Phi \left(r_{11}^5 \sin^2 \Theta + r_{00}^5 \cos^2 \Theta - \sqrt{2}\text{Re}\{r_{10}^5\} \sin 2\Theta \cos \phi - r_{1-1}^5 \sin^2 \Theta \cos 2\phi \right) \\ & \left. + \sqrt{2\epsilon(1 + \epsilon)} \sin \Phi \left(\sqrt{2}\text{Im}\{r_{10}^6\} \sin 2\Theta \sin \phi + \text{Im}\{r_{1-1}^6\} \sin^2 \Theta \sin 2\phi \right) \right], \\ \mathcal{W}^L(\Phi, \phi, \cos \Theta) = & \frac{3}{8\pi^2} \left[\sqrt{1 - \epsilon^2} \left(\sqrt{2}\text{Im}\{r_{10}^3\} \sin 2\Theta \sin \phi + \text{Im}\{r_{1-1}^3\} \sin^2 \Theta \sin 2\phi \right) \right. \\ & + \sqrt{2\epsilon(1 - \epsilon)} \cos \Phi \left(\sqrt{2}\text{Im}\{r_{10}^7\} \sin 2\Theta \sin \phi + \text{Im}\{r_{1-1}^7\} \sin^2 \Theta \sin 2\phi \right) \\ & \left. + \sqrt{2\epsilon(1 - \epsilon)} \sin \Phi \left(r_{11}^8 \sin^2 \Theta + r_{00}^8 \cos^2 \Theta - \sqrt{2}\text{Re}\{r_{10}^8\} \sin 2\Theta \cos \phi - r_{1-1}^8 \sin^2 \Theta \cos 2\phi \right) \right] \end{aligned}$$

➤ $\epsilon \rightarrow 1$, small W^L

2012 Exclusive ω Prod. on Unpolarized Proton



SCHC ($\lambda_\gamma = \lambda_V$)
(S-Channel Helicity Conservation)


SCHC implies:

- $r_{1-1}^1 + \text{Im} r_{1-1}^2 = 0$
 $= -0.010 \pm 0.032 \pm 0.047$ OK

- $\text{Re} r_{10}^5 + \text{Im} r_{10}^6 = 0$
 $= 0.014 \pm 0.011 \pm 0.013$ OK

- $\text{Im} r_{10}^7 - \text{Re} r_{10}^8 = 0$
 $= -0.088 \pm 0.110 \pm 0.196$ OK

all elements of classes C, D, E should be 0
for $\gamma_L^* \rightarrow \omega_T$ and $\gamma_T^* \rightarrow \omega_T$ OK within errors
NOT OBSERVED for transitions $\gamma_T^* \rightarrow \omega_L$



2012 Exclusive ρ^0 Prod. on Unpolarized Proton



SCHC ($\lambda_\gamma = \lambda_V$)

(S-Channel Helicity Conservation)


SCHC implies:

• $r_{1-1}^1 + \text{Im} r_{1-1}^2 = 0$ **OK**

• $\text{Re} r_{10}^5 + \text{Im} r_{10}^6 = 0$ **OK**

• $\text{Im} r_{10}^7 - \text{Re} r_{10}^8 = 0$ **✓OK**

• all elements of classes C, D, E should be 0 for $\gamma_L^* \rightarrow \rho_T^0$ and $\gamma_T^* \rightarrow \rho_{-T}^0$ OK within errors
NOT OBSERVED for transitions $\gamma_T^* \rightarrow \rho_L^0$



■ SDMEs COMPASS
 PRELIMINARY

Goloskokov and Kroll, EPJC 74 (2014) 2725

$$r_{00}^5 \propto \text{Re}[\langle \bar{E}_T \rangle_{LT}^* \langle H \rangle_{LL} + \frac{1}{2} \langle H_T \rangle_{LT}^* \langle E \rangle_{LL}]$$

First term dominates
→ Probes \bar{E}_T


NPE-to-UPE Asymmetry – 2012 data



$$P = \frac{2r_{1-1}^1}{1 - r_{00}^{04} - 2r_{1-1}^{04}} \approx \frac{d\sigma_T^N(\gamma_T^* \rightarrow V_T) - d\sigma_T^U(\gamma_T^* \rightarrow V_T)}{d\sigma_T^N(\gamma_T^* \rightarrow V_T) + d\sigma_T^U(\gamma_T^* \rightarrow V_T)}$$


- NPE: Natural Parity Exchange
- UPE: Unnatural Parity Exchange

NPE-to-UPE asymmetry of cross sections for transitions $\gamma_T^* \rightarrow V_T$



ρ^0

- NPE Dominance
- NPE \rightarrow GPDs E, H



ω

- NPE \approx UPE on average
- UPE Dominance at small W and p_T^2
- UPE \rightarrow GPDs \tilde{E}, \tilde{H}
- + Pion pole (dominant)

DVCS x-sections with polarized μ^+ and μ^-

- Beam charge-spin sum $\rightarrow \text{Im}\mathcal{H}(\xi,t) \rightarrow$ Transverse extension of partons as a function of x_{Bj}
- Beam charge-spin difference $\rightarrow \text{Re}\mathcal{H}(\xi,t) \rightarrow$ D-term, pressure distribution

HEMP of π^0 , ρ , ω , ϕ , J/ψ

- x-section of π^0 , SDME of ρ & $\omega \rightarrow$ Transversity GPDs \rightarrow Flavor Decomposition
- ϕ , $J/\psi \rightarrow$ Gluon GPDs


On-going analysis on 2016-17 data.





Backup Slides


Exclusive ρ^0 Production on Unpolarized Proton



COMPASS Setup for GPD Measurement




$$\text{DVCS} : \mu p \rightarrow \mu' p \gamma$$



ments:

- 2.5m LH2 target
- 4m ToF Barrel CAMERA
- ECALO



CAMERA

L=4m Ø=2m

24 inner & outer scintillators separated by about 1m
1 GHz SADC readout, 330ps ToF resolution

ECALO

ECALO: 2 × 2 m²

Shashlyk modules + MAPD readout
one module is made of 9 cells (4×4 cm²)
= 194 modules or 1746 cells

Letter of Intent - Draft 1.0: <https://arXiv.org/abs/1808.0084>

Expected to start at 2022

- Unique beam line with polarised μ^\pm and high-intensity **Pion** beam
- Possible high-intensity **antiproton** and **Kaon** beams, provided by RF-separation technique
- With upgraded apparatus

Proposed physics goals



Proton Radius
Meson PDF – gluon PDF
Proton spin structure
3D imaging (TMDs and GPDs)
Hadron spectroscopy
Anti-matter cross section

Program	Physics Goals	Beam Energy [GeV]	Beam Intensity [s^{-1}]	Trigger Rate [kHz]	Beam Type	Target	Earliest start time, duration	Hardware Additions
μp elastic scattering	Precision proton-radius measurement	100	$4 \cdot 10^6$	100	μ^\pm	high-pressure H2	2022 1 year	active TPC, SciFi trigger, silicon veto,
Hard exclusive reactions	GPD E	160	$2 \cdot 10^7$	10	μ^\pm	NH_3^\dagger	2022 2 years	recoil silicon, modified PT magnet
Input for Dark Matter Search	\bar{p} production cross section	20-280	$5 \cdot 10^5$	25	p	LH2, LHe	2022 1 month	LHe target
\bar{p} -induced Spectroscopy	Heavy quark exotics	12, 20	$5 \cdot 10^7$	25	\bar{p}	LH2	2022 2 years	target spectr.: tracking, calorimetry
Drell-Yan	Pion PDFs	190	$7 \cdot 10^7$	25	π^\pm	C/W	2022 1-2 years	
Drell-Yan (RF)	Kaon PDFs & Nucleon TMDs	~ 100	10^8	25-50	K^\pm, \bar{p}	NH_3^\dagger , C/W	2026 2-3 years	"active absorber", vertex det.
Primakoff (RF)	Kaon polarisability & pion life time	~ 100	$5 \cdot 10^6$	> 10	K^-	Ni	non-exclusive 2026 1 year	
Prompt Photons (RF)	Meson gluon PDFs	≥ 100	$5 \cdot 10^6$	10-100	K^\pm π^\pm	LH2, Ni	non-exclusive 2026 1-2 years	hodoscope
K -induced Spectroscopy (RF)	High-precision strange-meson spectrum	50-100	$5 \cdot 10^6$	25	K^-	LH2	2026 1 year	recoil TOF, forward PID
Vector mesons (RF)	Spin Density Matrix Elements	50-100	$5 \cdot 10^6$	10-100	K^\pm, π^\pm	from H to Pb	2026 1 year	

Possible RPD for COMPASS⁺⁺ / AMBER



A recoil proton detector (RPD) is mandatory to ensure the exclusivity. A Silicon detector is included *between* the target surrounded by the modified MW cavity *and* the polarizing magnet



A technology developed at JINR for NICA for the BM@N experiment

No possibility for ToF → PID of p/π with dE/dx
Momentum and trajectory measurements
 $|t|_{\min} \sim 0.1 \text{ GeV}$

ϕ Dep. of BH+DVCS with Unpol Target



$$\frac{d^4\sigma(\ell p \rightarrow \ell p \gamma)}{dx_B dQ^2 d|t| d\phi} = \underset{\text{Well known}}{d\sigma^{BH}} + \left(d\sigma_{unpol}^{DVCS} + P_\ell d\sigma_{pol}^{DVCS} \right) + (e_\ell \text{Re } I + e_\ell P_\ell \text{Im } I)$$

Σ	Σ
Σ	Σ
Δ	Δ
Σ	Δ
Δ	Σ
$\uparrow\downarrow$	$\uparrow\downarrow$
e^-	μ^\pm


$$d\sigma^{BH} \propto c_0^{BH} + c_1^{BH} \cos \phi + c_2^{BH} \cos 2\phi$$

$$d\sigma_{unpol}^{DVCS} \propto c_0^{DVCS} + c_1^{DVCS} \cos \phi + c_2^{DVCS} \cos 2\phi$$




$$d\sigma_{pol}^{DVCS} \propto s_1^{DVCS} \sin \phi$$

$$\text{Re } I \propto c_0^I + c_1^I \cos \phi + c_2^I \cos 2\phi + c_3^I \cos 3\phi$$

$$\text{Im } I \propto s_1^I \sin \phi + s_2^I \sin 2\phi$$



Twist-2 >>
 ■ Twist-3,
 ■ Twist-2
 double helicity flip
 for gluons (NLO)



$$s_1^I = \text{Im } \mathcal{F} \quad c_1^I = \text{Re } \mathcal{F} \quad \mathcal{F} = F_1 \mathcal{H} + \xi (F_1 + F_2) \tilde{\mathcal{H}} - t/4m^2 F_2 \mathcal{E} \xrightarrow{\text{at small } x_B} F_1 \mathcal{H}$$

for proton

$$c_0^{DVCS} \propto 4(\mathcal{H}\mathcal{H}^* + \tilde{\mathcal{H}}\tilde{\mathcal{H}}^*) + \frac{t}{M^2} \mathcal{E}\mathcal{E}^* \xrightarrow{\text{at small } x_B} 4 (\text{Im } \mathcal{H})^2$$




Beam Charge-spin Difference



$$D_{CS,U}(\phi) \equiv d\sigma(\mu^{+\leftarrow}) - d\sigma(\mu^{-\rightarrow}) \rightarrow c_0^I + c_1^I \cos \phi$$




$$c_1^I = \text{Re } F_1 \mathcal{H}$$

— KM10
— VGG



➤ With $\text{Re } F_1 \mathcal{H}$ and $\text{Im } F_1 \mathcal{H}$
 → Extraction of **D-term**

$\text{Re } \mathcal{H} > 0$ at H1
 < 0 at HERMES
 Value of x_{Bj} for the node?



← HERMES ← JLab
← COMPASS 2 years of data $E_\mu = 160 \text{ GeV}$ $1 < Q^2 < 8 \text{ GeV}^2$

Exclusive π^0 Production on Unpolarized Proton



$e p \rightarrow e \pi^0 p$

$$\frac{d^2\sigma}{dt d\phi_\pi} = \frac{1}{2\pi} \left[\left(\frac{d\sigma_T}{dt} + \epsilon \frac{d\sigma_L}{dt} \right) + \epsilon \cos 2\phi_\pi \frac{d\sigma_{TT}}{dt} + \sqrt{2\epsilon(1+\epsilon)} \cos \phi_\pi \frac{d\sigma_{LT}}{dt} \right]$$

$$\frac{d\sigma_L}{dt} = \frac{4\pi\alpha}{k'} \frac{1}{Q^6} \left\{ (1-\xi^2) |\langle \tilde{H} \rangle|^2 - 2\xi^2 \text{Re} [\langle \tilde{H} \rangle^* \langle \tilde{E} \rangle] - \frac{t'}{4m^2} \xi^2 |\langle \tilde{E} \rangle|^2 \right\}$$

Leading twist expected to be dominant
But measured as \approx only a few % of $\frac{d\sigma_T}{dt}$

The other contributions arise from coupling between chiral-odd (quark helicity flip) GPDs to the **twist-3** pion amplitude

$$\frac{d\sigma_T}{dt} = \frac{4\pi\alpha}{2k'} \frac{\mu_\pi^2}{Q^8} \left[(1-\xi^2) |\langle H_T \rangle|^2 - \frac{t'}{8m^2} |\langle \bar{E}_T \rangle|^2 \right]$$


$$\frac{d\sigma_{LT}}{dt} = \frac{4\pi\alpha}{\sqrt{2}k'} \frac{\mu_\pi}{Q^7} \xi \sqrt{1-\xi^2} \frac{\sqrt{-t'}}{2m} \text{Re} [\langle H_T \rangle^* \langle \tilde{E} \rangle]$$

$$\frac{d\sigma_{TT}}{dt} = \frac{4\pi\alpha}{k'} \frac{\mu_\pi^2}{Q^8} \frac{t'}{16m^2} |\langle \bar{E}_T \rangle|^2$$

A large contribution of \bar{E}_T can be identified:

- σ_{TT} contribution
- The dip at small $|t|$ of σ_T

COMPASS Acceptance of ϕ for DVCS



$$\frac{d^3\sigma_T^{\mu p}}{dQ^2 d\nu dt} = \int_{-\pi}^{\pi} d\phi (d\sigma - d\sigma^{BH}) \propto c_0^{DVCS}$$

Acceptance

- μ^- beam
- μ^+ beam

$$\left[-\pi, -\frac{4}{5}\pi, \dots, \frac{4}{5}\pi, \pi \right]$$

$\phi_{\gamma^* \gamma}$ (rad)

$$\frac{d\sigma^{\gamma^* p}}{dt} = \frac{1}{\Gamma(Q^2, \nu, E_\mu)} \frac{d^3\sigma_T^{\mu p}}{dQ^2 d\nu dt}$$

Flux of transverse virtual photons