

Exclusive π^0 muoproduction at COMPASS

Markéta Pešková (Charles University, Prague)

on behalf of the COMPASS collaboration



SPIN'21, October 18. - 22., 2021



Generalised Parton Distributions

- Proton spin sum rule: $\frac{1}{2} = \frac{1}{2}\Delta\Sigma + \Delta G + L_q + L_g$

Jaffe&Manohar Nucl. Phys. B337 (1990)



COMPASS experiment in μp DIS: $\Delta\Sigma = 0.32 \pm 0.03$

COMPASS Collaboration: Phys. Lett. B 693 (2010)

COMPASS, RHIC results: $\Delta G = 0.2^{+0.06}_{-0.07}$

de Florian et al. Phys.Rev.Lett. 113 (2014) no.1, 012001


Missing component: $L_{q,g} = ?? \rightarrow$ GPDs provides access




- Generalised Parton Distributions (GPD) give access to the 3D structure of a hadron
- GPDs encode the correlation between the longitudinal momentum of a parton and its position in the transverse plane

$$q^f(x, b_{\perp}) \xrightarrow{\int dx} \text{Form factors}$$

$$q^f(x, b_{\perp}) \xrightarrow{\int db_{\perp}} \text{PDFs}$$



Generalised Parton Distributions




- Most commonly used processes for GPDs parametrisation are Deeply Virtual Compton Scattering (DVCS) and Hard Exclusive Meson Production (HEMP)
- DVCS gives access to GPD $H \rightarrow$ 3D imaging of a hadron


$$H^q(x, \xi = 0, t) = \rho^q(x, b_\perp) \quad (\text{Burkardt 2000, 2003})$$

Definition of variables:

$q \dots \gamma^*$ four-momentum
 $x \dots$ average longitudinal momentum fraction of initial and final parton (NOT accessible)
 $\xi \dots$ difference of longitudinal-momentum fraction between initial and final parton
 $\approx x_B / (2 - x_B)$
 $t \dots$ four-momentum transfer



Generalised Parton Distributions




Definition of variables:

- $q \dots \gamma^*$ four-momentum
- $x \dots$ average longitudinal momentum fraction of initial and final parton (NOT accessible)
- $\xi \dots$ difference of longitudinal-momentum fraction between initial and final parton $\approx x_B / (2 - x_B)$
- $t \dots$ four-momentum transfer


- Most commonly used processes for GPDs parametrisation are Deeply Virtual Compton Scattering (DVCS) and Hard Exclusive Meson Production (HEMP)
- Vector meson production gives access to GPD $E \rightarrow$ helps constraining the total angular momentum of partons

$$J^f = \frac{1}{2} \lim_{t \rightarrow 0} \int_{-1}^1 dx x [H^f(x, \xi, t) + E^f(x, \xi, t)]$$

Phys. Rev. Lett. 78 (1997)



Generalised Parton Distributions




- 4 chiral-even GPDs (parton helicity conserved)
- 4 chiral-odd (or transversity) GPDs (parton helicity flipped)

		Quark Polarisation		
		Unpolarised (U)	Longitudinally polarised (L)	Transversely polarised (T)
Nucleon Polarisation	U	H		\bar{E}_T
	L		\tilde{H}	\tilde{E}_T
	T	E	\tilde{E}	H_T, \tilde{H}_T

GPDs enter the exclusive processes through **Compton Form Factors (CFF)**

$$\mathcal{H}(\xi, t) = \int_{-1}^1 dx \frac{H^q(x, \xi, t)}{x - \xi + i\epsilon} = \mathcal{P} \int_{-1}^1 dx \frac{H(x, \xi, t)}{x - \xi} + i\pi H$$

Generalised Parton Distributions



- 4 chiral-even GPDs (parton helicity conserved)
- 4 chiral-odd (or transversity) GPDs (parton helicity flipped)


		Quark Polarisation		
		Unpolarised (U)	Longitudinally polarised (L)	Transversely polarised (T)
Nucleon Polarisation	U	H		\bar{E}_T
	L		\tilde{H}	\tilde{E}_T
	T	E	\tilde{E}	H_T, \tilde{H}_T

GPDs enter the exclusive processes through **Compton Form Factors (CFF)**


$$\mathcal{H}(\xi, t) = \int_{-1}^1 dx \frac{H^q(x, \xi, t)}{x - \xi + i\epsilon} = \mathcal{P} \int_{-1}^1 dx \frac{H(x, \xi, t)}{x - \xi} + i\pi H$$

Generalised Parton Distributions

Quark contribution



Gluon contribution



Hard Exclusive Meson Production:

- Flavour separation for specific GPDs due to different partonic content of mesons
- Gluon and quark contributions at the same order in α_s for vector mesons
- DVCS sensitive to H^f , E^f , \tilde{H}^f , and \tilde{E}^f
- At the leading twist:
 - Vector meson production sensitive to H^f , and E^f
 - Pseudoscalar mesons production is described by GPDs \tilde{H}^f , and \tilde{E}^f
- Both vector meson and pseudoscalar mesons (as the π_0 presented in this talk) are also sensitive to $\tilde{E}_T^f = 2\tilde{H}_T^f + E_T^f$, and H_T^f

Road to HEMP cross-section


COMPASS measurement in 2012, and 2016/17 with μ^+ and μ^- beams of $E_\mu = 160$ GeV

Collected events corrected for:

- Luminosity of μ^+ and μ^- beams
- Background subtraction
- Acceptance of the spectrometer
- Reduction of μp cross-section to $\gamma^* p$:

$$\frac{d^4\sigma_{\mu p}}{dQ^2 dt d\nu d\phi} = \Gamma \frac{d^2\sigma_{\gamma^* p}}{dt d\phi}$$

with the virtual photon flux $\Gamma = \Gamma(E_\mu, Q^2, \nu)$



COMPASS 2012:

- 4 weeks \rightarrow results published:
PLB 805(2020) 135454

COMPASS 2016/17:

- 2 \times 6 months

Road to HEMP cross-section


COMPASS measurement in 2012, and 2016/17 with μ^+ and μ^- beams of $E_\mu = 160$ GeV

Collected events corrected for:

- Luminosity of μ^+ and μ^- beams
- **Background subtraction**
- Acceptance of the spectrometer
- Reduction of μp cross-section to $\gamma^* p$:

$$\frac{d^4\sigma_{\mu p}}{dQ^2 dt d\nu d\phi} = \Gamma \frac{d^2\sigma_{\gamma^* p}}{dt d\phi}$$

with the virtual photon flux $\Gamma = \Gamma(E_\mu, Q^2, \nu)$



COMPASS 2012:

- 4 weeks \rightarrow **results published:**
PLB 805(2020) 135454

COMPASS 2016/17:

- 2×6 months

Road to HEMP cross-section


COMPASS measurement in 2012, and 2016/17 with μ^+ and μ^- beams of $E_\mu = 160$ GeV

Collected events corrected for:

- Luminosity of μ^+ and μ^- beams
- Background subtraction
- **Acceptance of the spectrometer**
- Reduction of μp cross-section to $\gamma^* p$:

$$\frac{d^4\sigma_{\mu p}}{dQ^2 dt d\nu d\phi} = \Gamma \frac{d^2\sigma_{\gamma^* p}}{dt d\phi}$$

with the virtual photon flux $\Gamma = \Gamma(E_\mu, Q^2, \nu)$



COMPASS 2012:

- 4 weeks \rightarrow **results published:**
PLB 805(2020) 135454

COMPASS 2016/17:

- 2 \times 6 months

HEMP cross section


HEMP cross-section, reduced to γ^*p , for the **unpolarised target** and **polarised lepton beam** (relevant for COMPASS 2012, 2016/2017 measurements):

$$\frac{d^2\sigma_{\gamma^*p}^{\leftrightarrow}}{dt d\phi} = \frac{1}{2\pi} \left[\frac{d\sigma_T}{dt} + \epsilon \frac{d\sigma_L}{dt} + \epsilon \cos(2\phi) \frac{d\sigma_{TT}}{dt} + \sqrt{\epsilon(1+\epsilon)} \cos\phi \frac{d\sigma_{LT}}{dt} \right]$$

$$\mp |P_l| \sqrt{\epsilon(1-\epsilon)} \sin\phi \frac{d\sigma'_{LT}}{dt}$$

$$\epsilon = \frac{1 - y - \frac{y^2\gamma^2}{4}}{1 - y + \frac{y^2}{2} + \frac{y^2\gamma^2}{4}}$$

Factorization proven for σ_L ,
not for σ_T which is expected to
be suppressed by a factor $1/Q^2$
BUT large contributions are observed
at JLab



HEMP cross section

Spin independent HEMP cross-section after averaging the two spin-dependent cross-sections:

$$\frac{d^2\sigma_{\gamma^*p}}{dt d\phi} = \frac{1}{2} \left(\frac{d^2\sigma_{\gamma^*p}^{\leftarrow}}{dt d\phi} + \frac{d^2\sigma_{\gamma^*p}^{\rightarrow}}{dt d\phi} \right) =$$


$$\frac{1}{2\pi} \left[\frac{d\sigma_T}{dt} + \epsilon \frac{d\sigma_L}{dt} + \epsilon \cos(2\phi) \frac{d\sigma_{TT}}{dt} + \sqrt{\epsilon(1+\epsilon)} \cos\phi \frac{d\sigma_{LT}}{dt} \right]$$

~~$$\mp |P_I| \sqrt{\epsilon(1-\epsilon)} \sin\phi \frac{d\sigma'_{LT}}{dt}$$~~

$$\epsilon = \frac{1 - y - \frac{y^2\gamma^2}{4}}{1 - y + \frac{y^2}{2} + \frac{y^2\gamma^2}{4}}$$

⇒ study ϕ
dependence

Factorization proven for σ_L ,
not for σ_T which is expected to
be suppressed by a factor $1/Q^2$
BUT large contributions are observed
at JLab



HEMP cross section

Spin independent HEMP cross-section after averaging the two spin-dependent cross-sections:

$$\frac{d^2\sigma_{\gamma^*p}}{dt d\phi} = \frac{1}{2} \left(\frac{d^2\sigma_{\gamma^*p}^{\leftarrow}}{dt d\phi} + \frac{d^2\sigma_{\gamma^*p}^{\rightarrow}}{dt d\phi} \right) =$$

$$\frac{1}{2\pi} \left[\frac{d\sigma_T}{dt} + \epsilon \frac{d\sigma_L}{dt} + \epsilon \cos(2\phi) \frac{d\sigma_{TT}}{dt} + \sqrt{\epsilon(1+\epsilon)} \cos\phi \frac{d\sigma_{LT}}{dt} \right]$$


~~$$\mp |P_I| \sqrt{\epsilon(1-\epsilon)} \sin\phi \frac{d\sigma'_{LT}}{dt}$$~~

\Rightarrow study ϕ dependence

After integration in ϕ :

$$\frac{d\sigma_T}{dt} + \epsilon \frac{d\sigma_L}{dt}$$

\Rightarrow study t dependence



HEMP cross section

$$\frac{d^2\sigma_{\gamma^*p}}{dt d\phi} = \frac{1}{2\pi} \left[\frac{d\sigma_T}{dt} + \epsilon \frac{d\sigma_L}{dt} + \epsilon \cos(2\phi) \frac{d\sigma_{TT}}{dt} + \sqrt{\epsilon(1+\epsilon)} \cos\phi \frac{d\sigma_{LT}}{dt} \right]$$


GPDs in exclusive π^0 production

$$\frac{d\sigma_L}{dt} \propto \left[(1-\xi^2) |\langle \tilde{\mathcal{H}} \rangle|^2 - 2\xi^2 \text{Re}(\langle \tilde{\mathcal{H}} \rangle^* \langle \tilde{\mathcal{E}} \rangle) - \frac{t'}{4M^2} \xi^2 |\langle \tilde{\mathcal{E}} \rangle|^2 \right]$$

$$\frac{d\sigma_T}{dt} \propto \left[(1-\xi^2) |\langle \mathcal{H}_T \rangle|^2 - \frac{t'}{8M^2} |\langle \bar{\mathcal{E}}_T \rangle|^2 \right]$$

$$\frac{d\sigma_{TT}}{dt} \propto t' |\langle \bar{\mathcal{E}}_T \rangle|^2$$


$$\frac{d\sigma_{LT}}{dt} \propto \xi \sqrt{1-\xi^2} \sqrt{-t'} \text{Re}(\langle \mathcal{H}_T \rangle^* \langle \tilde{\mathcal{E}} \rangle)$$



Impact of $\bar{\mathcal{E}}_T$ should be visible in $\frac{d\sigma_{TT}}{dt}$,
and also a dip at small t of $\frac{d\sigma_T}{dt}$


$t' = t - t_{min}$, t_{min} smallest possible momentum transfer

COMPASS: Versatile facility to study QCD
with hadron (π^\pm , K^\pm , p ...) and lepton (polarized μ^\pm) beams
of ~ 200 GeV for hadron spectroscopy and
hadron structure studies using SIDIS, DY, DVCS, DVMP...



COMPASS GPD program

- Two stage magnetic spectrometer with large angular and momentum acceptance
- Versatile usage: hadron and muon beams
- Particle identification:
 - Ring Imaging Cherenkov (RICH) detector
 - Electromagnetic calorimeters (ECAL0, ECAL1, ECAL2)
 - Hadronic calorimeters (HCAL1, HCAL2)
 - 2 muon walls






COMPASS GPD program

- Target ToF system:
 - 24 inner and outer scintillators
 - 1 GHz readout
 - 310 ps ToF resolution
- ECAL0 calorimeter:
 - shaslyk modules
 - 2×2 m, 2200 channels


New equipments:

- 2.5m LH2 target
- 4m ToF Barrel CAMERA
- ECAL0



Exclusive π^0 production: Selection

- Incoming and outgoing μ connected to primary vertex
- Two photons in ECALs from π^0 decay, attached to the vertex
- Recoil proton candidate
- $1 < Q^2 < 5 \text{ (GeV/c)}^2$,
 $8.5 < \nu < 28 \text{ GeV}$,
 $0.08 < |t| < 0.64 \text{ (GeV/c)}^2$



Selections for exclusive π^0 events:

- Transverse momentum constraint:

$$\Delta p_T = p_{T,spect}^p - p_{T,recoil}^p$$

- $\Delta\varphi = \varphi_{spect}^p - \varphi_{recoil}^p$

- Z coordinate of inner CAMERA ring:

$$\Delta z = z_{spect}^p - z_{recoil}^p$$



- Energy-momentum conservation:

Exclusive π^0 production: Selection

- Incoming and outgoing μ connected to primary vertex
- Two photons in ECALs from π^0 decay, attached to the vertex
- Recoil proton candidate
- $1 < Q^2 < 5 \text{ (GeV/c)}^2$,
 $8.5 < \nu < 28 \text{ GeV}$,
 $0.08 < |t| < 0.64 \text{ (GeV/c)}^2$

Selections for exclusive π^0 events:

- Transverse momentum constraint:
 $\Delta p_T = p_{T,spect}^p - p_{T,recoil}^p$
- $\Delta\varphi = \varphi_{spect}^p - \varphi_{recoil}^p$
- Z coordinate of inner CAMERA ring:
 $\Delta z = z_{spect}^p - z_{recoil}^p$
- Energy-momentum conservation:
 $M_X^2 = (p_{\mu,in} + p_p - p_{\mu,out} - p_{p'} - p_{\pi^0})^2$
- Invariant mass $M_{\gamma\gamma}$ cut





Exclusive π^0 production: Selection

- Incoming and outgoing μ connected to primary vertex
- Two photons in ECALs from π^0 decay, attached to the vertex
- Recoil proton candidate
- $1 < Q^2 < 5 \text{ (GeV/c)}^2$,
 $8.5 < \nu < 28 \text{ GeV}$,
 $0.08 < |t| < 0.64 \text{ (GeV/c)}^2$

Selections for exclusive π^0 events:

- Transverse momentum constraint:
$$\Delta p_T = p_{T,spect}^p - p_{T,recoil}^p$$
- $\Delta\varphi = \varphi_{spect}^p - \varphi_{recoil}^p$
- Z coordinate of inner CAMERA ring:
$$\Delta z = z_{spect}^p - z_{recoil}^p$$
- Energy-momentum conservation:
$$M_X^2 = (p_{\mu,in} + p_p - p_{\mu,out} - p_{p'} - p_{\pi^0})^2$$
- Invariant mass $M_{\gamma\gamma}$ cut





Exclusive π^0 production: Selection

- Incoming and outgoing μ connected to primary vertex
- Two photons in ECALs from π^0 decay, attached to the vertex
- Recoil proton candidate
- $1 < Q^2 < 5 \text{ (GeV/c)}^2$,
 $8.5 < \nu < 28 \text{ GeV}$,
 $0.08 < |t| < 0.64 \text{ (GeV/c)}^2$

Selections for exclusive π^0 events:

- Transverse momentum constraint:
 $\Delta p_T = p_{T,spect}^p - p_{T,recoil}^p$
- $\Delta\varphi = \varphi_{spect}^p - \varphi_{recoil}^p$
- Z coordinate of inner CAMERA ring:
 $\Delta z = z_{spect}^p - z_{recoil}^p$
- Energy-momentum conservation:
 $M_X^2 = (p_{\mu,in} + p_p - p_{\mu,out} - p_{p'} - p_{\pi^0})^2$
- Invariant mass $M_{\gamma\gamma}$ cut





Exclusive π^0 production: Selection

- Incoming and outgoing μ connected to primary vertex
- Two photons in ECALs from π^0 decay, attached to the vertex
- Recoil proton candidate
- $1 < Q^2 < 5 \text{ (GeV/c)}^2$,
 $8.5 < \nu < 28 \text{ GeV}$,
 $0.08 < |t| < 0.64 \text{ (GeV/c)}^2$

Selections for exclusive π^0 events:

- Transverse momentum constraint:
$$\Delta p_T = p_{T,spect}^p - p_{T,recoil}^p$$
- $\Delta\varphi = \varphi_{spect}^p - \varphi_{recoil}^p$
- Z coordinate of inner CAMERA ring:
$$\Delta z = z_{spect}^p - z_{recoil}^p$$
- Energy-momentum conservation:
$$M_X^2 = (p_{\mu,in} + p_p - p_{\mu,out} - p_{p'} - p_{\pi^0})^2$$
- Invariant mass $M_{\gamma\gamma}$ cut





Exclusive π^0 production: Selection

- Incoming and outgoing μ connected to primary vertex
- Two photons in ECALs from π^0 decay, attached to the vertex
- Recoil proton candidate
- $1 < Q^2 < 5 \text{ (GeV/c)}^2$,
 $8.5 < \nu < 28 \text{ GeV}$,
 $0.08 < |t| < 0.64 \text{ (GeV/c)}^2$



Selections for exclusive π^0 events:

- Transverse momentum constraint:
 $\Delta p_T = p_{T,spect}^p - p_{T,recoil}^p$
- $\Delta\varphi = \varphi_{spect}^p - \varphi_{recoil}^p$
- Z coordinate of inner CAMERA ring:
 $\Delta z = z_{spect}^p - z_{recoil}^p$
- Energy-momentum conservation:
 $M_X^2 = (p_{\mu,in} + p_p - p_{\mu,out} - p_{p'} - p_{\pi^0})^2$
- Invariant mass $M_{\gamma\gamma}$ cut



Exclusive π^0 production: SIDIS background estimation

- Main background of π^0 production \Rightarrow non-exclusive DIS processes
- 2 reference samples (wider kinematic range) described by MC:
 - LEPTO for describing the shape of non-exclusive background distribution
 - HEPGEN++ for the shape of distributions of exclusive π^0 production (signal contribution)
- Search for best description of data in **signal region** and **background region**




Resulting fraction of non-exclusive background in data:


$$(29_{-6}^{+2} |_{\text{sys}}) \%$$

Exclusive π^0 production: COMPASS acceptance

- 4D acceptance in bins of ϕ_{π^0} , ν , $|t|$, Q^2
- figure shows 3D projection, as a function of ϕ_{π^0}



Exclusive π^0 cross-section as a function of $|t|$




PLB 805 (2020) 135454

Exclusive π^0 cross-section as a function of $|t|$


Dip would indicate contribution of \bar{E}_T

Factor of ~ 2 discrepancy with Goloskokov&Kroll model [EPJ A47 \(2011\) 112](#)




Exclusive π^0 cross-section as a function of ϕ

$$\frac{d^2\sigma_{\gamma^*p}}{dt d\phi} = \frac{1}{2\pi} \left[\frac{d\sigma_T}{dt} + \epsilon \frac{d\sigma_L}{dt} + \epsilon \cos(2\phi) \frac{d\sigma_{TT}}{dt} + \sqrt{\epsilon(1+\epsilon)} \cos\phi \frac{d\sigma_{LT}}{dt} \right]$$





Exclusive π^0 cross-section as a function of ϕ

$$\frac{d^2\sigma_{\gamma^*p}}{dt d\phi} = \frac{1}{2\pi} \left[\frac{d\sigma_T}{dt} + \epsilon \frac{d\sigma_L}{dt} + \epsilon \cos(2\phi) \frac{d\sigma_{TT}}{dt} + \sqrt{\epsilon(1+\epsilon)} \cos\phi \frac{d\sigma_{LT}}{dt} \right]$$





Outlook to COMPASS 2016/17 results

- Integrated 2012 luminosity: $L_{\mu^+} = 18.9 \text{ pb}^{-1}$, μ^- : $L_{\mu^-} = 23.5 \text{ pb}^{-1}$
- 2016+2017 data: $\sim 9\times$ statistics of 2012



Outlook to COMPASS 2016/17 results


- Integrated 2012 luminosity: $L_{\mu^+} = 18.9 \text{ pb}^{-1}$, μ^- : $L_{\mu^-} = 23.5 \text{ pb}^{-1}$
- 2016+2017 data: $\sim 9\times$ statistics of 2012
- 2016/7 data: Flatter acceptance, wider in large photon angles
- Analysis of 2016/17 data is ongoing, currently 2/3 of 2016 data processed
→ $2.5\times$ larger statistics than 2012



Summary

t -dependence and ϕ -dependence of exclusive π^0 cross-section on unpolarised proton target:

- First results at low ξ (or $\langle x_B \rangle = 0.093$) from COMPASS 2012 pilot measurement, input for constraining the Goloskokov&Kroll model



PLB 805 (2020) 135454

- New results soon expected from the measurement in 2016/2017 for DVCS, vector and pseudoscalar meson production
- Collected 2016/2017 statistics $\sim 9 \times$ larger than from 2012 run
- New results coming soon on differential cross-section of exclusive π^0 as a function Q^2 , ν , t , and ϕ !



Thank you for your attention!