


COMPASS Spin Physics Program



BAKUR PARSAMYAN

AANL, INFN section of Turin and CERN
on behalf of the COMPASS Collaboration



The 25th International Spin Symposium (SPIN 2023)
September 24-30, Duke University, Durham Convention Center, US

COMPASS collaboration

Common Muon and Proton Apparatus for Structure and Spectroscopy




25 institutions from 13 countries
– nearly 200 physicists (in 2022)

- CERN SPS north area
- Fixed target experiment
- Approved in 1997 (**25 years**)
- Taking data since 2002 (**20 years**)

International Workshop on Hadron Structure and Spectroscopy
IWHSS-2022 workshop (**anniversary edition**)
CERN Globe, August 29-31, 2022



<https://indico.cern.ch/e/IWHSS-2022>



COMPASS collaboration



Common Muon and Proton Apparatus for Structure and Spectroscopy



28 institutions from 14 countries

– nearly 210 physicists (in 2023: start of the Analysis Phase)

- CERN SPS north area
- Fixed target experiment
- Approved in 1997 (25 years)
- Taking data since 2002 (20 years)

Wide physics program


COMPASS-I

- Data taking 2002-2011
- Muon and hadron beams
- Nucleon spin structure
- Spectroscopy

COMPASS-II

- Data taking 2012-2022
- Primakoff
- DVCS (GPD+SIDIS)
- Polarized Drell-Yan
- Transverse deuteron SIDIS 2022

3 new groups joined the COMPASS collaboration in 2023
UCon (US), AANL (Armenia), NCU (Taiwan)



COMPASS web page: <http://wwwcompass.cern.ch>

See talks by: V. Andrieux, A. Kerbizi, A. Martin, J. Matousek, G. Reicherz, A. Vijayakumar

The COMPASS Experiment at the CERN SPS





Broad Physics Program to study Structure and Excitation Spectrum of Hadrons

Increasing resolution scale
(momentum transfer)


Nucleon structure

- Hard scattering of μ^\pm and π^- off (un)polarized P/D targets
- Study of nucleon spin structure
- Parton distribution functions and fragmentation functions




Hadron spectroscopy

- Diffractive $\pi(K)$ dissociation reaction with proton target
- PWA technique employed
- High-precision measurement of light-meson excitation spectrum
- Search for exotic states



Chiral dynamics

- Test chiral perturbation theory in $\pi(K)\gamma$ reactions
- π^\pm and K^\pm polarizabilities
- Chiral anomaly $F_{3\pi}$



The COMPASS Experiment at the CERN SPS





Broad Physics Program to study Structure and Excitation Spectrum of Hadrons

Increasing resolution scale
(momentum transfer)

Nucleon structure

- Hard scattering of μ^\pm and π^- off (un)polarized P/D targets
- Study of nucleon spin structure
- Parton distribution functions and fragmentation functions




Hadron spectroscopy

- Diffractive $\pi(K)$ dissociation reaction with proton target
- PWA technique employed
- High-precision measurement of light-meson excitation spectrum
- Search for exotic states


Chiral dynamics

- Test chiral perturbation theory in $\pi(K)\gamma$ reactions
- π^\pm and K^\pm polarizabilities
- Chiral anomaly $F_{3\pi}$





Nucleon 3D structure

- Transverse position \vec{b}_T of partons
 - Correlation between \vec{b}_T and x
 - Complementary to TMD PDFs
- 8 generalized parton distribution functions (GPDs)
 - Contain information about parton orbital angular momentum
 - Mostly unknown
- Measured in exclusive processes:
 - Deeply virtual Compton scattering (DVCS): $\mu + N \rightarrow \mu + \gamma + N$
 - Hard exclusive meson production (HEMP): $\mu + N \rightarrow \mu + VM + N$ with $VM = \pi^0, \rho(770), \omega(782), \dots$



See the COMPASS GPD program overview talk by J. Matousek



COMPASS experimental setup




COmmon Muon Proton Apparatus for Structure and Spectroscopy

CERN SPS North Area (building 888)


Two-stage spectrometer LAS+SAS

- Large Angle Spectrometer (SM1 magnet)
- Small Angle Spectrometer (SM2 magnet)



- Primary beam - 400 GeV p from SPS
 - impinging on Be production target (T6)
- 190 GeV secondary hadron beams
 - h^- beam: 97% π^- , 2% K^- , 1% p
 - h^+ beam: 75% π^+ , 24% p , 1% K^+
- 160 GeV tertiary muon beams
 - μ^\pm longitudinally polarized

B. Parsamyan




Large-acceptance forward spectrometer

- Precise tracking (350 planes)
SciFi, Silicon, MicroMegas, GEM, MWPC, DC, Straw, Muon walls
- PID - CEDARs, RICH, calorimeters, MWs

Various targets:

- Polarized solid-state NH_3 or 6LiD
- Liquid H_2
- Solid-state nuclear targets (e.g. Ni, W, Pb)



COMPASS experimental setup: Phase II (SIDIS programme)




COmmon Muon Proton Apparatus for Structure and Spectroscopy




CERN SPS North Area (building 888)

Two-stage spectrometer LAS+SAS

- Large Angle Spectrometer (SM1 magnet)
- Small Angle Spectrometer (SM2 magnet)



- Primary beam - 400 GeV p from SPS
 - impinging on Be production target (T6)
- 190 GeV secondary hadron beams
 - h^- beam: 97% π^- , 2% K^- , 1% p
 - h^+ beam: 75% π^+ , 24% p , 1% K^+
- 160 GeV tertiary muon beams
 - μ^+ longitudinally polarized



see Gerhard Reicherz' talk

COMPASS experimental setup: Phase II (DY programme)




COmmon Muon Proton Apparatus for Structure and Spectroscopy





CERN SPS North Area (building 888)

Two-stage spectrometer LAS+SAS

- Large Angle Spectrometer (SM1 magnet)
- Small Angle Spectrometer (SM2 magnet)



- Primary beam - 400 GeV p from SPS
 - impinging on Be production target (T6)
- 190 GeV secondary hadron beams
 - h^- beam: 97% π^- , 2% K^- , 1% p
 - h^+ beam: 75% π^+ , 24% p , 1% K^+
- 160 GeV tertiary muon beams
 - μ^\pm longitudinally polarized



COMPASS data taking campaigns



Beam	Target	year	Physics programme
μ^+	Polarized deuteron (${}^6\text{LiD}$)	2002	
		2003	80% Longitudinal 20% Transverse SIDIS
		2004	
		2006	Longitudinal SIDIS
	Polarized proton (NH_3)	2007	50% Longitudinal 50% Transverse SIDIS
$\pi K p$	$\text{LH}_2, \text{Ni}, \text{Pb}, \text{W}$	2008 2009	Spectroscopy
μ^+	Polarized proton (NH_3)	2010	Transverse SIDIS
		2011	Longitudinal SIDIS
$\pi K p$	Ni	2012	Primakoff
μ^\pm	LH_2	2012	Pilot DVCS & HEMP & unpolarized SIDIS
π^-	Polarized proton (NH_3)	2014	Pilot Drell-Yan
		2015	Transverse Drell-Yan
		2018	
μ^\pm	LH_2	2016 2017	DVCS & HEMP & unpolarized SIDIS
μ^+	Polarized deuteron (${}^6\text{LiD}$)	2021 2022	Transverse SIDIS

Nucleon spin structure: collinear approach \leftrightarrow TMDs



quark

	U	L	T
U	$f_1^q(x)$ number density		
L		$g_1^q(x)$ helicity	
T			$h_1^q(x)$ transversity



nucleon

↔



	U	L	T
U	$f_1^q(x, \mathbf{k}_T^2)$ number density		$h_1^{\perp q}(x, \mathbf{k}_T^2)$ Boer-Mulders
L		$g_1^q(x, \mathbf{k}_T^2)$ helicity	$h_{1L}^{\perp q}(x, \mathbf{k}_T^2)$ worm-gear L
T	$f_{1T}^{\perp q}(x, \mathbf{k}_T^2)$ Sivers	$g_{1T}^q(x, \mathbf{k}_T^2)$ worm-gear T	$h_1^q(x, \mathbf{k}_T^2)$ transversity $h_{1T}^{\perp q}(x, \mathbf{k}_T^2)$ pretzelosity

nucleon


- PDFs – universal (process independent) objects; T-odd PDFs – conditionally universal




Nucleon spin structure: helicity $g_{1,d}^q(x)$





- COMPASS contribution:
lowest x and highest Q^2 regions





COMPASS PLB 769(2017) 34





Nucleon spin structure: helicity $g_{1,p}^q(x)$




- COMPASS contribution:
lowest x and highest Q^2 regions
- Both deuteron and proton target data



Nucleon spin structure: helicity $g_{1,p}^q(x)$




- COMPASS contribution:
lowest x and highest Q^2 regions
- Both deuteron and proton target data
- For the first time non-zero spin effects at smallest x and Q^2 – positive signal for $g_1^p(x)$





28 September 2023



B. Parsamyan




Nucleon spin structure: helicity $g_{1,d(p)}^q(x)$





- COMPASS contribution:
lowest x and highest Q^2 regions
- Both deuteron and proton target data
- For the first time non-zero spin effects at smallest x and Q^2 – positive signal for $g_1^p(x)$
- Gluon polarization measurements via open charm and SIDIS
- COMPASS - first to rule out a large gluon polarization in the nucleon!
Precise test of Bjorken sum rule (9% level)




Nucleon spin structure: helicity $g_{1,d(p)}^q(x)$



- COMPASS contribution:
lowest x and highest Q^2 regions
- Both deuteron and proton target data
- For the first time non-zero spin effects at
smallest x and Q^2 – positive signal for $g_1^p(x)$
- Both inclusive and semi-inclusive
measurements – access to flavor




COMPASS PLB 693 (2010) 227




Nucleon spin structure



- 1964 Quark model





- 1969 Parton model



- 1973 asymptotic freedom and QCD


- 1978 intrinsic transverse motion of quarks and azimuthal asymmetries



Cahn effect in SIDIS


$$\frac{d\sigma}{dxdydzdp_T^2d\phi_h d\phi_s} = \left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L}) \times (1 + \sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h + \dots)$$

$f_1^q(x, \mathbf{k}_T^2)$
number density





Cahn effect - R. N. Cahn, PLB 78 (1978)

The point that there are azimuthal dependences, which arise from the transverse momenta of the partons was clearly stated in this papers: T.P. Cheng and A. Zee, Phys. Rev. D6 (1972) 885; F. Ravndal, Phys. Lett. 43B (1973) 301.R.L. Kingsley, Phys. Rev. D10 (1974) 1580; A.M. Kotsynyan, Teor. Mat. Fiz. 24 (1975) 206;



See L. Gamberg's talk





Cahn effect in SIDIS

$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_s} = \left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L}) \\ \times (1 + \sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos \phi_h} \cos \phi_h + \dots)$$

$$F_{UU}^{\cos \phi_h} = \frac{2M}{Q} C \left\{ -\frac{\hat{\cdot}}{M_h} \left(\textcolor{red}{xh} H_{1q}^{\perp h} + \frac{M_h}{M} f_1^q \frac{D_q^{\perp h}}{z} \right) - \frac{\hat{\cdot}}{M} \left(\textcolor{red}{xf}^{\perp q} D_{1q}^h + \frac{M_h}{M} h_1^{\perp q} \frac{H_q^h}{z} \right) \right\}$$

Cahn effect in SIDIS

$$\frac{d\sigma}{dxdydzdp_T^2d\phi_hd\phi_s} = \left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L}) \times (1 + \sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h + \dots)$$

Cahn effect




$f_1^q(x, k_T^2)$
number density

As of 1978 – simplistic kinematic effect:

- non-zero k_T induces an azimuthal modulation

As of 2023 – complex SF (twist-2/3 functions)

- Measurements by different experiments
- Complex multi-D kinematic dependences
 - So far, no clear interpretation



Cahn effect in SIDIS

$$\frac{d\sigma}{dxdydzdp_T^2d\phi_hd\phi_s} = \left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L}) \times (1 + \sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h + \dots)$$

Cahn effect




$f_1^q(x, k_T^2)$
number density

As of 1978 – simplistic kinematic effect:

- non-zero k_T induces an azimuthal modulation

As of 2023 – complex SF (twist-2/3 functions)

- Measurements by different experiments
- Complex multi-D kinematic dependences
 - So far, no clear interpretation
- A set of complex corrections:
 - Acceptance, diffractively produced VMs, radiative corrections, etc.



Cahn effect in SIDIS

$$\frac{d\sigma}{dxdydzdp_T^2d\phi_h d\phi_s} = \left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L}) \times (1 + \sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h + \dots)$$



Cahn effect

$f_1^q(x, k_T^2)$
number density


As of 1978 – simplistic kinematic effect:

- non-zero k_T induces an azimuthal modulation

As of 2023 – complex SF (twist-2/3 functions)

- Measurements by different experiments
- Complex multi-D kinematic dependences
 - So far, no clear interpretation
- A set of complex corrections:
 - Acceptance, diffractively produced VMs, radiative corrections, etc.
- Strong Q^2 dependence – unexplained
 - Do not seem to come from RCs
 - Transition between TMD – collinear regions?



Recent COMPASS results



see J. Matousek's talk


Cahn effect in SIDIS

$$\frac{d\sigma}{dxdydzdp_T^2d\phi_hd\phi_s} = \left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L})$$



P_T -dependent distributions

- Extracted in multi-D kinematic bins
- A set of complex corrections:
 - Acceptance, diffractively produced VMs, radiative corrections, etc.
- Global fits by different groups (SIDIS-DY)
 - Normalization issues
(See A. Bacchetta's talk)
- COMPASS measurements
 - isoscalar target data - published
 - proton data – ongoing analysis
- COMPASS-2022 data
 - More deuteron data points to be expected



SIDIS x-section and TMDs at twist-2



$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_s} =$$


All measured by COMPASS

$$\left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L})$$

$$1 + \sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h + \varepsilon A_{UU}^{\cos 2\phi_h} \cos 2\phi_h \\ + \lambda \sqrt{2\varepsilon(1-\varepsilon)} A_{LU}^{\sin\phi_h} \sin\phi_h$$

$$+ S_L \left[\sqrt{2\varepsilon(1+\varepsilon)} A_{UL}^{\sin\phi_h} \sin\phi_h + \varepsilon A_{UL}^{\sin 2\phi_h} \sin 2\phi_h \right] \\ + S_L \lambda \left[\sqrt{1-\varepsilon^2} A_{LL} + \sqrt{2\varepsilon(1-\varepsilon)} A_{LL}^{\cos\phi_h} \cos\phi_h \right]$$

$$\times \left\{ \begin{array}{l} A_{UT}^{\sin(\phi_h - \phi_s)} \sin(\phi_h - \phi_s) \\ + \varepsilon A_{UT}^{\sin(\phi_h + \phi_s)} \sin(\phi_h + \phi_s) \\ + \varepsilon A_{UT}^{\sin(3\phi_h - \phi_s)} \sin(3\phi_h - \phi_s) \\ + \sqrt{2\varepsilon(1+\varepsilon)} A_{UT}^{\sin\phi_s} \sin\phi_s \\ + \sqrt{2\varepsilon(1+\varepsilon)} A_{UT}^{\sin(2\phi_h - \phi_s)} \sin(2\phi_h - \phi_s) \end{array} \right\} \\ + S_T \left\{ \begin{array}{l} \sqrt{(1-\varepsilon^2)} A_{LT}^{\cos(\phi_h - \phi_s)} \cos(\phi_h - \phi_s) \\ + \sqrt{2\varepsilon(1-\varepsilon)} A_{LT}^{\cos\phi_s} \cos\phi_s \\ + \sqrt{2\varepsilon(1-\varepsilon)} A_{LT}^{\cos(2\phi_h - \phi_s)} \cos(2\phi_h - \phi_s) \end{array} \right\}$$



Quark Nucleon	U	L	T
U	number density		Boer-Mulders
L		helicity	worm-gear L
T		Kotzinian-Mulders worm-gear T	transversity pretzelosity
Sivers			

spin of the nucleon spin of the quark k_T

SIDIS: target longitudinal spin dependent asymmetries



$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_s} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots \right.$$

$$\left. + S_L \left[\sqrt{2\varepsilon(1+\varepsilon)} A_{UL}^{\sin \phi_h} \sin \phi_h + \varepsilon A_{UL}^{\sin 2\phi_h} \sin 2\phi_h \right] \right\}$$


$$+ S_L \lambda \left[\sqrt{1-\varepsilon^2} A_{LL} + \sqrt{2\varepsilon(1-\varepsilon)} A_{LL}^{\cos \phi_h} \cos \phi_h \right]$$

$$F_{LL}^1 = \mathcal{C} \left\{ g_{1L}^q D_{1q}^h \right\}$$

$$F_{UL}^{\sin \phi_h} = \frac{2M}{Q} \mathcal{C} \left\{ -\frac{\hat{h} \cdot p_T}{M_h} \left(x h_L^q H_{1q}^{\perp h} + \frac{M_h}{M} g_{1L}^q \frac{\tilde{G}_q^{\perp h}}{z} \right) \right. \\ \left. + \frac{\hat{h} \cdot k_T}{M} \left(x f_L^{\perp q} D_{1q}^h - \frac{M_h}{M} h_{1L}^{\perp q} \frac{\tilde{H}_q^h}{z} \right) \right\}$$

$$F_{UL}^{\sin 2\phi_h} = \mathcal{C} \left\{ -\frac{2(\hat{h} \cdot p_T)(\hat{h} \cdot k_T) - p_T \cdot k_T}{MM_h} h_{1L}^{\perp q} H_{1q}^{\perp h} \right\}$$

$$F_{LL}^{\cos \phi_h} = \frac{2M}{Q} \mathcal{C} \left\{ -\frac{\hat{h} \cdot p_T}{M_h} \left(x e_L^q H_{1q}^{\perp h} + \frac{M_h}{M} g_{1L}^q \frac{\tilde{D}_q^{\perp h}}{z} \right) \right. \\ \left. + \frac{\hat{h} \cdot k_T}{M} \left(x g_L^{\perp q} D_{1q}^h - \frac{M_h}{M} h_{1L}^{\perp q} \frac{\tilde{E}_q^h}{z} \right) \right\}$$



SIDIS: target longitudinal spin dependent asymmetries



$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_s} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots \right.$$

$$\left. + S_L \left[\sqrt{2\varepsilon(1+\varepsilon)} A_{UL}^{\sin\phi_h} \sin\phi_h + \varepsilon A_{UL}^{\sin 2\phi_h} \sin 2\phi_h \right] \right\}$$

$$+ S_L \lambda \left[\sqrt{1-\varepsilon^2} A_{LL} + \sqrt{2\varepsilon(1-\varepsilon)} A_{LL}^{\cos\phi_h} \cos\phi_h \right]$$

COMPASS collected large amount of L-SIDIS data
Unprecedented precision for some amplitudes!

$$A_{UL}^{\sin\phi_h}$$

- Q-suppression, Various different “twist” ingredients
- Sizable TSA-mixing
- Significant h^+ asymmetry, clear z -dependence
- h^- compatible with zero


$$A_{UL}^{\sin 2\phi_h}$$

- Only “twist-2” ingredients
- Additional p_T -suppression
- Compatible with zero, in agreement with models
- Collins-like behavior?

$$A_{LL}^{\cos\phi_h}$$

- Q-suppression, Various different “twist” ingredients
- Compatible with zero, in agreement with models

B. Parsamyan (for COMPASS) [arXiv:1801.01488](https://arxiv.org/abs/1801.01488) [hep-ex]





SIDIS: target longitudinal spin dependent asymmetries



$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_s} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_L \lambda \sqrt{2\varepsilon(1-\varepsilon)} A_{LL}^{\cos\phi_h} \cos\phi_h + \dots \right\}$$

$$F_{LL}^{\cos\phi_h} = \frac{2M}{Q} C \left\{ -\frac{\hat{h} \cdot p_T}{M_h} \left(xe_L^q H_{1q}^{\perp h} + \frac{M_h}{M} g_{1L}^q \frac{\tilde{D}_q^{\perp h}}{z} \right) + \frac{\hat{h} \cdot k_T}{M} \left(x g_L^{\perp q} D_{1q}^h - \frac{M_h}{M} h_{1L}^{\perp q} \frac{\tilde{E}_q^h}{z} \right) \right\}$$




- Q-suppression, various different “twist” ingredients
- **Measured to be non zero at CLAS6, what about CLAS12?**
- HERMES/COMPASS - small and compatible with zero, in agreement with model predictions

Selected results for di-hadron LSAs




COMPASS (NH_3) 2007+2011 data: preliminary




- Alternative way to access various twist-2/-3 distributions
- Non zero signal for $A_{UL}^{sin\phi_R}$ and A_{LL}^1
- CLAS-COMPASS: different behavior for $A_{UL}^{sin2\phi_R}$ at large x?

CLAS 6 GeV (NH_3)
S. A. Pereira: PoS (DIS 2014) 231



$Q^2 > 1 (\text{GeV}/c)^2$
 $0.0025 < x < 0.7$
 $0.1 < y < 0.9$
 $W > 5 \text{ GeV}/c^2$



SIDIS x-section and TMDs at twist-2: TSAs



$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_s} =$$

All measured by COMPASS


$$\left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L})$$

$$\left[1 + \sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h + \varepsilon A_{UU}^{\cos 2\phi_h} \cos 2\phi_h \right. \\ \left. + \lambda \sqrt{2\varepsilon(1-\varepsilon)} A_{LU}^{\sin\phi_h} \sin\phi_h \right]$$

$$+ S_L \left[\sqrt{2\varepsilon(1+\varepsilon)} A_{UL}^{\sin\phi_h} \sin\phi_h + \varepsilon A_{UL}^{\sin 2\phi_h} \sin 2\phi_h \right] \\ + S_L \lambda \left[\sqrt{1-\varepsilon^2} A_{LL} + \sqrt{2\varepsilon(1-\varepsilon)} A_{LL}^{\cos\phi_h} \cos\phi_h \right]$$

$$\times \left\{ \begin{array}{l} A_{UT}^{\sin(\phi_h - \phi_s)} \sin(\phi_h - \phi_s) \\ + \varepsilon A_{UT}^{\sin(\phi_h + \phi_s)} \sin(\phi_h + \phi_s) \\ + \varepsilon A_{UT}^{\sin(3\phi_h - \phi_s)} \sin(3\phi_h - \phi_s) \\ + \sqrt{2\varepsilon(1+\varepsilon)} A_{UT}^{\sin\phi_s} \sin\phi_s \\ + \sqrt{2\varepsilon(1+\varepsilon)} A_{UT}^{\sin(2\phi_h - \phi_s)} \sin(2\phi_h - \phi_s) \end{array} \right\}$$

$$+ S_T \lambda \left[\begin{array}{l} \sqrt{(1-\varepsilon^2)} A_{LT}^{\cos(\phi_h - \phi_s)} \cos(\phi_h - \phi_s) \\ + \sqrt{2\varepsilon(1-\varepsilon)} A_{LT}^{\cos\phi_s} \cos\phi_s \\ + \sqrt{2\varepsilon(1-\varepsilon)} A_{LT}^{\cos(2\phi_h - \phi_s)} \cos(2\phi_h - \phi_s) \end{array} \right]$$



$$A_{UT}^{\sin(\phi_h - \phi_s)} \propto f_{1T}^{\perp q} \otimes D_{1q}^h \quad \text{Sivers}$$

$$A_{UT}^{\sin(\phi_h + \phi_s)} \propto h_1^q \otimes H_{1q}^{\perp h} \quad \text{Collins}$$

Twist-2

$$A_{UT}^{\sin(3\phi_h - \phi_s)} \propto h_{1T}^{\perp q} \otimes H_{1q}^{\perp h}$$

$$A_{UT}^{\sin(\phi_s)} \stackrel{WW}{\propto} Q^{-1} (h_1^q \otimes H_{1q}^{\perp h} + f_{1T}^{\perp q} \otimes D_{1q}^h + \dots)$$

$$A_{UT}^{\sin(2\phi_h - \phi_s)} \stackrel{WW}{\propto} Q^{-1} (h_{1T}^{\perp q} \otimes H_{1q}^{\perp h} + f_{1T}^{\perp q} \otimes D_{1q}^h + \dots)$$

$$A_{LT}^{\cos(\phi_h - \phi_s)} \propto g_{1T}^q \otimes D_{1q}^h$$

$$A_{LT}^{\cos(\phi_s)} \stackrel{WW}{\propto} Q^{-1} (g_{1T}^q \otimes D_{1q}^h + \dots)$$

$$A_{LT}^{\cos(2\phi_h - \phi_s)} \stackrel{WW}{\propto} Q^{-1} (g_{1T}^q \otimes D_{1q}^h + \dots)$$

SIDIS TSAs: Collins and Sivers effects (deuteron)



$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T A_{UT}^{\sin(\phi_h - \phi_S)} \sin(\phi_h - \phi_S) + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_S)} \sin(\phi_h + \phi_S) \dots \right\}$$



$$F_{UT}^{\sin(\phi_h + \phi_S)} = C \left[-\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$



$$F_{UT,T}^{\sin(\phi_h - \phi_S)} = C \left[-\frac{\hat{h} \cdot k_T}{M} f_{1T}^{\perp q} D_{1q}^h \right], F_{UT,L}^{\sin(\phi_h - \phi_S)} = 0$$



COMPASS PLB 673 (2009) 127



- 1st COMPASS deuteron measurements
- Collins and Sivers asymmetries compatible with zero within uncertainties.

SIDIS TSAs: Collins and Sivers effects (proton)



$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T A_{UT}^{\sin(\phi_h - \phi_S)} \sin(\phi_h - \phi_S) + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_S)} \sin(\phi_h + \phi_S) \dots \right\}$$


$$F_{UT}^{\sin(\phi_h + \phi_S)} = C \left[-\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$




$$F_{UT,T}^{\sin(\phi_h - \phi_S)} = C \left[-\frac{\hat{h} \cdot k_T}{M} f_{1T}^{\perp q} D_{1q}^h \right], F_{UT,L}^{\sin(\phi_h - \phi_S)} = 0$$



COMPASS PLB 744(2015)250




Collins



Sivers

- 1st COMPASS deuteron measurements – Collins and Sivers asymmetries compatible with zero
- COMPASS proton measurements – clear non-zero signal for both asymmetries

SIDIS TSAs: Collins effect and Transversity






$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_s} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_s)} \sin(\phi_h + \phi_s) + \dots \right\}$$

$$F_{UT}^{\sin(\phi_h + \phi_s)} = C \left[-\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$




- Measured on P/D in SIDIS and in dihadron SIDIS
- Compatible results COMPASS/HERMES
(Q^2 is different by a factor of ~2-3)
- No impact from Q^2 -evolution?

COMPASS PLB 744 (2015) 250



SIDIS TSAs: Collins effect and Transversity

$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_s} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_s)} \sin(\phi_h + \phi_s) + \dots \right\}$$




$$F_{UT}^{\sin(\phi_h + \phi_s)} = C \left[-\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$




- Measured on P/D in SIDIS and in dihadron SIDIS
- Compatible results COMPASS/HERMES
(Q^2 is different by a factor of ~2-3)
- No impact from Q^2 -evolution?

HERMES, JHEP 12 (2020) 010



COMPASS, PBL 770 (2017) 138



SIDIS TSAs: Collins effect and Transversity



$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_s} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_s)} \sin(\phi_h + \phi_s) + \dots \right\}$$

$$F_{UT}^{\sin(\phi_h + \phi_s)} = C \left[-\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$






- Measured on P/D in SIDIS and in dihadron SIDIS
- Compatible results COMPASS/HERMES
(Q^2 is different by a factor of ~2-3)
- No impact from Q^2 -evolution?


PLB 824 (2022) 136834



- indication for a positive asymmetry
- opposite to π^+ and π^0 as predicted by the models
- Large effect at small P_T



SIDIS TSAs: Collins effect and Transversity






$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_s} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_s)} \sin(\phi_h + \phi_s) + \dots \right\}$$


$$F_{UT}^{\sin(\phi_h + \phi_s)} = C \left[-\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$



- Measured on P/D in SIDIS and in dihadron SIDIS
- Compatible results COMPASS/HERMES
(Q^2 is different by a factor of ~2-3)
- No impact from Q^2 -evolution?
- Extensive phenomenological studies and various global fits by different groups



SIDIS TSAs: Collins effect and Transversity



$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_s} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_s)} \sin(\phi_h + \phi_s) + \dots \right\}$$


$$F_{UT}^{\sin(\phi_h + \phi_s)} = C \left[-\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$



- Measured on P/D in SIDIS and in dihadron SIDIS
- Compatible results COMPASS/HERMES
(Q^2 is different by a factor of ~2-3)
- No impact from Q^2 -evolution?
- Extensive phenomenological studies and various global fits by different groups


[Addendum to the COMPASS-II Proposal]

Projected uncertainties for Collins asymmetry



COMPASS-II (2022)

- 2nd COMPASS deuteron measurements performed
- Crucial to constrain the transversity TMD PDF for the d-quark



SIDIS TSAs: Collins effect and Transversity




$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_s} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_s)} \sin(\phi_h + \phi_s) + \dots \right\}$$

$$F_{UT}^{\sin(\phi_h + \phi_s)} = C \left[-\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$



- Measured on P/D in SIDIS and in dihadron SIDIS
- Compatible results COMPASS/HERMES
(Q^2 is different by a factor of ~2-3)
- No impact from Q^2 -evolution?
- Extensive phenomenological studies and various global fits by different groups


Total protons delivered on the production target: $\sim 5.95 \times 10^{18}$ (98% of the request) in ~ 150 days



SPS efficiency: $\sim 73\%$
Spectrometer efficiency: $\sim 90\%$
Physics data collection efficiency: $\sim 75\%$

Highly successful Run in 2022!

SIDIS TSAs: Collins effect and Transversity





$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_s} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_s)} \sin(\phi_h + \phi_s) + \dots \right\}$$

$$F_{UT}^{\sin(\phi_h + \phi_s)} = C \left[-\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$



- Measured on P/D in SIDIS and in dihadron SIDIS
- Compatible results COMPASS/HERMES
(Q^2 is different by a factor of ~2-3)
- No impact from Q^2 -evolution?
- Extensive phenomenological studies and various global fits by different groups



COMPASS-II (2022)

- 2nd COMPASS deuteron measurements performed
- Crucial to constrain the transversity TMD PDF for the d-quark

SIDIS TSAs: Collins effect and Transversity






$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_s} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_s)} \sin(\phi_h + \phi_s) + \dots \right\}$$

$$F_{UT}^{\sin(\phi_h + \phi_s)} = C \left[-\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$






- Measured on P/D in SIDIS and in dihadron SIDIS
- Compatible results COMPASS/HERMES
(Q^2 is different by a factor of ~2-3)
- No impact from Q^2 -evolution?
- Extensive phenomenological studies and various global fits by different groups



COMPASS-II (2022)


- 2nd COMPASS deuteron measurements performed
- Crucial to constrain the transversity TMD PDF for the d-quark

COMPASS 2022 run: new unique deuteron data




COMPASS 2022 deuteron run

S. Bhattacharya, Z. B. Kang, A. Metz, G. Penn and D. Pitonyak
PRD **105** (2022) 3, 034007



SIDIS TSAs: Kotzinian-Mulders asymmetry

$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + \lambda S_T \sqrt{(1-\varepsilon^2)} A_{LT}^{\cos(\phi_h - \phi_S)} \cos(\phi_h - \phi_S) + \dots \right\}$$



$$A_{LT}^{\cos(\phi_h - \phi_S)} = C \left[\frac{\hat{h} \cdot k_T}{M} g_{1T}^q D_{1q}^h \right]$$




COMPASS/HERMES/CLAS6 results


$$A_{LT}^{\cos(\phi_h - \phi_S)}$$

- Only “twist-2” ingredients
- Sizable non-zero effect for h^+ !**
- Similar effect at HERMES


COMPASS, PBL 770 (2017) 138; PoS QCDEV2017 (2018) 042



See also, PRD 107, (2023) 034016 – global fit by:
M. Horstmann, A. Schafer and A. Vladimirov




First global QCD analysis of the g_{1T} TMD PDF using SIDIS data



SIDIS TSAs: Sivers effect





$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T A_{UT}^{\sin(\phi_h - \phi_S)} \sin(\phi_h - \phi_S) + \dots \right\}$$



$$F_{UT,T}^{\sin(\phi_h - \phi_S)} = C \left[-\frac{\hat{h} \cdot k_T}{M} f_{1T}^{\perp q} D_{1q}^h \right], F_{UT,L}^{\sin(\phi_h - \phi_S)} = 0$$



- COMPASS-HERMES discrepancy
 - Q^2 -evolution?
- T-odd TMD PDF: Expected to change sign between SIDIS and Drell-Yan
- New precise deuteron data





SIDIS Sivers TSA in COMPASS Drell-Yan Q²-ranges




$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T A_{UT}^{\sin(\phi_h - \phi_S)} \sin(\phi_h - \phi_S) + \dots \right\}$$

$$F_{UT,T}^{\sin(\phi_h - \phi_S)} = C \left[-\frac{\hat{h} \cdot k_T}{M} f_{1T}^{\perp q} D_{1q}^h \right], F_{UT,L}^{\sin(\phi_h - \phi_S)} = 0$$



28 September 2023

COMPASS PLB 770 (2017) 138



1st COMPASS multi-D fit done for all eight TSAs

B. Parsamyan

43

COMPASS Multi-D TSA analyses




$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_s} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T A_{UT}^{\sin(\phi_h - \phi_s)} \sin(\phi_h - \phi_s) + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_s)} \sin(\phi_h + \phi_s) \dots \right\}$$

$$F_{UT,T}^{\sin(\phi_h - \phi_s)} = C \left[-\frac{\hat{h} \cdot k_T}{M} f_{1T}^{\perp q} D_{1q}^h \right], F_{UT,L}^{\sin(\phi_h - \phi_s)} = 0$$




$$F_{UT}^{\sin(\phi_h + \phi_s)} = C \left[-\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$




3D x:Q²:z or x:Q²:p_T x:z:p_T

- No clear Q²-dependence within statistical accuracy
- Possible decreasing trend for Sivers TSA?

B.Parsamyan (for COMPASS) [arXiv:1504.01599](https://arxiv.org/abs/1504.01599) [hep-ex] (SPIN-2014)



B. Parsamyan




SIDIS and single-polarized DY x-sections at twist-2 (LO)



$$\frac{d\sigma^{LO}}{dx dy dz dp_T^2 d\phi_h d\phi_s} \propto (F_{UU,T} + \varepsilon F_{UU,L})$$

SIDIS

$$\times \left\{ \begin{array}{l} 1 + \varepsilon A_{UU}^{\cos 2\phi_h} \cos 2\phi_h \\ + S_L \varepsilon A_{UL}^{\sin 2\phi_h} \sin 2\phi_h + S_L \lambda \sqrt{1-\varepsilon^2} A_{LL} \\ \times \left[\begin{array}{l} A_{UT}^{\sin(\phi_h-\phi_s)} \sin(\phi_h-\phi_s) \\ + \varepsilon A_{UT}^{\sin(\phi_h+\phi_s)} \sin(\phi_h+\phi_s) \\ + \varepsilon A_{UT}^{\sin(3\phi_h-\phi_s)} \sin(3\phi_h-\phi_s) \end{array} \right] \\ + S_T \lambda \left[\sqrt{(1-\varepsilon^2)} A_{LT}^{\cos(\phi_h-\phi_s)} \cos(\phi_h-\phi_s) \right] \end{array} \right\}$$




$$\frac{d\sigma^{LO}}{dq^4 d\Omega} \propto F_U^1 (1 + \cos^2 \theta_{CS})$$

DY

$$\times \left\{ \begin{array}{l} 1 + D_{[\sin^2 \theta_{CS}]} A_U^{\cos 2\phi_{CS}} \cos 2\phi_{CS} \\ + S_L \sin^2 \theta_{CS} A_L^{\sin 2\phi_{CS}} \sin 2\phi_{CS} \\ \times \left[\begin{array}{l} A_T^{\sin \varphi_s} \sin \varphi_s \\ + S_T \left[+ D_{[\sin^2 \theta_{CS}]} \left(\begin{array}{l} A_T^{\sin(2\phi_{CS}-\varphi_s)} \sin(2\phi_{CS}-\varphi_s) \\ + A_T^{\sin(2\phi_{CS}+\varphi_s)} \sin(2\phi_{CS}+\varphi_s) \end{array} \right) \right] \end{array} \right] \end{array} \right\}$$


where $D_{[\sin^2 \theta_{CS}]} = \sin^2 \theta_{CS} / (1 + \cos^2 \theta_{CS})$



SIDIS and single-polarized DY x-sections at twist-2 (LO)



$$\frac{d\sigma^{LO}}{dx dy dz dp_T^2 d\phi_h d\phi_s} \propto (F_{UU,T} + \varepsilon F_{UU,L})$$




$$1 + \varepsilon A_{UU}^{\cos 2\phi_h} \cos 2\phi_h \\ + S_L \varepsilon A_{UL}^{\sin 2\phi_h} \sin 2\phi_h + S_L \lambda \sqrt{1-\varepsilon^2} A_{LL}$$



$$\times \left\{ \begin{array}{l} A_{UT}^{\sin(\phi_h - \phi_s)} \sin(\phi_h - \phi_s) \\ + \varepsilon A_{UT}^{\sin(\phi_h + \phi_s)} \sin(\phi_h + \phi_s) \\ + \varepsilon A_{UT}^{\sin(3\phi_h - \phi_s)} \sin(3\phi_h - \phi_s) \\ + S_T \lambda \left[\sqrt{(1-\varepsilon^2)} A_{LT}^{\cos(\phi_h - \phi_s)} \cos(\phi_h - \phi_s) \right] \end{array} \right\}$$

$$\frac{d\sigma^{LO}}{dq^4 d\Omega} \propto F_U^1 (1 + \cos^2 \theta_{CS})$$



$$1 + D_{[\sin^2 \theta_{CS}]} A_U^{\cos 2\phi_{CS}} \cos 2\phi_{CS} \\ + S_L \sin^2 \theta_{CS} A_L^{\sin 2\phi_{CS}} \sin 2\phi_{CS}$$

$$+ S_T \left[\begin{array}{l} A_T^{\sin \varphi_s} \sin \varphi_s \\ + D_{[\sin^2 \theta_{CS}]} \left(A_T^{\sin(2\phi_{CS} - \varphi_s)} \sin(2\phi_{CS} - \varphi_s) \right. \\ \left. + A_T^{\sin(2\phi_{CS} + \varphi_s)} \sin(2\phi_{CS} + \varphi_s) \right) \end{array} \right]$$



where $D_{[\sin^2 \theta_{CS}]} = \sin^2 \theta_{CS} / (1 + \cos^2 \theta_{CS})$

$A_{UU}^{\cos 2\phi_h} \propto h_1^{\perp q} \otimes H_{1q}^{\perp h} + \dots$	Boer-Mulders	$A_U^{\cos 2\phi_{CS}} \propto h_{1,\pi}^{\perp q} \otimes h_{1,p}^{\perp q}$
$A_{UT}^{\sin(\phi_h - \phi_s)} \propto f_{1T}^{\perp q} \otimes D_{1q}^h$	Sivers	$A_T^{\sin \varphi_s} \propto f_{1,\pi}^q \otimes f_{1T,p}^{\perp q}$
$A_{UT}^{\sin(\phi_h + \phi_s)} \propto h_1^q \otimes H_{1q}^{\perp h}$	Transversity	$A_T^{\sin(2\phi_{CS} - \varphi_s)} \propto h_{1,\pi}^{\perp q} \otimes h_{1,p}^q$
$A_{UT}^{\sin(3\phi_h - \phi_s)} \propto h_{1T}^{\perp q} \otimes H_{1q}^{\perp h}$	Pretzelosity	$A_T^{\sin(2\phi_{CS} + \varphi_s)} \propto h_{1,\pi}^{\perp q} \otimes h_{1T,p}^{\perp q}$

- Sign-change of T-odd Sivers and Boer-Mulders TMD PDFs;
- Multiple access to Collins FF $H_{1q}^{\perp h}$ and pion Boer-Mulders PDF $h_{1,\pi}^{\perp q}$

SIDIS and single-polarized DY x-sections at twist-2 (LO)

- $2.5 < M/(\text{GeV}/c^2) < 4.3$ “Charmonia mass”
 - Strong J/ψ -signal → study of J/ψ physics
 - Good signal/background
- $4.3 < M/(\text{GeV}/c^2) < 8.5$ “High mass”
 - Low DY cross-section
 - Beyond charmonium region, background < 3%
 - Valence region → largest asymmetries



$$\frac{d\sigma^{LO}}{dq^4 d\Omega} \propto F_U^1 \left(1 + \cos^2 \theta_{CS} \right)$$

$$1 + D_{[\sin^2 \theta_{CS}]} A_U^{\cos 2\phi_{CS}} \cos 2\phi_{CS}$$


$$+ S_L \sin^2 \theta_{CS} A_L^{\sin 2\phi_{CS}} \sin 2\phi_{CS}$$

$$+ S_T \left[A_T^{\sin \varphi_S} \sin \varphi_S \right.$$


$$\left. + D_{[\sin^2 \theta_{CS}]} \left(A_T^{\sin(2\phi_{CS} - \varphi_S)} \sin(2\phi_{CS} - \varphi_S) \right. \right.$$

$$\left. \left. + A_T^{\sin(2\phi_{CS} + \varphi_S)} \sin(2\phi_{CS} + \varphi_S) \right) \right]$$


$D_{[\sin^2 \theta_{CS}]} = \sin^2 \theta_{CS} / (1 + \cos^2 \theta_{CS})$



Single-polarized DY measurements at COMPASS



HM events are in the valence quark range



28 September 2023

B. Parsamyan

$$\frac{d\sigma^{LO}}{dq^4 d\Omega} \propto F_U^1 \left(1 + \cos^2 \theta_{CS} \right)$$


$$1 + D_{[\sin^2 \theta_{CS}]} A_U^{\cos 2\varphi_{CS}} \cos 2\varphi_{CS}$$

$$+ S_L \sin^2 \theta_{CS} A_L^{\sin^2 \varphi_{CS}} \sin 2\varphi_{CS}$$

$$+ S_T \left[A_T^{\sin \varphi_S} \sin \varphi_S + D_{[\sin^2 \theta_{CS}]} \left(A_T^{\sin(2\varphi_{CS} - \varphi_S)} \sin(2\varphi_{CS} - \varphi_S) + A_T^{\sin(2\varphi_{CS} + \varphi_S)} \sin(2\varphi_{CS} + \varphi_S) \right) \right]$$

$$D_{[\sin^2 \theta_{CS}]} = \sin^2 \theta_{CS} / (1 + \cos^2 \theta_{CS})$$

$4.3 < M / (\text{GeV}/c^2) < 8.5$ “High mass” range
Beyond charmonium region, background < 3%
Valence region → largest asymmetries





49

Drell-Yan TSAs – Transversity

$$\frac{d\sigma}{dq^4 d\Omega} \propto 1 + \dots + S_T \left[D_{[\sin^2 \theta_{CS}]} A_T^{\sin(2\varphi_{CS} - \varphi_s)} \sin(2\varphi_{CS} - \varphi_s) + \dots \right]$$

Transversity DY TSA


$$A_T^{\sin(2\varphi_{CS} - \varphi_s)} \propto h_{1,\pi}^{\perp q} \otimes h_{1,p}^q$$



Collins SIDIS TSA

$$A_{UT}^{\sin(\phi_h + \phi_s - \pi)} \propto h_1^q \otimes H_{1q}^{\perp h}$$

$$A_{UT}^{\sin(\phi_h + \phi_s - \pi)}$$




Drell-Yan TSAs – Transversity

$$\frac{d\sigma}{dq^4 d\Omega} \propto 1 + \dots + S_T \left[D_{[\sin^2 \theta_{CS}]} A_T^{\sin(2\varphi_{CS} - \varphi_s)} \sin(2\varphi_{CS} - \varphi_s) + \dots \right]$$

Transversity DY TSA


$$A_T^{\sin(2\varphi_{CS} - \varphi_s)} \propto h_{1,\pi}^{\perp q} \otimes h_{1,p}^q$$



Collins SIDIS TSA

$$A_{UT}^{\sin(\phi_h + \phi_s - \pi)} \propto h_1^q \otimes H_{1q}^{\perp h}$$

COMPASS PLB 770 (2017) 138




Drell-Yan TSAs – Sivers

$$\frac{d\sigma}{dq^4 d\Omega} \propto 1 + \dots + S_T [A_T^{\sin \varphi_S} \sin \varphi_S + \dots]$$

Sivers DY TSA


$$A_T^{\sin \varphi_S} \propto f_{1,\pi}^q \otimes f_{1T,p}^{\perp q}$$



Sivers SIDIS TSA

$$A_{UT}^{\sin(\phi_h - \phi_s)} \propto f_{1T}^{\perp q} \otimes D_{1q}^h$$



COMPASS
PLB 770 (2017) 138



DY TSAs at COMPASS (high-mass range)




Theory curves based on S. Bastami et al. JHEP 02, (2021), 166



- General agreement with available theory predictions

3D unpolarized Drell-Yan cross section on NH₃ and W



Conclusions




- Importance of careful understanding and confrontation of experimental data from different experiments
 - Different kinematic domains and phase-space limitations
 - Experiments employ complex analysis techniques, Monte-Carlo simulations, and sophisticated corrections (acceptance, VMs, radiative corrections)
- Close collaboration between different experiments → general benefit for the field
 - Knowledge transfer, comparison of the analysis techniques, tools, and methodology, cross-analyses between different experiments
- Close collaboration between experiment and phenomenology/theory
 - Flexibility in adapting on the analysis side to the choice of the observables, phase-space selections, etc. (before publishing the data)
 - Different possibilities for common paper projects, external membership
- Possibility to organize effective and fruitful collaborative work

Conclusions



- COMPASS holds the record for the longest-running CERN experiment
(20 years of data-taking)
- Series of successful and important measurements addressing nucleon spin-structure
 - Inclusive measurements, unpolarized and polarized SIDIS (longitudinal/transverse)
 - First-ever polarized Drell-Yan measurements
- A wealth of (SI)DIS, Drell-Yan, DVCS, HEMP data collected across the years
 - **Petabytes of data available for analysis**
- Wide and unique kinematic domain accessing low x and large Q^2
 - **Will remain unique for at least another decade**
- World-unique SIDIS deuteron data collected in 2022
 - **Highly successful run, promising preliminary results**
- Since 2023 the experiment entered the Analysis Phase
 - The spectrometer has been transferred to the COMPASS successor in the M2 beamline – the AMBER collaboration
 - **3 new groups joined COMPASS in the course of 2023 for the Analysis Phase**
 - **If you are interested – don't hesitate to get in touch!**

Thank You!



Joint XX-th International Workshop on Hadron Structure and Spectroscopy




and 5-th Workshop on Correlations in
Partonic and Hadronic Interactions

Yerevan, Armenia

30 September – 4 October, 2024





Joint XX-th International Workshop on Hadron Structure and Spectroscopy



and 5-th Workshop on Correlations in
Partonic and Hadronic Interactions

Yerevan, Armenia

30 September – 4 October, 2024

3896 m

5137 m

