

Nucleon spin and TMD studies with COMPASS experiment: selected highlights

BAKUR PARSAMYAN

AANL, CERN, INFN section of Turin

on behalf of the COMPASS Collaboration



Strong QCD from Hadron Structure Experiments - VI (SQCD VI)
Nanjing University, Science and Technology building
May 14 – 17, 2024, Nanjing, China

COMPASS collaboration



Common Muon and Proton Apparatus for Structure and Spectroscopy



28 institutions from 14 countries

– nearly 210 physicists (in 2023: start of the Analysis Phase)

- CERN SPS North Area
- Fixed target experiment
- Approved in 1997 (25 years)
- Taking data since 2002 (20 years)

Wide physics program


COMPASS phase-I

- Data taking 2002-2011
- Muon and hadron programs
- Nucleon spin structure (SI)DIS
- Spectroscopy and exotics

COMPASS phase-II

- Data taking 2012-2022
- Primakoff
- DVCS (GPD+SIDIS)
- Polarized Drell-Yan
- Transverse deuteron SIDIS 2022

3 new groups joined the COMPASS collaboration in 2023
UConn (US), AANL (Armenia), NCU (Taiwan)



COMPASS web page: <http://wwwcompass.cern.ch>

COMPASS experimental setup




COmmon Muon Proton Apparatus for Structure and Spectroscopy



CERN SPS North Area (building 888)

Two-stage spectrometer LAS+SAS

- Large Angle Spectrometer (SM1 magnet)
- Small Angle Spectrometer (SM2 magnet)




- Primary beam - 400 GeV p from SPS
 - impinging on Be production target (T6)
- 190 GeV secondary hadron beams
 - h^- beam: 97% π^- , 2% K^- , 1% p
 - h^+ beam: 75% π^+ , 24% p , 1% K^+
- 160 GeV tertiary muon beams
 - μ^\pm longitudinally polarized



Large-acceptance forward spectrometer

- Precise tracking (350 planes)
SciFi, Silicon, MicroMegas, GEM, MWPC, DC, Straw, Muon walls
- PID - CEDARs, RICH, calorimeters, MWs
- Various targets:
 - Polarized solid-state NH_3 or 6LiD
 - Liquid H_2
 - Solid-state nuclear targets (e.g. Ni, W, Pb)



COMPASS experimental setup: Phase II (SIDIS programme)




COmmon Muon Proton Apparatus for Structure and Spectroscopy


CERN SPS North Area (building 888)

Two-stage spectrometer LAS+SAS


- Large Angle Spectrometer (SM1 magnet)
- Small Angle Spectrometer (SM2 magnet)



- Primary beam - 400 GeV p from SPS
 - impinging on Be production target (T6)
- 190 GeV secondary hadron beams
 - h^- beam: 97% π^- , 2% K^- , 1% p
 - h^+ beam: 75% π^+ , 24% p , 1% K^+
- 160 GeV tertiary muon beams
 - μ^+ longitudinally polarized



- Polarized solid-state NH_3 or 6LiD
- Two or three oppositely polarized cells
- Longitudinal and transverse polarization



COMPASS experimental setup: Phase II (DY programme)




COmmon Muon Proton Apparatus for Structure and Spectroscopy




CERN SPS North Area (building 888)

Two-stage spectrometer LAS+SAS

- Large Angle Spectrometer (SM1 magnet)
- Small Angle Spectrometer (SM2 magnet)




- Primary beam - 400 GeV p from SPS
 - impinging on Be production target (T6)
- 190 GeV secondary hadron beams
 - h^- beam: 97% π^- , 2% K^- , 1% p
 - h^+ beam: 75% π^+ , 24% p , 1% K^+
- 160 GeV tertiary muon beams
 - μ^\pm longitudinally polarized







- Solid-state NH_3 material
- Two cells
- oppositely polarized
- Transverse polarization

COMPASS timeline






COMPASS timeline






Turning the needle A snapshot of part of the COMPASS spectrometer. Credit: P. Traczyk / CERN-PHOTO-202104-060-2

The COMPASS experiment at CERN has reported the first direct evidence for a long-lived resonance in pion decays which can masquerade as a resonance.




COMPASS Legacy



FLAVOUR PHYSICS | NEWS



COMPASS points to triangle singularity

23 August 2021




Turning the needle A snapshot of part of the COMPASS spectrometer. Credit: P. PHOTO-202104-060-2

The COMPASS experiment at CERN has reported the first direct evidence of an exotic state which can masquerade as a pion.



A000BER
Apparatus for Meson and Baryon
Experimental Research



SIDIS L/T

Spectroscopy SIDIS

S L/T
Primakoff
DVCS (pilot)
D

AMBER

58

2021
S-3



The COMPASS Experiment at the CERN SPS

Broad Physics Program to study Structure and Excitation Spectrum of Hadrons

Increasing resolution scale
(momentum transfer)


Nucleon structure

- Hard scattering of μ^\pm and π^- off (un)polarized P/D targets
- Study of nucleon spin structure
- Parton distribution functions and fragmentation functions




Hadron spectroscopy

- Diffractive $\pi(K)$ dissociation reaction with proton target
- PWA technique employed
- High-precision measurement of light-meson excitation spectrum
- Search for exotic states



Chiral dynamics

- Test chiral perturbation theory in $\pi(K)\gamma$ reactions
- π^\pm and K^\pm polarizabilities
- Chiral anomaly $F_{3\pi}$





The COMPASS Experiment at the CERN SPS

Broad Physics Program to study Structure and Excitation Spectrum of Hadrons

Increasing resolution scale
(momentum transfer)

Nucleon structure

- Hard scattering of μ^\pm and π^- off (un)polarized P/D targets
- Study of nucleon spin structure
- Parton distribution functions and fragmentation functions




Hadron spectroscopy

- Diffractive $\pi(K)$ dissociation reaction with proton target
- PWA technique employed
- High-precision measurement of light-meson excitation spectrum
- Search for exotic states




Chiral dynamics

- Test chiral perturbation theory in $\pi(K)\gamma$ reactions
- π^\pm and K^\pm polarizabilities
- Chiral anomaly $F_{3\pi}$



Nucleon 3D structure: GPDs

- Transverse position \vec{b}_T of partons
 - Correlation between \vec{b}_T and x
 - Complementary to TMD PDFs
- 8 generalized parton distribution functions (GPDs)
 - Contain information about parton orbital angular momentum
 - Mostly unknown
- COMPASS exclusive process measurements:
 - Deeply virtual Compton scattering (DVCS): $\mu + N \rightarrow \mu + \gamma + N$
 - Hard exclusive meson production (HEMP): $\mu + N \rightarrow \mu + \text{Meson} + N$ with $\pi^0, \rho(770), \omega(782), \dots$



Nucleon spin structure: collinear approach \leftrightarrow TMDs





quark			
	U	L	T
U	$f_1^q(x)$ number density		
L		$g_1^q(x)$ helicity	
T			$h_1^q(x)$ transversity



quark			
	U	L	T
U	$f_1^q(x, \mathbf{k}_T^2)$ number density		$h_1^{\perp q}(x, \mathbf{k}_T^2)$ Boer-Mulders
L		$g_1^q(x, \mathbf{k}_T^2)$ helicity	$h_{1L}^{\perp q}(x, \mathbf{k}_T^2)$ worm-gear L
T	$f_{1T}^{\perp q}(x, \mathbf{k}_T^2)$ Sivers	$g_{1T}^q(x, \mathbf{k}_T^2)$ worm-gear T	$h_1^q(x, \mathbf{k}_T^2)$ transversity $h_{1T}^{\perp q}(x, \mathbf{k}_T^2)$ pretzelosity

- PDFs – universal (process independent) objects; T-odd PDFs – conditionally universal





Hadron multiplicities; h^\pm , π^\pm and K^\pm (2016 data)

collinear

TMD


A set of complex corrections:

- Acceptance, rad. corrections, PID, diffractive VMs, etc.



New radiative corrections

The article is in a final drafting stage






Hadron multiplicities; h^\pm , π^\pm and K^\pm (2016 data)

collinear




A set of complex corrections:

- Acceptance, rad. corrections, PID, diffractive VMs, etc.




New radiative corrections being applied
Drafting started for a dedicated article

3D unpolarized Drell-Yan cross section on NH₃ and W




- First new results in 30 years!
- Data from light/heavy targets
 - NH₃-He, Al, W
 - Nuclear dependence
- 1D/2D/3D representations
- $x_F:q_T:M$
- Unique data to access collinear and TMD distributions
e.g. pion TMD PDF





B. Parsamyan

3D unpolarized Drell-Yan cross section on NH₃ and W




MAP collaboration
Phys. Rev. D. 107, 014014




- First new results in 30 years!
- Data from light/heavy targets
 - NH₃-He, Al, W
 - Nuclear dependence
- 1D/2D/3D representations
- $x_F:q_T:M$
- Unique data to access collinear and TMD distributions
e.g. pion TMD PDF
- To be included in future global fits (MAP, JAM, etc.)

Nucleon spin structure: azimuthal effects

- 1964 Quark model





- 1969 Parton model



- 1973 asymptotic freedom and QCD



- 1978 intrinsic transverse motion of quarks and azimuthal asymmetries




Cahn effect in SIDIS



$$\frac{d\sigma}{dxdydzdp_T^2d\phi_hd\phi_S} = \left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L}) \times (1 + \underbrace{\sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h}_{+ \dots})$$




$f_1^q(x, k_T^2)$
number density



Cahn effect - R.N. Cahn, PLB 78 (1978)



The point that there are azimuthal dependences, which arise from the transverse momenta of the partons was clearly stated in this papers: T.P. Cheng and A. Zee, Phys. Rev. D6 (1972) 885; F. Ravndal, Phys. Lett. 43B (1973) 301; R.L. Kingsley, Phys. Rev. D10 (1974) 1580; A.M. Kotsynyan, Teor. Mat. Fiz. 24 (1975) 206;



Cahn effect in SIDIS

$$\frac{d\sigma}{dxdydzdp_T^2d\phi_h d\phi_S} = \left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L}) \times (1 + \underbrace{\sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h}_{\text{Cahn effect}} + \dots)$$



Cahn effect

$$f_1^q(x, \mathbf{k}_T^2)$$




number density

As of 1978 – simplistic kinematic effect:

- non-zero \mathbf{k}_T induces an azimuthal modulation

As of 2023 – complex SF (twist-2/3 functions)

- Measurements by different experiments



$$F_{UU}^{\cos\phi_h} = \frac{2M}{Q} C \left\{ -\frac{\hat{\mathbf{h}} \cdot \mathbf{p}_T}{M_h} \left(\textcolor{red}{xh} H_{1q}^{\perp h} + \frac{M_h}{M} f_1^q \frac{\tilde{D}_q^{\perp h}}{z} \right) - \frac{\hat{\mathbf{h}} \cdot \mathbf{k}_T}{M} \left(\textcolor{red}{xf}^{\perp q} D_{1q}^h + \frac{M_h}{M} h_1^{\perp q} \frac{\tilde{H}_q^h}{z} \right) \right\}$$

Cahn effect in SIDIS

$$\frac{d\sigma}{dxdydzdp_T^2d\phi_hd\phi_S} = \left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L}) \times (1 + \underbrace{\sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h}_{\text{Cahn effect}} + \dots)$$



Cahn effect


$f_1^q(x, k_T^2)$
number density

As of 1978 – simplistic kinematic effect:

- non-zero k_T induces an azimuthal modulation

As of 2023 – complex SF (twist-2/3 functions)

- Measurements by different experiments
- Complex multi-D kinematic dependences
 - So far, no comprehensive interpretation



Cahn effect in SIDIS

$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_s} =$$

$$\left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L})$$

$$\times (1 + \sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h + \dots)$$



Cahn effect

$$f_1^q(x, \mathbf{k}_T^2)$$

number density



As of 1978 – simplistic kinematic effect:


- non-zero k_T induces an azimuthal modulation

As of 2023 – complex SF (twist-2/3 functions)


- Measurements by different experiments
 - Complex multi-D kinematic dependences
 - So far, no comprehensive interpretation
 - A set of complex corrections:
 - Acceptance, diffractively produced VMs, radiative corrections (RC), etc.



$\gamma^* \rightarrow \rho^0$ spin components



VM contribution



Cahn effect in SIDIS

$$\frac{d\sigma}{dxdydzdp_T^2d\phi_h d\phi_S} = \left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L}) \times (1 + \underbrace{\sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h}_{\text{Cahn effect}} + \dots)$$



Cahn effect

$f_1^q(x, k_T^2)$
number density


As of 1978 – simplistic kinematic effect:

- non-zero k_T induces an azimuthal modulation


As of 2023 – complex SF (twist-2/3 functions)

- Measurements by different experiments
- Complex multi-D kinematic dependences
 - So far, no comprehensive interpretation
- A set of complex corrections:
 - Acceptance, **diffractively produced VMs**, radiative corrections (RC), etc.

COMPASS, EPJC (2023) 83 924



VM corrections, applied



Cahn effect in SIDIS

COMPASS, EPJC (2023) 83 924



$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_s} = \left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L}) \\ \times (1 + \underbrace{\sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h}_{\text{Cahn effect}} + \dots)$$



Cahn effect

$f_1^q(x, k_T^2)$
number density

As of 1978 – simplistic kinematic effect:


- non-zero k_T induces an azimuthal modulation

As of 2023 – complex SF (twist-2/3 functions)


- Measurements by different experiments
- Complex multi-D kinematic dependences
 - So far, no comprehensive interpretation
- A set of complex corrections:
 - Acceptance, **diffractively produced VMs, radiative corrections (RC), etc.**

NEW!

RC corrections, applied



VM corrections, applied



Cahn effect in SIDIS

$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} = \left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L}) \times (1 + \underbrace{\sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h}_{\text{Cahn effect}} + \dots)$$



Cahn effect

$f_1^q(x, k_T^2)$
number density


As of 1978 – simplistic kinematic effect:

- non-zero k_T induces an azimuthal modulation

As of 2023 – complex SF (twist-2/3 functions)

- Measurements by different experiments
- Complex multi-D kinematic dependences
 - So far, no comprehensive interpretation
- A set of complex corrections:
 - Acceptance, diffractively produced VMs, radiative corrections (RC), etc.
- Strong Q^2 dependence – unexplained
 - Do not seem to come from RCs
 - Transition between TMD \leftrightarrow collinear regions?

Recent COMPASS results




Boer-Mulders effect in SIDIS

$$\frac{d\sigma}{dxdydzdp_T^2d\phi_h d\phi_s} = \left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L}) \\ \times (1 + \sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h + \varepsilon A_{UU}^{\cos 2\phi_h} \cos 2\phi_h + \dots)$$

Arises due to the correlation between quark transverse spin and intrinsic transverse momentum



Quark Nucleon	U		T
U	$f_1^q(x, k_T^2)$ number density 		$h_1^{\perp q}(x, k_T^2)$ Boer-Mulders




$$F_{UU}^{\cos 2\phi_h} = C \left[- \frac{2(\hat{h} \cdot p_T)(\hat{h} \cdot k_T) - p_T \cdot k_T}{MM_h} h_1^{\perp q} H_{1q}^{\perp h} \right]$$

Boer-Mulders effect in SIDIS


$$\frac{d\sigma}{dxdydzdp_T^2d\phi_h d\phi_s} = \left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L}) \\ \times (1 + \sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h + \varepsilon A_{UU}^{\cos 2\phi_h} \cos 2\phi_h + \dots)$$

Quark Nucleon	U		T
U	$f_1^q(x, k_T^2)$ number density 		$h_1^{\perp q}(x, k_T^2)$ Boer-Mulders

Arises due to the correlation between quark transverse spin and intrinsic transverse momentum



$$F_{UU}^{\cos 2\phi_h} = C \left[-\frac{2(\hat{h} \cdot p_T)(\hat{h} \cdot k_T) - p_T \cdot k_T}{MM_h} h_1^{\perp q} H_{1q}^{\perp h} \right]$$




- Collins-like behavior (h^-h^+ - mirror symmetry)?

Boer-Mulders effect in SIDIS

$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_s} = \left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L}) \\ \times (1 + \sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h + \varepsilon A_{UU}^{\cos 2\phi_h} \cos 2\phi_h + \lambda \sqrt{2\varepsilon(1-\varepsilon)} A_{LU}^{\sin\phi_h} \sin\phi_h + \dots)$$

Target spin independent part of the cross-section: three asymmetries



Cahn effect
Different for h^+ , h^-
Non-trivial Q^2 dependence

Boer-Mulders effect
Collins-like behavior
(h^+h^- - mirror symmetry)

Beam-spin asymmetry
higher-twist effect
non-zero, positive trend

Working on 3D kinematic dependences


SIDIS x-section: twist-2 terms and TMDs

$$\frac{d\sigma}{dxdydzdp_T^2d\phi_hd\phi_s} = \left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L})$$

All measured by COMPASS

$1 + \varepsilon A_{UU}^{\cos 2\phi_h} \cos 2\phi_h$ $+ S_L \left[\varepsilon A_{UL}^{\sin 2\phi_h} \sin 2\phi_h \right]$ $+ S_L \lambda \left[\sqrt{1-\varepsilon^2} A_{LL} \right]$	\times $+ S_T \left[\begin{array}{l} A_{UT}^{\sin(\phi_h - \phi_s)} \sin(\phi_h - \phi_s) \\ + \varepsilon A_{UT}^{\sin(\phi_h + \phi_s)} \sin(\phi_h + \phi_s) \\ + \varepsilon A_{UT}^{\sin(3\phi_h - \phi_s)} \sin(3\phi_h - \phi_s) \end{array} \right]$ $+ S_T \lambda \left[\sqrt{(1-\varepsilon^2)} A_{LT}^{\cos(\phi_h - \phi_s)} \cos(\phi_h - \phi_s) \right]$
---	--

Twist-2



Quark	U	L	T
Nucleon	number density		Boer-Mulders
U			
L		helicity	worm-gear L
T	Sivers	Kotzinian-Mulders worm-gear T	transversity pretzelosity
	spin of the nucleon	spin of the quark	k_T

SIDIS x-section: twist-2 terms and twist-3 terms

$$\frac{d\sigma}{dxdydzdp_T^2d\phi_h d\phi_s} =$$

All measured by COMPASS

$$\left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L})$$


$$1 + \sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h + \varepsilon A_{UU}^{\cos 2\phi_h} \cos 2\phi_h \\ + \lambda \sqrt{2\varepsilon(1-\varepsilon)} A_{LU}^{\sin\phi_h} \sin\phi_h$$

$$+ S_L \left[\sqrt{2\varepsilon(1+\varepsilon)} A_{UL}^{\sin\phi_h} \sin\phi_h + \varepsilon A_{UL}^{\sin 2\phi_h} \sin 2\phi_h \right]$$

$$+ S_L \lambda \left[\sqrt{1-\varepsilon^2} A_{LL} + \sqrt{2\varepsilon(1-\varepsilon)} A_{LL}^{\cos\phi_h} \cos\phi_h \right]$$

$$\times \left[\begin{array}{l} A_{UT}^{\sin(\phi_h - \phi_s)} \sin(\phi_h - \phi_s) \\ + \varepsilon A_{UT}^{\sin(\phi_h + \phi_s)} \sin(\phi_h + \phi_s) \\ + \varepsilon A_{UT}^{\sin(3\phi_h - \phi_s)} \sin(3\phi_h - \phi_s) \\ + \sqrt{2\varepsilon(1+\varepsilon)} A_{UT}^{\sin\phi_s} \sin\phi_s \\ + \sqrt{2\varepsilon(1+\varepsilon)} A_{UT}^{\sin(2\phi_h - \phi_s)} \sin(2\phi_h - \phi_s) \end{array} \right] \\ \hline \left[\begin{array}{l} \sqrt{(1-\varepsilon^2)} A_{LT}^{\cos(\phi_h - \phi_s)} \cos(\phi_h - \phi_s) \\ + \sqrt{2\varepsilon(1-\varepsilon)} A_{LT}^{\cos\phi_s} \cos\phi_s \\ + \sqrt{2\varepsilon(1-\varepsilon)} A_{LT}^{\cos(2\phi_h - \phi_s)} \cos(2\phi_h - \phi_s) \end{array} \right]$$

Twist-2
Twist-3



Quark	U	L	T
Nucleon			
U			
L			
T			

spin of the nucleon spin of the quark k_T

SIDIS: target longitudinal spin dependent asymmetries

$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots \right.$$

$$\left. + S_L \left[\sqrt{2\varepsilon(1+\varepsilon)} A_{UL}^{\sin \phi_h} \sin \phi_h + \varepsilon A_{UL}^{\sin 2\phi_h} \sin 2\phi_h \right] \right\}$$


$$+ S_L \lambda \left[\sqrt{1-\varepsilon^2} A_{LL} + \sqrt{2\varepsilon(1-\varepsilon)} A_{LL}^{\cos \phi_h} \cos \phi_h \right]$$

$$F_{LL}^1 = \mathcal{C} \left\{ g_{1L}^q D_{1q}^h \right\}$$

$$F_{UL}^{\sin \phi_h} = \frac{2M}{Q} \mathcal{C} \left\{ -\frac{\hat{\mathbf{h}} \cdot \mathbf{p}_T}{M_h} \left(x h_L^q H_{1q}^{\perp h} + \frac{M_h}{M} g_{1L}^q \frac{\tilde{G}_q^{\perp h}}{z} \right) \right. \\ \left. + \frac{\hat{\mathbf{h}} \cdot \mathbf{k}_T}{M} \left(x f_L^{\perp q} D_{1q}^h - \frac{M_h}{M} h_{1L}^{\perp q} \frac{\tilde{H}_q^h}{z} \right) \right\}$$

$$F_{UL}^{\sin 2\phi_h} = \mathcal{C} \left\{ -\frac{2(\hat{\mathbf{h}} \cdot \mathbf{p}_T)(\hat{\mathbf{h}} \cdot \mathbf{k}_T) - \mathbf{p}_T \cdot \mathbf{k}_T}{MM_h} h_{1L}^{\perp q} H_{1q}^{\perp h} \right\}$$

$$F_{LL}^{\cos \phi_h} = \frac{2M}{Q} \mathcal{C} \left\{ -\frac{\hat{\mathbf{h}} \cdot \mathbf{p}_T}{M_h} \left(x e_L^q H_{1q}^{\perp h} + \frac{M_h}{M} g_{1L}^q \frac{\tilde{D}_q^{\perp h}}{z} \right) \right. \\ \left. + \frac{\hat{\mathbf{h}} \cdot \mathbf{k}_T}{M} \left(x g_L^{\perp q} D_{1q}^h - \frac{M_h}{M} h_{1L}^{\perp q} \frac{\tilde{E}_q^h}{z} \right) \right\}$$



SIDIS: target longitudinal spin dependent asymmetries

$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots \right.$$

$$+ S_L \left[\sqrt{2\varepsilon(1+\varepsilon)} A_{UL}^{\sin \phi_h} \sin \phi_h + \varepsilon A_{UL}^{\sin 2\phi_h} \sin 2\phi_h \right] \left. \right\}$$

$$+ S_L \lambda \left[\sqrt{1-\varepsilon^2} A_{LL} + \sqrt{2\varepsilon(1-\varepsilon)} A_{LL}^{\cos \phi_h} \cos \phi_h \right]$$

COMPASS collected large amount of L-SIDIS data
Unprecedented precision for some amplitudes!

$$A_{UL}^{\sin \phi_h}$$

- Q-suppression, Various different “twist” ingredients
- Sizable TSA-mixing
- Significant h^+ asymmetry, clear z -dependence
- h^- compatible with zero


$$A_{UL}^{\sin 2\phi_h}$$

- Only “twist-2” ingredients
- Additional p_T -suppression
- Compatible with zero, in agreement with models
- Collins-like behavior?

$$A_{LL}^{\cos \phi_h}$$

- Q-suppression, Various different “twist” ingredients
- Compatible with zero, in agreement with models

B. Parsamyan (for COMPASS) [arXiv:1801.01488](https://arxiv.org/abs/1801.01488) [hep-ex]






SIDIS: target longitudinal spin dependent asymmetries



$$\frac{d\sigma}{dx dy dz dp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_L \lambda \sqrt{1-\varepsilon^2} A_{LL} + \dots \right\}$$

$$F_{LL}^1 = C \left\{ g_{1L}^q D_{1q}^h \right\}$$

- Measurement of (semi-)inclusive $A_1(A_{LL})$ is one of the key physics topics of HERMES/COMPASS
- Large amount of P/D data
- No P_T -dependence observed




SIDIS x-section and TMDs at twist-2: TSAs

$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_s} =$$

All measured by COMPASS

$$\left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L})$$

$$\begin{aligned} & \left[1 + \sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h + \varepsilon A_{UU}^{\cos 2\phi_h} \cos 2\phi_h \right. \\ & \quad \left. + \lambda \sqrt{2\varepsilon(1-\varepsilon)} A_{LU}^{\sin\phi_h} \sin\phi_h \right] \\ & + S_L \left[\sqrt{2\varepsilon(1+\varepsilon)} A_{UL}^{\sin\phi_h} \sin\phi_h + \varepsilon A_{UL}^{\sin 2\phi_h} \sin 2\phi_h \right] \\ & + S_L \lambda \left[\sqrt{1-\varepsilon^2} A_{LL} + \sqrt{2\varepsilon(1-\varepsilon)} A_{LL}^{\cos\phi_h} \cos\phi_h \right] \\ & \times \left[\begin{aligned} & A_{UT}^{\sin(\phi_h - \phi_s)} \sin(\phi_h - \phi_s) \\ & + \varepsilon A_{UT}^{\sin(\phi_h + \phi_s)} \sin(\phi_h + \phi_s) \\ & + \varepsilon A_{UT}^{\sin(3\phi_h - \phi_s)} \sin(3\phi_h - \phi_s) \\ & + \sqrt{2\varepsilon(1+\varepsilon)} A_{UT}^{\sin\phi_s} \sin\phi_s \\ & + \sqrt{2\varepsilon(1+\varepsilon)} A_{UT}^{\sin(2\phi_h - \phi_s)} \sin(2\phi_h - \phi_s) \end{aligned} \right] \\ & + S_T \left[\begin{aligned} & \sqrt{(1-\varepsilon^2)} A_{LT}^{\cos(\phi_h - \phi_s)} \cos(\phi_h - \phi_s) \\ & + \sqrt{2\varepsilon(1-\varepsilon)} A_{LT}^{\cos\phi_s} \cos\phi_s \\ & + \sqrt{2\varepsilon(1-\varepsilon)} A_{LT}^{\cos(2\phi_h - \phi_s)} \cos(2\phi_h - \phi_s) \end{aligned} \right] \end{aligned}$$



$$A_{UT}^{\sin(\phi_h - \phi_s)} \propto f_{1T}^{\perp q} \otimes D_{1q}^h$$

Sivers

$$A_{UT}^{\sin(\phi_h + \phi_s)} \propto h_1^q \otimes H_{1q}^{\perp h}$$

Collins

$$A_{UT}^{\sin(3\phi_h - \phi_s)} \propto h_{1T}^{\perp q} \otimes H_{1q}^{\perp h}$$

Twist-2

$$A_{UT}^{\sin(\phi_s)} \stackrel{WW}{\propto} Q^{-1} (h_1^q \otimes H_{1q}^{\perp h} + f_{1T}^{\perp q} \otimes D_{1q}^h + \dots)$$

$$A_{UT}^{\sin(2\phi_h - \phi_s)} \stackrel{WW}{\propto} Q^{-1} (h_{1T}^{\perp q} \otimes H_{1q}^{\perp h} + f_{1T}^{\perp q} \otimes D_{1q}^h + \dots)$$

$$A_{LT}^{\cos(\phi_h - \phi_s)} \propto g_{1T}^q \otimes D_{1q}^h$$

$$A_{LT}^{\cos(\phi_s)} \stackrel{WW}{\propto} Q^{-1} (g_{1T}^q \otimes D_{1q}^h + \dots)$$

$$A_{LT}^{\cos(2\phi_h - \phi_s)} \stackrel{WW}{\propto} Q^{-1} (g_{1T}^q \otimes D_{1q}^h + \dots)$$

SIDIS TSAs: Collins and Sivers effects (deuteron)



$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T A_{UT}^{\sin(\phi_h - \phi_S)} \sin(\phi_h - \phi_S) + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_S)} \sin(\phi_h + \phi_S) \dots \right\}$$



$$F_{UT}^{\sin(\phi_h + \phi_S)} = C \left[-\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$



$$F_{UT,T}^{\sin(\phi_h - \phi_S)} = C \left[-\frac{\hat{h} \cdot k_T}{M} f_{1T}^{\perp q} D_{1q}^h \right], F_{UT,L}^{\sin(\phi_h - \phi_S)} = 0$$



COMPASS PLB 673 (2009) 127



- 1st COMPASS deuteron measurements
- Collins and Sivers asymmetries compatible with zero within uncertainties.

SIDIS TSAs: Collins and Sivers effects (proton)



$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T A_{UT}^{\sin(\phi_h - \phi_S)} \sin(\phi_h - \phi_S) + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_S)} \sin(\phi_h + \phi_S) \dots \right\}$$



$$F_{UT}^{\sin(\phi_h + \phi_S)} = C \left[-\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$



$$F_{UT,T}^{\sin(\phi_h - \phi_S)} = C \left[-\frac{\hat{h} \cdot k_T}{M} f_{1T}^{\perp q} D_{1q}^h \right], F_{UT,L}^{\sin(\phi_h - \phi_S)} = 0$$




COMPASS PLB 744(2015)250



- 1st COMPASS deuteron measurements – Collins and Sivers asymmetries compatible with zero
- COMPASS proton measurements – clear non-zero signal for both asymmetries

SIDIS TSAs: Collins effect and Transversity






$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_S)} \sin(\phi_h + \phi_S) + \dots \right\}$$

$$F_{UT}^{\sin(\phi_h + \phi_S)} = C \left[-\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$




- Measured on P/D in SIDIS and in dihadron SIDIS
- Compatible results COMPASS/HERMES
(Q^2 is different by a factor of $\sim 2-3$)
- No impact from Q^2 -evolution?

COMPASS PLB 744 (2015) 250



SIDIS TSAs: Collins effect and Transversity




$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_S)} \sin(\phi_h + \phi_S) + \dots \right\}$$

$$F_{UT}^{\sin(\phi_h + \phi_S)} = C \left[-\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$




- Measured on P/D in SIDIS and in dihadron SIDIS
- Compatible results COMPASS/HERMES
(Q^2 is different by a factor of ~ 2 -3)
- No impact from Q^2 -evolution?

HERMES, JHEP 12 (2020) 010




B. Parsamyan

COMPASS, PBL 770 (2017) 138



SIDIS TSAs: Collins effect and Transversity






$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_S)} \sin(\phi_h + \phi_S) + \dots \right\}$$


$$F_{UT}^{\sin(\phi_h + \phi_S)} = C \left[-\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$



- Measured on P/D in SIDIS and in dihadron SIDIS
- Compatible results COMPASS/HERMES (Q^2 is different by a factor of ~ 2 -3)
- No impact from Q^2 -evolution?
- Extensive phenomenological studies and various global fits by different groups



SIDIS TSAs: Collins effect and Transversity



$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_S)} \sin(\phi_h + \phi_S) + \dots \right\}$$


$$F_{UT}^{\sin(\phi_h + \phi_S)} = C \left[-\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$



- Measured on P/D in SIDIS and in dihadron SIDIS
- Compatible results COMPASS/HERMES (Q^2 is different by a factor of ~ 2 -3)
- No impact from Q^2 -evolution?
- Extensive phenomenological studies and various global fits by different groups


[Addendum to the COMPASS-II Proposal]

Projected uncertainties for Collins asymmetry




COMPASS-II (2022)

- 2nd COMPASS deuteron measurements performed
- Crucial to constrain the transversity TMD PDF for the d-quark



SIDIS TSAs: Collins effect and Transversity




$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_S)} \sin(\phi_h + \phi_S) + \dots \right\}$$

$$F_{UT}^{\sin(\phi_h + \phi_S)} = C \left[-\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$



- Measured on P/D in SIDIS and in dihadron SIDIS
- Compatible results COMPASS/HERMES (Q^2 is different by a factor of ~ 2 -3)
- **No impact from Q^2 -evolution?**
- Extensive phenomenological studies and various global fits by different groups


Total protons delivered on the production target: $\sim 5.95 \times 10^{18}$ (98% of the request) in ~ 150 days



SPS efficiency: $\sim 73\%$
 Spectrometer efficiency: $\sim 90\%$
 Physics data collection efficiency: $\sim 75\%$

Highly successful Run in 2022!

SIDIS TSAs: Collins effect and Transversity




$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_S)} \sin(\phi_h + \phi_S) + \dots \right\}$$

$$F_{UT}^{\sin(\phi_h + \phi_S)} = C \left[-\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$




- Measured on P/D in SIDIS and dihadron SIDIS
- Extensive phenomenological studies and various global fits by different groups
- New deuteron data crucial to constrain d -quark transversity



COMPASS 2022 run – highly successful data-taking!

- 2nd COMPASS deuteron measurements conducted in 2022: unique SIDIS data for the next decades

SIDIS TSAs: Collins effect and Transversity




$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_S)} \sin(\phi_h + \phi_S) + \dots \right\}$$

$$F_{UT}^{\sin(\phi_h + \phi_S)} = C \left[-\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$




- Measured on P/D in SIDIS and dihadron SIDIS
- Extensive phenomenological studies and various global fits by different groups
- New deuteron data crucial to constrain d -quark transversity



COMPASS 2022 run – highly successful data-taking!

- 2nd COMPASS deuteron measurements conducted in 2022: unique SIDIS data for the next decades

Dihadron Collins effect and Transversity





$$\frac{d^7 \sigma}{d \cos \theta d M_{hh} d \phi_R d z d x d y d \phi_S} =$$

$$\frac{\alpha^2}{2\pi Q^2 y} \left((1-y + \frac{y^2}{2}) \sum_q e_q^2 f_1^q(x) D_{1,q}(z, M_{hh}^2, \cos \theta) + \right.$$

$$\left. S_\perp (1-y) \sum_q e_q^2 \frac{|\mathbf{p}_1 - \mathbf{p}_2|}{2M_{hh}} \sin \theta \sin \phi_{RS} h_1^q(x) H_{1,q}^\triangleleft(z, M_{hh}^2, \cos \theta) \right)$$

$$A_{UT}^{\sin \phi_{RS}} = \frac{|\mathbf{p}_1 - \mathbf{p}_2|}{2M_{hh}} \frac{\sum_q e_q^2 h_1^q(x) H_{1,q}^\triangleleft(z, M_{hh}^2, \cos \theta)}{\sum_q e_q^2 f_1^q(x) D_{1,q}(z, M_{hh}^2, \cos \theta)}$$




COMPASS 2022 run – highly successful data-taking!

- 2nd COMPASS deuteron measurements conducted in 2022: unique SIDIS data for the next decades
- **New results – dihedron Collins-like asymmetries**
- Access to collinear transversity PDF; Non-zero trend at large x
- Precision comparable with proton results

SIDIS TSAs: Sivers effect



$$\frac{d\sigma}{dxdydzdp_T^2d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T A_{UT}^{\sin(\phi_h - \phi_S)} \sin(\phi_h - \phi_S) + \dots \right\}$$







$$F_{UT,T}^{\sin(\phi_h - \phi_S)} = C \left[-\frac{\hat{\mathbf{h}} \cdot \mathbf{k}_T}{M} f_{1T}^{\perp q} D_{1q}^h \right], F_{UT,L}^{\sin(\phi_h - \phi_S)} = 0$$






- COMPASS-HERMES discrepancy
- T-oddness: sign-change (SIDIS \leftrightarrow Drell-Yan)
 - Explored by COMPASS
- New precise deuteron data from COMPASS
 - Unique input to constrain Sivers PDF




COMPASS 2022 run: new unique deuteron data





Drell-Yan



SIDIS


SIDIS and single-polarized DY x-sections at twist-2 (LO)



$$\frac{d\sigma^{LO}}{dx dy dz dp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L})$$

SIDIS

$$\times \left\{ \begin{array}{l} 1 + \varepsilon A_{UU}^{\cos 2\phi_h} \cos 2\phi_h \\ + S_L \varepsilon A_{UL}^{\sin 2\phi_h} \sin 2\phi_h + S_L \lambda \sqrt{1-\varepsilon^2} A_{LL} \\ \times \left[\begin{array}{l} A_{UT}^{\sin(\phi_h - \phi_s)} \sin(\phi_h - \phi_s) \\ + \varepsilon A_{UT}^{\sin(\phi_h + \phi_s)} \sin(\phi_h + \phi_s) \\ + \varepsilon A_{UT}^{\sin(3\phi_h - \phi_s)} \sin(3\phi_h - \phi_s) \end{array} \right] \\ + S_T \lambda \left[\sqrt{(1-\varepsilon^2)} A_{LT}^{\cos(\phi_h - \phi_s)} \cos(\phi_h - \phi_s) \right] \end{array} \right\}$$




$$\frac{d\sigma^{LO}}{dq^4 d\Omega} \propto F_U^1 (1 + \cos^2 \theta_{CS})$$


DY

$$\times \left\{ \begin{array}{l} 1 + D_{[\sin^2 \theta_{CS}]} A_U^{\cos 2\phi_{CS}} \cos 2\phi_{CS} \\ + S_L \sin^2 \theta_{CS} A_L^{\sin 2\phi_{CS}} \sin 2\phi_{CS} \\ \times \left[\begin{array}{l} A_T^{\sin \phi_s} \sin \phi_s \\ + D_{[\sin^2 \theta_{CS}]} \left(\begin{array}{l} A_T^{\sin(2\phi_{CS} - \phi_s)} \sin(2\phi_{CS} - \phi_s) \\ + A_T^{\sin(2\phi_{CS} + \phi_s)} \sin(2\phi_{CS} + \phi_s) \end{array} \right) \end{array} \right] \end{array} \right\}$$

$$\text{where } D_{[\sin^2 \theta_{CS}]} = \sin^2 \theta_{CS} / (1 + \cos^2 \theta_{CS})$$



Single-polarized Drell-Yan cross-section at twist-2 (LO)



$$\frac{d\sigma^{LO}}{dq^4 d\Omega} \propto F_U^1 \left(1 + \cos^2 \theta_{CS}\right)$$




$$\times \left\{ \begin{array}{l} 1 + D_{[\sin^2 \theta_{CS}]} A_U^{\cos 2\varphi_{CS}} \cos 2\varphi_{CS} \\ + S_L \sin^2 \theta_{CS} A_L^{\sin 2\varphi_{CS}} \sin 2\varphi_{CS} \\ + \boxed{S_T \left[\begin{array}{l} A_T^{\sin \varphi_S} \sin \varphi_S \\ + D_{[\sin^2 \theta_{CS}]} \left(\begin{array}{l} A_T^{\sin(2\varphi_{CS} - \varphi_S)} \sin(2\varphi_{CS} - \varphi_S) \\ + A_T^{\sin(2\varphi_{CS} + \varphi_S)} \sin(2\varphi_{CS} + \varphi_S) \end{array} \right) \end{array} \right]} \end{array} \right\}$$

$A_U^{\cos 2\varphi_{CS}} \propto \underline{h}_{1,\pi}^{\perp q} \otimes \underline{h}_{1,p}^{\perp q}$	Boer-Mulders (T-odd)
$A_T^{\sin \varphi_S} \propto f_{1,\pi}^q \otimes \underline{f}_{1T,p}^{\perp q}$	Sivers (T-odd)
$A_T^{\sin(2\varphi_{CS} - \varphi_S)} \propto \underline{h}_{1,\pi}^{\perp q} \otimes \underline{h}_{1,p}^q$	Transversity
$A_T^{\sin(2\varphi_{CS} + \varphi_S)} \propto \underline{h}_{1,\pi}^{\perp q} \otimes \underline{h}_{1T,p}^{\perp q}$	Pretzelosity
SIDIS \leftrightarrow Drell-Yan sign-change of the T-odd TMD PDFs	

COMPASS phase-II proposal submitted in 2010 (Drell-Yan, DVCS,...)

Predictions for a large Sivers effect in Drell-Yan and J/psi at COMPASS \rightarrow sign change test

Single-polarized Drell-Yan cross-section at twist-2 (LO)



$$\frac{d\sigma^{LO}}{dq^4 d\Omega} \propto F_U^1 (1 + \cos^2 \theta_{CS})$$

$$\times \left\{ \begin{array}{l} 1 + D_{[\sin^2 \theta_{CS}]} A_U^{\cos 2\varphi_{CS}} \cos 2\varphi_{CS} \\ + S_L \sin^2 \theta_{CS} A_L^{\sin 2\varphi_{CS}} \sin 2\varphi_{CS} \\ + \boxed{S_T \left[\begin{array}{l} A_T^{\sin \varphi_S} \sin \varphi_S \\ + D_{[\sin^2 \theta_{CS}]} \left(A_T^{\sin(2\varphi_{CS} - \varphi_S)} \sin(2\varphi_{CS} - \varphi_S) \right. \right. \\ \left. \left. + A_T^{\sin(2\varphi_{CS} + \varphi_S)} \sin(2\varphi_{CS} + \varphi_S) \right) \end{array} \right]} \end{array} \right\}$$

$A_U^{\cos 2\varphi_{CS}} \propto h_{1,\pi}^{\perp q} \otimes h_{1,p}^{\perp q}$	Boer-Mulders (T-odd)
$A_T^{\sin \varphi_S} \propto f_{1,\pi}^q \otimes f_{1T,p}^{\perp q}$	Sivers (T-odd)
$A_T^{\sin(2\varphi_{CS} - \varphi_S)} \propto h_{1,\pi}^{\perp q} \otimes h_{1,p}^q$	Transversity
$A_T^{\sin(2\varphi_{CS} + \varphi_S)} \propto h_{1,\pi}^{\perp q} \otimes h_{1T,p}^{\perp q}$	Pretzelosity

SIDIS \leftrightarrow Drell-Yan sign-change of the T-odd TMD PDFs

COMPASS phase-II proposal submitted in 2010 (Drell-Yan, DVCS,...)




Predictions for a large Sivers effect in Drell-Yan and J/psi at COMPASS \rightarrow sign change test

Drell-Yan TSAs – Sivers effect

$$\frac{d\sigma}{dq^4 d\Omega} \propto 1 + \dots + S_T [A_T^{\sin \varphi_S} \sin \varphi_S + \dots]$$

Sivers DY TSA

$$A_T^{\sin \varphi_S} \propto f_{1,\pi}^q \otimes f_{1T,p}^{\perp q}$$




Sivers SIDIS TSA

$$A_{UT}^{\sin(\phi_h - \phi_s)} \propto f_{1T}^{\perp q} \otimes D_{1q}^h$$

COMPASS proton Sivers measurements

- Clear signal in the matching Q² ranges






Drell-Yan TSAs – Sivers effect

$$\frac{d\sigma}{dq^4 d\Omega} \propto 1 + \dots + S_T [A_T^{\sin \varphi_S} \sin \varphi_S + \dots]$$

Sivers DY TSA

$$A_T^{\sin \varphi_S} \propto f_{1,\pi}^q \otimes f_{1T,p}^{\perp q}$$




Sivers SIDIS TSA

$$A_{UT}^{\sin(\phi_h - \phi_s)} \propto f_{1T}^{\perp q} \otimes D_{1q}^h$$

COMPASS proton Sivers measurements

- Clear signal in the matching Q^2 ranges






Drell-Yan TSAs – Transversity

$$\frac{d\sigma}{dq^4 d\Omega} \propto 1 + \dots + S_T \left[D_{[\sin^2 \theta_{CS}]} A_T^{\sin(2\phi_{CS} - \phi_s)} \sin(2\phi_{CS} - \phi_s) + \dots \right]$$



Transversity DY TSA

$$A_T^{\sin(2\phi_{CS} - \phi_s)} \propto h_{1,\pi}^{\perp q} \otimes h_{1,p}^q$$




Collins SIDIS TSA

$$A_{UT}^{\sin(\phi_h + \phi_s)} \propto h_1^q \otimes H_{1q}^{\perp h}$$


COMPASS proton Collins measurements

- Clear signal in the matching Q^2 ranges



Transverse-spin asymmetries in $\pi^- p^\uparrow$ scattering

Theory curves based on S. Bastami et al. JHEP 02, (2021), 166



Drell-Yan measurements


- Ruled out predictions for large asymmetries
- General agreement with currently available model calculations
- COMPASS data favors the sign-change hypothesis for the Sivers TMD PDF**
- COMPASS data also favors proton Boer-Mulders TMD PDF sign-change (indirect, model-based)**

J/ψ production channel

- All TSAs are small and compatible with zero
- Hint that J/ψ production might go via gluon-gluon fusion in COMPASS**
- Access to small gluon TMDs?

Conclusions

- Importance of careful understanding and confrontation of experimental data from different experiments
 - Different kinematic domains and phase-space limitations
 - Experiments employ complex analysis techniques, Monte-Carlo simulations, and sophisticated corrections (acceptance, VMs, radiative corrections)
- Close collaboration between different experiments → general benefit for the field
 - Knowledge transfer, comparison of the analysis techniques, tools, and methodology, cross-analyses between different experiments
- Close collaboration between experiment and phenomenology/theory
 - Flexibility in adapting on the analysis side to the choice of the observables, phase-space selections, etc. (before publishing the data)
 - Different possibilities for common paper projects, external membership
- Possibility to organize effective and fruitful collaborative work





The 8th edition of COMAP mini-workshop dedicated to synergies between COMPASS, LHCspin and AMBER projects

22 May 2024, CERN, Switzerland
for registration/info see:
<https://indico.cern.ch/e/COMAP-LHCspin-AMBER>

Organizers: Bakur Parsamyan, Pasquale Di Nezza,
Fulvio Tessarotto, Jan Matousek, Luciano Libero Pappalardo,
Marco Santimaria, Thomas Poschl

Welcome and Introduction	Dr Bakur Parsamyan et al.
892/1-D20, CERN	09:00 - 09:05
The LHCspin proposal	Pasquale Di Nezza
892/1-D20, CERN	09:05 - 09:30
Spin asymmetries for quarkonium production as a probe of gluon distributions	Cristian Pisano
892/1-D20, CERN	09:30 - 09:55
Spin-dependent gluon TMDs	Dr Francesco Giovanni Celiberto
892/1-D20, CERN	09:55 - 10:20
Coffee Break	
892/1-D20, CERN	10:20 - 10:40
Pion-induced Drell-Yan and J/psi production measurements of COMPASS	Catarina Quintans
892/1-D20, CERN	10:40 - 11:05
Quarkonium Polarization Measurements: Challenges and Opportunities	Ilse Kraetschmer
892/1-D20, CERN	11:05 - 11:30
LHCspin UPC physcis opportunities	
892/1-D20, CERN	11:30 - 11:55
The physics case of LHCspin	Luciano Libero Pappalardo
892/1-D20, CERN	14:00 - 14:25
LHCspin simulations	Marco Santimaria
892/1-D20, CERN	14:25 - 14:50
Pion and kaon PDFs confronted by fixed-target charmonium production	Jen-Chieh Peng
892/1-D20, CERN	14:50 - 15:15
Drell-Yan and J/psi measurements programm at AMBER	
892/1-D20, CERN	15:15 - 15:40
Round table	
892/1-D20, CERN	15:40 - 16:20



Yerevan
Armenia



Joint XX-th International Workshop on Hadron Structure and Spectroscopy and 5-th Workshop on Correlations in Partonic and Hadronic Interactions



Yerevan, Armenia

30 September – 4 October, 2024

<https://indico.cern.ch/e/IWHSS-CPHI-2024>



Confirmed speakers

Abhay Deshpande

Albi Kerbizi

Alessandro Bacchetta

Alessandro Pilloni

Alexander Ilyichev

Alexey Prokudin

Alexey Vladimirov

Asmita Mukherjee

Audrey Francisco

Charlotte Van Hulse

Cristian Pisano

Dennis Sivers

Eric Voutier

Gregory Matousek

Giulio Mezzadri

Gunar Schnell

Igor Denisenko

Ishara Fernando

Jen-Chieh Peng

Jinlong Zhang

Lamiaa El Fassi

Latifa Elouadrhiri

Leonard Gamberg

Liliet Diaz

Marco Radici

Misak Sargsian

Nobuo Sato

Oleg Eyser

Pasquale Di Nezza

Patrizia Rossi

Paweł Sznajder

Shohini Bhattacharya

Silvia Niccolai

Stefan Diehl

Stephane Peigné

Holly S zumila-Vance

Timothy Hayward

Valery Kubarovskiy

Valerio Bertone

Xuan Tong

Whitney Armstrong

Xiao-Rui Lyu

Yuri Kovchegov

Zein-Eddine Meziani



Conclusions

- COMPASS holds the record for the longest-running CERN experiment
(20 years of data-taking)
- Series of successful and important measurements addressing nucleon spin-structure
 - Inclusive measurements, unpolarized and polarized SIDIS (longitudinal/transverse)
 - First-ever polarized Drell-Yan measurements
- A wealth of (SI)DIS, Drell-Yan, DVCS, HEMP data collected across the years
 - **Petabytes of data available for analysis**
- Wide and unique kinematic domain accessing low x and large Q^2
 - **Will remain unique for at least another decade**
- World-unique SIDIS deuteron data collected in 2022
 - **Highly successful run, promising first results – soon in PRL**
- Since 2023 the experiment entered the Analysis Phase
 - The spectrometer has been transferred to the COMPASS successor in the M2 beamline – the AMBER collaboration
 - **3 new groups joined COMPASS in the course of 2023 for the Analysis Phase**
 - **If you are interested – don't hesitate to get in touch!**

Thank You!



- Spare slides

COMPASS collaboration



Common Muon and Proton Apparatus for Structure and Spectroscopy



25 institutions from 13 countries
– nearly 200 physicists (in 2022)

- CERN SPS North Area
- Fixed target experiment
- Approved in 1997 (**25 years**)
- Taking data since 2002 (**20 years**)


International Workshop on Hadron Structure and Spectroscopy
IWHSS-2022 workshop (**anniversary edition**)

CERN Globe, August 29-31, 2022



<https://indico.cern.ch/e/IWHSS-2022>

B. Parsamyan



COMPASS experimental setup: Phase II (DVCS programme)




COmmon Muon Proton Apparatus for Structure and Spectroscopy



CERN SPS North Area (building 888)

Two-stage spectrometer LAS+SAS


- Large Angle Spectrometer (SM1 magnet)
- Small Angle Spectrometer (SM2 magnet)



- Primary beam - 400 GeV p from SPS
 - impinging on Be production target (T6)
- 190 GeV secondary hadron beams
 - h^- beam: 97% π^- , 2% K^- , 1% p
 - h^+ beam: 75% p , 24% π^+ , 1% K^+
- 160 GeV tertiary muon beams
 - μ^\pm longitudinally polarized



- Unpolarized target
- Liquid H_2
- Recoil detector "CAMERA"








COMPASS data taking campaigns

Beam	Target	year	Physics programme
μ^+	Polarized deuteron (${}^6\text{LiD}$)	2002	
		2003	80% Longitudinal 20% Transverse SIDIS
		2004	
		2006	Longitudinal SIDIS
	Polarized proton (NH_3)	2007	50% Longitudinal 50% Transverse SIDIS
$\pi^- \text{K}^- \text{p}$	LH_2 , Ni, Pb, W	2008 2009	Spectroscopy
μ^+	Polarized proton (NH_3)	2010	Transverse SIDIS
		2011	Longitudinal SIDIS
$\pi^- \text{K}^- \text{p}$	Ni	2012	Primakoff
μ^\pm	LH_2	2012	Pilot DVCS & HEMP & unpolarized SIDIS
π^-	Polarized proton (NH_3)	2014	Pilot Drell-Yan
		2015	
		2018	Transverse Drell-Yan
μ^\pm	LH_2	2016	
		2017	DVCS & HEMP & unpolarized SIDIS
μ^+	Polarized deuteron (${}^6\text{LiD}$)	2021 2022	Transverse SIDIS


COMPASS Legacy





FLAVOUR PHYSICS | NEWS

COMPASS points to triangle singularity


23 August 2021



The COMPASS experiment at CERN has reported the first direct evidence on decays which can masquerade as a re



COMPASS proposal



The logo for COMPASS, featuring the word "COMPASS" in a bold, italicized serif font inside a blue circle with a red compass needle pointing towards the top right.

COMPASS
approval

COMPASS 1st data taking

SIDIS L/T

Spectroscopy
SIDIS

L/T
Primakoff
DVCS(pilot)

AMBER
1st data taking

Pilot run

Phase I

B. Parsamyan

Phase II

Analysis Phase

Nucleon spin structure: collinear approach \leftrightarrow TMDs





quark			
	U	L	T
U	$f_1^q(x)$ number density		
L		$g_1^q(x)$ helicity	
T			$h_1^q(x)$ transversity




quark			
	U	L	T
U	$f_1^q(x, \mathbf{k}_T^2)$ number density		$h_1^{\perp q}(x, \mathbf{k}_T^2)$ Boer-Mulders
L		$g_1^q(x, \mathbf{k}_T^2)$ helicity	$h_{1L}^{\perp q}(x, \mathbf{k}_T^2)$ worm-gear L
T	$f_{1T}^{\perp q}(x, \mathbf{k}_T^2)$ Sivers	$g_{1T}^q(x, \mathbf{k}_T^2)$ worm-gear T	$h_1^q(x, \mathbf{k}_T^2)$ transversity $h_{1T}^{\perp q}(x, \mathbf{k}_T^2)$ pretzelosity


- PDFs – universal (process independent) objects; T-odd PDFs – conditionally universal




Nucleon spin structure: helicity $g_{1,d}^q(x)$




- COMPASS contribution:
lowest x and highest Q^2 regions




COMPASS PLB 769(2017) 34




Nucleon spin structure: helicity $g_{1,p}^q(x)$




- COMPASS contribution:
lowest x and highest Q^2 regions
- Both deuteron and proton target data





COMPASS PLB 753(2016)18




Nucleon spin structure: helicity $g_{1,p}^q(x)$




- COMPASS contribution:
lowest x and highest Q^2 regions
- Both deuteron and proton target data
- For the first time non-zero spin effects at
smallest x and Q^2 – positive signal for $g_1^p(x)$




Nucleon spin structure: helicity $g_{1,d(p)}^q(x)$




- COMPASS contribution:
lowest x and highest Q^2 regions
- Both deuteron and proton target data
- For the first time non-zero spin effects at
smallest x and Q^2 – positive signal for $g_1^p(x)$
- Both inclusive and semi-inclusive
measurements – access to flavor



COMPASS PLB 680 (2009) 217



COMPASS PLB 693 (2010) 227




SIDIS: target longitudinal spin dependent asymmetries



$$\frac{d\sigma}{dx dy dz dp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_L \sqrt{2\varepsilon(1+\varepsilon)} A_{UL}^{\sin \phi_h} \sin \phi_h + \dots \right\}$$

$$F_{UL}^{\sin \phi_h} = \frac{2M}{Q} \mathcal{C} \left\{ -\frac{\hat{\mathbf{h}} \cdot \mathbf{p}_T}{M_h} \left(x h_L^q H_{1q}^{\perp h} + \frac{M_h}{M} g_{1L}^q \frac{\tilde{G}_q^{\perp h}}{z} \right) \right. \\ \left. + \frac{\hat{\mathbf{h}} \cdot \mathbf{k}_T}{M} \left(x f_L^{\perp q} D_{1q}^h - \frac{M_h}{M} h_{1L}^{\perp q} \frac{\tilde{H}_q^h}{z} \right) \right\}$$



- Q-suppression, various different “twist” ingredients
- **Measured to be non zero COMPASS and HERMES**
- **Non-zero trend for h^+ , h^- - compatible with zero, clear z -dependence**


SIDIS: target longitudinal spin dependent asymmetries



$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_L \sqrt{2\varepsilon(1+\varepsilon)} A_{UL}^{\sin\phi_h} \sin\phi_h + \dots \right\}$$


$$F_{UL}^{\sin\phi_h} = \frac{2M}{Q} \mathcal{C} \left\{ -\frac{\hat{\mathbf{h}} \cdot \mathbf{p}_T}{M_h} \left(x h_L^q H_{1q}^{\perp h} + \frac{M_h}{M} g_{1L}^q \frac{\tilde{G}_q^{\perp h}}{z} \right) + \frac{\hat{\mathbf{h}} \cdot \mathbf{k}_T}{M} \left(x f_L^{\perp q} D_{1q}^h - \frac{M_h}{M} h_{1L}^{\perp q} \frac{\tilde{H}_q^h}{z} \right) \right\}$$

S. Bastami et al. JHEP 1906 (2019) 007:
“SIDIS in Wandzura-Wilczek-type approximation”




- Q-suppression, various different “twist” ingredients
- Measured to be non zero COMPASS and HERMES
- Non-zero trend for h^+ , h^- - compatible with zero, clear z -dependence

B. Parsamyan (for COMPASS)
[arXiv:1801.01488 \[hep-ex\]](https://arxiv.org/abs/1801.01488)




Zhun Lu
Phys. Rev. D 90, 014037(2014)




Selected results for di-hadron LSAs

COMPASS (NH₃) 2007+2011 data: preliminary




- Alternative way to access various twist-2/-3 distributions
- Non zero signal for $A_{UL}^{\sin\phi_R}$ and A_{LL}^1
- CLAS-COMPASS: different behavior for $A_{UL}^{\sin2\phi_R}$ at large x?


CLAS 6 GeV (NH₃)
S. A. Pereira: PoS (DIS 2014) 231





$Q^2 > 1 (\text{GeV}/c)^2$
 $0.0025 < x < 0.7$
 $0.1 < y < 0.9$
 $W > 5 \text{ GeV}/c^2$




Nucleon spin structure: helicity $g_{1,d(p)}^q(x)$



- COMPASS contribution:
lowest x and highest Q^2 regions
- Both deuteron and proton target data
- For the first time non-zero spin effects at
smallest x and Q^2 – positive signal for $g_1^p(x)$
- Gluon polarization measurements
via open charm and SIDIS
- COMPASS - first to rule out a large gluon
polarization in the nucleon!
Precise test of Bjorken sum rule (9% level)



SIDIS TSAs: Collins effect and Transversity





$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_S)} \sin(\phi_h + \phi_S) + \dots \right\}$$

$$F_{UT}^{\sin(\phi_h + \phi_S)} = C \left[-\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$





- Measured on P/D in SIDIS and in dihadron SIDIS
- Compatible results COMPASS/HERMES (Q^2 is different by a factor of ~ 2 -3)
- No impact from Q^2 -evolution?


PLB 824 (2022) 136834



- indication for a positive asymmetry
- opposite to π^+ and π^0 as predicted by the models
- Large effect at small P_T



SIDIS TSAs: Collins effect and Transversity





$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_S)} \sin(\phi_h + \phi_S) + \dots \right\}$$

$$F_{UT}^{\sin(\phi_h + \phi_S)} = C \left[-\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$



- Measured on P/D in SIDIS and dihadron SIDIS
- Extensive phenomenological studies and various global fits by different groups
- New deuteron data crucial to constrain d -quark transversity




COMPASS 2022 run – highly successful data-taking!

- 2nd COMPASS deuteron measurements conducted in 2022: unique SIDIS data for the next decades

SIDIS TSAs: Sivers effect




$$\frac{d\sigma}{dxdydzdp_T^2d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T A_{UT}^{\sin(\phi_h - \phi_S)} \sin(\phi_h - \phi_S) + \dots \right\}$$




$$F_{UT,T}^{\sin(\phi_h - \phi_S)} = C \left[-\frac{\hat{\mathbf{h}} \cdot \mathbf{k}_T}{M} f_{1T}^{\perp q} D_{1q}^h \right], F_{UT,L}^{\sin(\phi_h - \phi_S)} = 0$$



- COMPASS-HERMES discrepancy
- T-oddness: sign-change (SIDIS \leftrightarrow Drell-Yan)
 - Explored by COMPASS
- New precise deuteron data from COMPASS
 - Unique input to constrain Sivers PDF



SIDIS TSAs: Kotzinian-Mulders asymmetry



$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + \lambda S_T \sqrt{(1-\varepsilon^2)} A_{LT}^{\cos(\phi_h - \phi_S)} \cos(\phi_h - \phi_S) + \dots \right\}$$

$$F_{LT}^{\cos(\phi_h - \phi_S)} = C \left[\frac{\hat{\mathbf{h}} \cdot \mathbf{k}_T}{M} g_{1T}^q D_{1q}^h \right]$$




COMPASS/HERMES/CLAS6 results


$$A_{LT}^{\cos(\phi_h - \phi_S)}$$

- Only “twist-2” ingredients
- Sizable non-zero effect for h^+ !
- Similar effect at HERMES



COMPASS, PBL 770 (2017) 138; PoS QCDEV2017 (2018) 042



See also, PRD 107, (2023) 034016 – global fit by:
M. Horstmann, A. Schafer and A. Vladimirov






First global QCD analysis of the g_{1T} TMD PDF using SIDIS data



Nucleon 3D structure: GPDs

- Transverse position \vec{b}_T of partons
 - Correlation between \vec{b}_T and x
 - Complementary to TMD PDFs
- 8 generalized parton distribution functions (GPDs)
 - Contain information about parton orbital angular momentum
 - Mostly unknown
- COMPASS exclusive process measurements:
 - Deeply virtual Compton scattering (DVCS): $\mu + N \rightarrow \mu + \gamma + N$
 - Hard exclusive meson production (HEMP): $\mu + N \rightarrow \mu + \text{Meson} + N$ with $\pi^0, \rho(770), \omega(782), \dots$





SIDIS: target longitudinal spin dependent asymmetries



$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_L \lambda \sqrt{2\varepsilon(1-\varepsilon)} A_{LL}^{\cos\phi_h} \cos\phi_h + \dots \right\}$$


$$F_{LL}^{\cos\phi_h} = \frac{2M}{Q} C \left\{ -\frac{\hat{\mathbf{h}} \cdot \mathbf{p}_T}{M_h} \left(xe_L^q H_{1q}^{\perp h} + \frac{M_h}{M} g_{1L}^q \frac{\tilde{D}_q^{\perp h}}{z} \right) \right. \\ \left. + \frac{\hat{\mathbf{h}} \cdot \mathbf{k}_T}{M} \left(x g_L^{\perp q} D_{1q}^h - \frac{M_h}{M} h_{1L}^{\perp q} \frac{\tilde{E}_q^h}{z} \right) \right\}$$





- Q-suppression, various different “twist” ingredients
- **Measured to be non zero at CLAS6, what about CLAS12?**
- HERMES/COMPASS - small and compatible with zero, in agreement with model predictions



Transverse-spin asymmetries in $\pi^- p^\uparrow$ scattering




Transverse-spin asymmetries in $\pi^- p^\uparrow$ scattering




$$\text{DY: } A_T^{\sin(2\varphi_{CS} - \varphi_S)} \propto -\left\{ h_{1,\pi^-}^{\perp\bar{u}} \otimes h_{1,p}^u \right\} < 0 \Rightarrow h_{1,\pi^-}^{\perp\bar{u}} > 0$$


M. Lambertsen, W. Vogelsang
PRD93, 114013 (2016)



$$A_U^{\cos 2\varphi_{CS}} \propto h_{1,\pi}^{\perp q} \otimes h_{1,p}^{\perp q}$$



S. Bastami et al. JHEP 02, (2021), 166



$$\text{DY: } A_T^{\sin 2\varphi_{CS}} \propto \left\{ h_{1,\pi^-}^{\perp\bar{u}} \otimes h_{1,p}^{\perp u} \right\} > 0 \Rightarrow h_{1,p}^{\perp u} > 0 \stackrel{\text{sign-change}}{\Leftrightarrow} \text{SIDIS: } h_{1,p}^{\perp u} < 0$$

$h_{1,p}^{\perp u} < 0 \rightarrow \text{SIDIS fits}$
V. Barone, et al.
PRD 82 (2010) 114025

- COMPASS data also favors proton Boer-Mulders TMD PDF sign-change